Contents lists available at ScienceDirect

Infection, Genetics and Evolution

journal homepage: www.elsevier.com/locate/meegid



Research paper

Identification and characterization of new broad host-range rV5-like coliphages C203 and P206 directed against enterobacteria



Domonkos Sváb^a, Linda Falgenhauer^b, Manfred Rohde^c, Trinad Chakraborty^b, István Tóth^{a,*}

^a Institute for Veterinary Medical Research, Centre for Agricultural Research, Hungarian Academy of Sciences, Budapest, Hungary

^b Institute of Medical Microbiology, Justus Liebig University Giessen, German Center for Infection Research (DZIF), Partner site Giessen-Marburg-Langen, Giessen, Germany

^c Central Facility for Microscopy, Helmholtz Centre for Infection Research, HZI, Braunschweig, Germany

ARTICLE INFO

Keywords: Bacteriophage rV5-like phage Whole phage genome Phylogeny Host specificity E. coli 0157 Food sample

ABSTRACT

We isolated and characterized two novel rV5-like lytic bacteriophages from independently collected food samples. Nucleotide sequence analysis revealed that these phages have linear double-stranded DNA genomes comprising 138,073 bp with 213 CDS and 5 tRNA genes. The two genomes contain completely identical nucleotide sequence, albeit there is a 10,718 bp-long shift in the sequence. The GC content of the phage genomes was 43.7% and they showed high general homology to rV5-like phages. The new phages were termed C203 and P206. The genome of both phages contains a unique ORF that encodes for a putative phage homing endonuclease. The phage produced clear plaques with a burst size of approx. 1000 viral particles and a latent period of 60 min. Morphological investigation indicated that the new phages are members of the family Myoviridae with an approximate head length of 85 nm, tail length of 75 nm, and a head width of 96 nm. C203 and P206 exhibit a broad and uniform host range, which included enterohemorrhagic Escherichia coli strains of serogroup O157, multi drug resistant (MDR) E. coli strains of various sero- and pathotypes, and both Shigella sonnei and S. dysenteriae strains. C203 and P206 both effectively reduced the number of living EHEC 0157:H7 Sakai in experimentally inoculated minced meat. The same broad host range, the lack of any virulence related genes, the stability and its short latent period suggest that these newly found phages could be suitable candidates as a bio-control agents against food-borne pathogenic Enterobacteria.

1. Introduction

With the increasing resistance against antibiotics, there is now renewed interest in bacteriophages (shortly phages) capable of lysing important pathogenic bacteria (reviewed by Hagens and Loessner, 2010). Enterohemorrhagic Escherichia coli (EHEC) strains of the O157:H7 serotype are considered to be among the most dangerous foodborne pathogens (reviewed in (Croxen et al., 2013; Gyles, 2007). They can cause serious hemorrhagic colitis (HC), in some cases with the life-threatening complication of hemolytic-uremic syndrome (HUS) and thrombocytopenia (reviewed in (Bielaszewska and Karch, 2005; Croxen et al., 2013). Because of these features, E. coli O157:H7 strains have been prime targets in studies aiming to identify bacteriophages capable of eradicating them from its host or from food products. Several groups have isolated and characterized bacteriophages capable of lysis or in vitro growth inhibition of E. coli O157:H7 type strains. These include both T5- and T4-like phages (Lee et al., 2016; Liu et al., 2015; Raya et al., 2011).

rV5-like phages (V5virus genus) are a recently established genus of tailed bacteriophages belonging to the family Myoviridae with a large genome of over 100 kb long (Kropinski et al., 2013; Santos et al., 2011) and exhibiting a relatively wide host spectrum (Kropinski et al., 2013). Several members of the group are notable for their capacity to lyse E. coli O157:H7 strains (Kropinski et al., 2013; Truncaite et al., 2012). In the study which suggested the establishment of the rV5 genus, the authors highlighted the lack of restriction sites, the usage of inner core lipopolysaccharide receptors and the absence of lysogeny-associated genes that contribute to the phages' broad host range and make phage PVP-SE1 good candidate for biocontrol against Salmonella (Santos et al., 2011).

In an earlier study, we assessed the risk presented by food-borne pathogens present in foodstuff illegally imported into Europe, with a special emphasis on Shiga-toxin producing E. coli (STEC; Nagy et al., 2015). We hypothesized that phages capable of lysing these pathogens may also be present in the same foodstuff.

In the current study, we characterized two new rV5-like phages

* Corresponding author.

E-mail addresses: svab.domonkos@agrar.mta.hu (D. Sváb), toth.istvan@agrar.mta.hu (I. Tóth).

https://doi.org/10.1016/j.meegid.2018.07.004

Received 27 April 2018; Received in revised form 8 June 2018; Accepted 3 July 2018 Available online 04 July 2018

1567-1348/ © 2018 Elsevier B.V. All rights reserved.

designated C203 and P206 originating from foodstuff. The phages isolated are founding members of a new genotype, and apart from several *E. coli* O157:H7 strains, they are capable of lysing an unusually wide spectrum of pathogenic *E. coli* from other serotypes as well.

2. Materials and methods

2.1. Bacteriophage isolation

Bacteriophages were isolated from two independent sources, from cottage cheese and from poultry liver confiscated on the Hungarian border, the samples of which were designated C203 and P206, respectively. The samples underwent the first steps of the ISO 16654:2001 method for isolating *E. coli* O157. Briefly, 5 g pieces of the food samples were homogenized at 1:10 weight to volume ratio of tryptic soy broth supplemented with bile salts, and incubated for 24 h at 42 °C. After removing the bacteria by centrifugation the samples' supernatants were spread or spotted onto layered agar plates containing *E. coli* K-12 derivative strains C600 and MG1655. After overnight incubation at 37 °C, single plaques were picked up and purified by amplification on *E. coli* MG1655 at least three times, until high titer (at least 10^{11} PFU/ml) phage stocks were produced.

2.2. Bacterial strains

E. coli K-12 derivative strain MG1655 was mostly used for propagation of the phages. The efficiency of plating (EOP) was tested on various enterobacterial strains listed in Tables 1 and 2. A rifampicin-resistant mutant of the EHEC O157:H7 Sakai strain was used in the *in situ* bacterial challenge test (2.8).

2.3. Phage DNA isolation

Phage DNA was isolated from phage stocks with a concentration of at least 10^{11} PFU/ml. The phenol-chloroform method described by

Sambrook et al. (1987) was used for DNA isolation, with the modifications outlined as described before (Tóth et al., 2016).

2.4. Genome sequence determination and analysis

Genomic DNA sequencing libraries were prepared using the Nextera XT kit (Illumina, Eindhoven, NL). Sequencing was performed using Nextseq Mid-output reagent kit v2 (2×150 bp) on an Illumina NextSeq 500. Assembly was performed with CLC Genomic Workbench 9.0. The genome was annotated using the RAST server (Overbeek et al., 2014). Homology searches were conducted with the BLAST tools available at the NCBI website, with PSI-BLAST results supplementing the annotation.

The genome sequences of phage C203 and P206 were deposited in GenBank under the accession nos. MG022439 and MG022440, respectively.

2.5. Phylogenetic analysis

Whole-genome based phylogenetic analysis was conducted with VICTOR (Meier-Kolthoff and Goeker, 2017). A progressive Mauve alignment (Darling et al., 2011) including bacteriophage genomes C203, P206, FFH2, FV3, Murica, slur16 and rV5 (GenBank nos. MG022440, LN881727.1, KJ190158, NC_019517, KT001917, DQ832317) was also conducted.

2.6. Host specificity and efficiency of plating

Host specificity and EOP was tested on a wide array of pathogenic *E. coli, Salmonella* and *Shigella* strains, with an emphasis on *E. coli* O157 strains. Among *E. coli* pathotypes, EHEC, enteropathogenic (EPEC), atypical *E. coli* O157 and multidrug resistant (MDR) strains of human origin were also included. *E. coli* O157:H7 strain C83/00 representing phage type 55 was kindly provided by Ivelina Damjanova (National Institute of Hygiene, Budapest). The serotype, pathotype and other

Table 1

Host specificity and efficiency of plating of bacteriophages C203 and P206. EOP values are given relative to the titer of the phages on *E. coli* MG1655. Zero (0) values mean that no lysis was observable on the given strain.

| Strain | Pathotype/serovar/species | Serogroup or serotype | EOP | Strain reference |
|----------|---------------------------|-----------------------|----------------------|------------------------------|
| MG1655 | E. coli K-12 | O16:H48 | 1 | (Blattner et al., 1997) |
| Sakai | EHEC | O157:H7 | 7×10^{-1} | (Hayashi et al., 2001) |
| EDL933 | EHEC | O157:H7 | 2×10^{-5} | (Perna et al., 2001) |
| E22 | EHEC | O103:H2 | 0 | (Marchès et al., 2003) |
| E2348/69 | EPEC | O127:H6 | 1.3×10^{-2} | (Iguchi et al., 2009) |
| II95–36 | EIEC | 0121 | 2×10^{-8} | (Sváb et al., 2018) |
| 20 | EIEC | 0124 | 4×10^{-1} | (Sváb et al., 2018) |
| Bra2 26 | EIEC | 0152 | 2×10^{-7} | (Sváb et al., 2018) |
| Saigon | EIEC | O164 | 8×10^{-8} | (Sváb et al., 2018) |
| T22 | atypical | O157:H43 | 2×10^{-6} | (Tóth et al., 2009b) |
| 536 | UPEC | O6:K15:H31 | 0 | (Hochhut et al., 2006) |
| IHE3034 | ExPEC | O18:K1:H7 | 0 | (Moriel et al., 2010) |
| E250 | APEC | 0115 | 2×10^{-6} | (Tóth et al., 2009a) |
| 5871 | MDR E. coli | 015 | 2×10^{-6} | (Sváb et al., 2018) |
| 18531 | MDR E. coli | 073 | 0 | (Sváb et al., 2018) |
| 29095 | MDR E. coli | O90 | 2×10^{-9} | (Sváb et al., 2018) |
| ILC169 | Citrobacter rodentium | N/A | 0 | (Petty et al., 2011) |
| 20080 | Shigella dysenteriae 1A | N/A | 10^{-1} | (Sváb et al., 2018) |
| M90 T | Shigella flexneri | N/A | 10^{-2} | (Sváb et al., 2018) |
| 20038 | Shigella boydii | N/A | 0 | (Sváb et al., 2018) |
| 866-F | Shigella sonnei | N/A | 4×10^{-7} | (Allué-Guardia et al., 2011) |
| 20045 | Shigella sonnei | N/A | 0 | (Sváb et al., 2018) |
| 75/02 | Shigella sonnei | N/A | $3.3 	imes 10^{-8}$ | (Sváb et al., 2017) |
| 1201 | Salmonella Typhimurium 1 | N/A | 0 | (Sváb et al., 2018) |
| 1202 | Salmonella Infantis | N/A | 0 | (Sváb et al., 2018) |
| 1203 | Salmonella Panama | N/A | 0 | (Sváb et al., 2018) |
| 1199 | Salmonella Typhi | N/A | 0 | (Sváb et al., 2018) |
| 1198 | Salmonella Gallinarum | N/A | 0 | (Sváb et al., 2018) |
| 1200 | Salmonella Enteritidis | N/A | 0 | (Sváb et al., 2018) |
| | | | | |

Table 2

Lysis pattern and efficiency of plating of bacteriophage C203 and P206 on *E. coli* strains of the O157 serogroup, with *E. coli* MG1655 as reference efficiency. As in the case of Table 1, the lysis pattern of C203 and P206 was completely identical. Exclusive lysis by C203/P206 is marked with gray.

| Strain | Pathotpye | Serotype | Phage type ¹ | EOP ² | Lysed by typing phage ⁴ | | Reference | |
|--------|-----------------|----------|-------------------------|------------------|------------------------------------|---------|-----------|-------------------------|
| | | | | | TP4 | TP5 | TP14 | |
| MG1655 | E. coli K-12 | O16:H48 | ND | 1 | | | | (Blattner et al., 1997) |
| Sakai | EHEC | O157:H7 | PT14 | $> 10^{-4}$ | + | + | + | (Hayashi et al., 2001) |
| EDL933 | EHEC | O157:H7 | PT21 | $> 10^{-4}$ | - | - | - | (Perna et al., 2001) |
| 34 | EHEC | O157:H7 | PT21 | 0 | - | - | - | (Tóth et al., 2009b) |
| 52 | EHEC | O157:H7 | PT33 | 0 | - | - | - | (Tóth et al., 2009b) |
| 318 | EHEC | O157:NM | PT8 | $> 10^{-5}$ | + | + | + | (Tóth et al., 2009b) |
| 65 | EPEC | O157:H7 | PT33 | 0 | - | - | - | (Tóth et al., 2009b) |
| 68 | EPEC | O157:H7 | PT8 | $> 10^{-3}$ | + | + | + | (Tóth et al., 2009b) |
| 103 | EPEC | O157:H7 | PT50d | $> 10^{-5}$ | - | - | - | (Tóth et al., 2009b) |
| 144 | EPEC | O157:H7 | NT-R | 0 | - | - | - | (Tóth et al., 2009b) |
| B20 | AT ³ | O157:H12 | NT-R | $> 10^{-5}$ | - | - | - | (Tóth et al., 2009b) |
| B47 | AT ³ | O157:NM | NT-R | 0 | - | - | - | (Tóth et al., 2009b) |
| T22 | AT ³ | O157:H43 | NC | $> 10^{-5}$ | No data | No data | No data | (Tóth et al., 2009b) |
| T34 | AT ³ | O157:H9 | PT21 | $> 10^{-5}$ | - | - | - | (Tóth et al., 2009b) |
| T49 | AT ³ | O157:H37 | NC | $> 10^{-5}$ | No data | No data | No data | (Tóth et al., 2009b) |
| C83/00 | EHEC | O157:H7 | PT55 | $> 10^{-4}$ | + | + | + | present study |

¹ NT: non-typeable, R: phage resistant, NC: non-characteristic phage type; ND: not done.

² Efficiency of plating.

³ Atypical pathotype (*stx* and *eae* negative);

 4 + = lysis; - = no lysis.

characteristics of these bacterial strains are given in Tables 1 and 2.

Host spectrum and EOP was determined by applying serial dilutions of the phage suspension in spot assays on layered soft agar containing cultures of the test strains, according to the protocol described by Strauch et al. (2001). The ratio of phage titer on the test strain divided by the titer measured on *E. coli* MG1655 was considered the EOP of the phage on the given strain.

2.7. One step growth experiments

One step growth experiment to determine burst size was performed on *E. coli* MG1655 according the protocol of Lee et al. (2016) with some modifications. Briefly, 2×10^8 bacteria were mixed with 2×10^6 phages, setting multiplicity of infection (MOI) to 0.01 in Luria-Bertani broth (LB), and incubated for 20 min at room temperature. After incubation the mixture was centrifuged at 6000 x g for 10 min, the pellet was resuspended in 50 ml of fresh LB, and incubated at 37 °C for 1 h. Samples were taken every 5 min, and plated on layered soft agar for counting. Three independent experiments were run in two parallels. Burst size was determined as a ratio of the phage count before and after the burst.

2.8. In vitro and in situ bacterial challenge tests

In order to assess the activity of the phages against EHEC 0157 strains, a bacterial challenge test was performed. *E. coli* 0157:H7 Sakai strain was grown overnight in liquid LB culture by shaking with 180 rpm at 37 °C, then diluted 100-fold and grown again to $OD_{600} = 1$. Culture was diluted 10-fold and aliquoted into 48 separate cultures. Twenty-four of the cultures received P206 in MOI = 10 and all cultures were grown again at 37 °C by shaking with 180 rpm for 12 h. At 1-h intervals the OD_{600} of two cultures each of the treated and non-treated sets was measured.

In order to isolate rifampicin resistant mutants of EHEC O157:H7 Sakai strain, two hundred microliters of overnight LB culture of *E. coli* O157:H7 Sakai strain was spread onto rifampicin (150 µg/ml) containing LB agar plate and incubated at 37 °C for 16 h. Single colonies were selected and further purified by several passages on selective plates. One of the resistant colonies was used for further experiments.

In situ biocontrol potential of P206 was investigated in the following

way: One gram of minced meat, bought at a local supermarket was homogenized by using 1 ml LB and inoculated with high number (2.5×10^{10}) of the rifampicin resistant derivative of *E. coli* O157:H7 Sakai cells. After inoculation, 1.5×10^{10} and 1.5×10^{9} phage particles of P206 were applied to the mixture, which corresponded to multiplicity of infection (MOI) 0.6 and 0.06, respectively. Control tubes with bacterial cells alone were also used. The experiment was conducted in triplicate. After two hours incubation at 37 °C samples were homogenized and serial dilutions were plated out on LB plates containing 150 µg/ml of rifampicin in duplicate. Number of surviving bacterial colonies was determined after 16 h incubation at 37 °C.

2.9. Determination of phage morphology

Bacteriophages were investigated by transmission electron microscopy (TEM). Briefly, drops of high titer bacteriophage suspensions were placed on parafilm, absorbed onto carbon film, washed in TE buffer (10 mM TRIS, 1 mM EDTA, pH 6.9) and negatively-stained with 2% aqueous uranyl acetate, pH 5.0. Carbon film was collected with 300 mesh copper grids and excess negative-stain was removed with filter paper and subsequently air-dried. Samples were examined in a TEM 910 transmission electron microscope (Carl Zeiss, Oberkochen) at an acceleration voltage of 80 kV. Images were recorded digitally at calibrated magnifications with a Slow-Scan CCD-Camera (ProScan, 1024 × 1024, Scheuring, Germany) with ITEM-Software (Olympus Soft Imaging Solutions, Münster, Germany). Contrast and brightness were adjusted with Adobe Photoshop CS3.

2.10. Heat stability tests

Heat stability of phages was tested as follows: 1 ml of LB containing 10^9 PFU/ml phage was incubated for 1 h at 25, 37, 42 and 80 °C, respectively. After incubation, the titer of the treated stocks was determined with a spot assay on *E. coli* MG1655. Experiments were performed in two parallels on each temperature.

2.11. pH tolerance tests

Stability of the phages under different pH values was tested by incubating 10⁹ PFU/ml phages for 1 h at 37 °C in 1 ml LB stocks pH-



Fig. 1. Transmission electronmicrograph of negatively-stained (2% uranyl acetate) bacteriophages C203 (A and B) and P206 (C and D). Both phages exhibit the typical Myoviridae morphology of non-bended and contractile tails. Bars represent 100 nm.

adjusted to 3, 5, 7, 9 and 10 with HCl and NaOH solutions. After incubation, the titer of the treated stocks was determined with spot assay on *E. coli* MG1655. Experiments were performed in two parallels on each pH value.

3. Results

3.1. Morphology

Two bacteriophages were isolated from two independent food sources by using *E. coli* K12 derivatives as propagating strains. Both phages showed Myoviridae morphology, with head length around 85 nm and width around 75 nm. The characteristic contractile tail was around 96 nm long (Fig. 1).

3.2. Genome characteristics

The new bacteriophages were designated as C203 and P206 (with C and P referring to the cheese and poultry sources). The genome sequences of the phages, as determined on the Illumina platform, are 138,073 bp in length, and contain 213 ORFs and 5 tRNA genes. The GC content of the genomes is 43.7%. Average read length was 183 for phage C203 and 177 for P206, the average coverage was 1173.97 and 1453.33, respectively. Homology searches showed that genome sequence of the phages are identical, but the sequence start and endpoints are different. This causes a 10,718 bp-long shift between the two phages, therefore ORFs 1–7 in the C203 genome correspond to ORFs 207–213 in the P206 genome. Therefore ORF1 in P206 is a truncated version of ORF8 in C203, the rest of the gene was transferred to the end

of the genome in P206 as non-coding region. Similarly, ORF1 in C203 is a truncated version of ORF207 in P206. The RAST and PSI-BLAST homology searches enabled to assign a function to 36% (78/213) of the ORFs, with the rest encoding proteins with unknown functions. The complete list of ORFs for both phages is given in Supplementary Tables 1 and 2.

The overall genome structure of C203 and P206 regarding the order of genes is similar to the rV5-like bacteriophage genomes sequenced earlier, with sequence shifts as indicated by the MAUVE alignment (Fig. 2). On the nucleotide level according to whole-genome BLAST search it showed the highest similarity to Escherichia phage vB_EcoM_FFH2 (GenBank KJ190158.1) with 97% coverage and 97% average sequence identity, albeit some genes from C203/P206 rather showed significant homology to those carried by other rV5-like phages, these are indicated in Supplementary Tables 1 and 2. A large region comprising of tail fiber protein CDSs AHN83624.1 and AHN83629.1 in vB_EcoM_FFH2, is absent in C203/P206. Individual sequence homology searches showed that ORF12 and 5 of C203 and P206 respectively, are highly homologous to the tail fiber genes carried by the Salmonella specific phages slur12 and slur16, and the phage EcoM_FV3 (GenBank nos. LN881735.1, LN881727.1, and NC_019517). Additionally, ORF16/ 9 in C203/P206 is homologous to the corresponding gene of the original rv5 phage (CDS ABI79112.1) and those of three O157-typing phages (TP4, TP 5 and TP14). Further differences between C203/P206 and vB_EcoM_FFH2 are the absence of genes encoding a putative NTP reductase small subunit (AHN83629.1) together with two putative HNH endonucleases. One of the endonuclease genes (AHN83797.1 in phage EcoM_FFH2) disrupts the phage terminase large subunit, which is intact in C203/P206 (ORF39/32), like in the case of the Murica phage



Fig. 2. Progressive Mauve alignment of rV5-like bacteriophage genomes C203, P206, slur16, FFH2, FV3, Murica and rV5 (GenBank nos. MG022439, MG022440, LN881727.1, KJ190158, NC_019517, KT001917, DQ832317). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

(KT001917). The other endonuclease in EcoM_FFH2 (AHN83797.1) is situated between a DNA polymerase and a DNA primase (corresponding ORFs 181–182 in C203 and 174–175 in P206) similarly to the original rV5-like phage (GenBank DQ832317.1) as well as to EcoM_FV3. Table 3 summarizes the relations of ORFs annotated as tail fiber genes, comparing them on the amino acid level to other rV5-like phages.

3.3. Phylogenetic relations

Phylogenetic tree of rV5-like phages including bacteriophage C203 and P206 was constructed based with the VICTOR whole genome phylogenetic analysis It showed that throughout the rV-like phages which were included, whole genome sequences had a pairwise identity of at least 90%. The resulting tree suggests the slur12 and slur16 phages as closest relatives to the new phages (Fig. 3.).

3.4. Host spectrum, efficiency of plating, burst size and stability

Phages C203 and P206 propagated on a wide array of commensal

and pathogenic E. coli strains, most notably on the prototype strains of EHEC O157:H7 Sakai and EDL933. The lysis pattern of C203 and P206 was completely identical. Additionally to E. coli strains, phages C203 and P206 also propagated on four out of six Shigella strains representing three Shigella species, while at the same time none of six Salmonella strains belonging to different serovars was lysed by these phages. The full host spectrum of phages C203 and P206 together with EOP values is listed in Table 1. In the one step growth experiments the burst size proved to be around 1000 with a latent period of approximately 60 min (Fig. 4). The lysis pattern of C203/P206 on various O157:H7 strains paired with phage types is shown separately in Table 2. Note that the plaque morphology of the phages was cloudy on the O157 strains, hence only an approximate minimum EOP value is given, as only the endpoint could be determined without exactly countable plaque numbers. Differences between the six CDSs identified as tail fibers (Supplementary Tables 1 and 2) could account for the different lysis pattern between C203/P206 and the O157 typing phages. All these CDSs had at least one amino acid switch when compared to the corresponding CDSs of the typing phages, and in one case, ORF12/6 in C203/P206 shows

Table 3

Amino acid sequence based comparison of tail fiber encoding ORFs in rV5-like phages.

| Phage name | GenBank no. | Amino acid changes compared to tail fiber encoding ORFs of phage C203 | | | | | | |
|----------------------|-------------|---|------|------|------------------|-------------------|-------------------|-------|
| | | ORF4 | ORF5 | ORF7 | ORF8 | ORF12 | ORF16 | ORF17 |
| Murica | KT001917.1 | 7 | 2 | 2 | 1 | 105 | Only 99% coverage | 5 |
| V18 | KY683736.1 | 3 | 5 | 0 | 4 + 1 frameshift | Only 78% coverage | Only 78% coverage | 11 |
| APCEc02 | KR698074.1 | 7 + a 5 AA-long gap | 3 | 3 | 0 | 99 | Only 84% coverage | 7 |
| O157 typing phage 14 | KP869112.1 | 6 | 2 | 2 | 3 | 65 | Only 98% coverage | 7 |
| slur12 | LN881735.1 | 3 | 0 | 2 | 2 | 14 | 25 | 9 |
| slur16 | LN881727.1 | 3 | 0 | 2 | 2 | 14 | 25 | 9 |
| O157 typing phage 5 | KP869103.1 | 4 | 4 | 2 | 1 | 16 | Only 98% coverage | 7 |
| rV5 | DQ832317.1 | 5 | 4 | 2 | 1 | 16 | Only 98% coverage | 7 |
| O157 typing phage 4 | KP869102.1 | 1 | 6 | 1 | 1 | 20 | Only 84% coverage | 8 |
| 2 JES-2013 | KC690136.1 | 6 | 9 | 2 | 0 | 103 | Only 98% coverage | 8 |
| vB_EcoM_FFH2 | KJ190158.1 | 6 | 3 | 2 | 1 | 99 | Only 84% coverage | 6 |
| vB_EcoM-FV3 | JQ031132.1 | 8 | 15 | 1 | 3 | 26 | 56 | 26 |



H 0.005

Fig. 3. Whole-genome based phylogenetic relations of rV5-like bacteriophages made with VICTOR. The bacteriophages and their accession numbers are: Murica, KT001917.1; V18, KY683736.1; APCEc02, KR698074.1; O157 typing phage 14, KP869112.1; slur12, LN881735.1; slur16, LN881727.1; O157 typing phage 5, KP869103.1; rV5, DQ832317.1; O157 typing phage 4, KP869102.1; 2 JES-2013, KC690136.1; vB_EcoM_FFH2, KJ190158.1; vB_EcoM-FV3, JQ031132.1.

only 78% identity to the corresponding CDS in typing phage 14 on the amino acid level.

A stock of P206 propagated on the *E. coli* O157:H7 Sakai strain was re-plated on all bacterial strains which proved to be sensitive to the phage earlier. All of them were still sensitive with roughly the same EOP values, but in the case of the EIEC O152 strain Bra2 26, a 10^4 -fold EOP increase could be observed, while in the case of bovine EHEC O157:H7 strain 318 a 10^3 -fold EOP decrease occurred.

Heat stability experiments showed that incubation at 25, 37 and 42 °C did not affect the EOP of the phages. Incubation at 80 °C, however, caused 10^4 -fold average decrease in the phage titer. Throughout the experiments in general, the phage proved to be stable after weeks of incubation at 4 °C.

At pH value of 3 the phages were completely inactivated, as no plaques were observable even in the spots of the concentrated suspension. On the other hand, no significant decrease in titer was observed at pH values 5, 7, 9 and 10.

3.5. In vitro and in situ bacterial challenges

In a MOI of 10, phage P206 could effectively inhibit the growth of *E. coli* O157:H7 Sakai in LB for 5 h, and the treated samples showed lagged growth even 12 h following phage addition (Fig. 5).

Since O157 contaminated meat is the leading source of human enterohemorrhagic infections, we aimed to explore the elimination potential of phage C203/P206 in minced meat experimentally contaminated with a rifampicin resistant derivative of *E. coli* O157:H7 Sakai strain. The *in situ* bacterial challenge experiments showed that phage P206 effectively reduced the extremely high number (2.5×10^{10}) of EHEC O157:H7 cells in the contaminated meat samples. By using phages in MOI 0.6 and 0.06, after 2 h of infection, the number of surviving EHEC cells was reduced by 10^4 and 10^2 times, respectively. The unincoulated meat did not contain rifampicin resistant bacteria (data not shown).

4. Discussion

In the current study we isolated and characterized two rV5-like bacteriophages designated C203 and P206 from independent food samples, carrying identical genome content. Interestingly the phage genomes have a 10,718 bp-long shift in their sequences. These phages are capable of lysing a broad spectrum of pathogenic *E. coli* strains, including most notably the reference EHEC O157:H7 strains (Hayashi et al., 2001; Perna et al., 2001). It is worth mentioning that while the phages lysed numerous *E. coli* O157 strains, strains of this serogroup were not detected in the original food samples from which they were isolated (Nagy et al., 2015). C203 and P206 effectively lysed a rifampicin resistant derivative of EHEC O157 Sakai strain *in situ*.

In the recent years, rV5-like phages have been established as a 'singleton' genus within the Myoviridae family, showing similar genomic organization and phenotypic characteristics to each other, but exhibiting significant differences to other coliphages (Smith et al., 2015; Truncaite et al., 2012). Based on their morphology, all of them are members of the Myoviridae family, and phage C203 and P206 are no exception from this notion.

The genomes of C203 and P206 have identical gene content except for truncated genes at the genome ends. Their closest homologue based on BLAST searches is phage vB_EcoM_FFH2, but there are striking differences in the tail fiber genes, and genes encoding DNA modifying enzymes present in vB_EcoM_FFH2, are absent from C203/P206. These genes, as shown by BLAST searches, are rather similar to those found in the *Salmonella* phages slur12 and slur16 (Smith et al., 2015), the phage EcoM_FV3 (Truncaite et al., 2012) and Murica (Wilder et al., 2015). It has to be noted that the Mauve alignment of rV5-like genomes indicated that the order of genes is generally uniform in rV5 phages, with sequence shifts responsible for different organizations (Fig. 2).

It is also noteworthy that the genetic relationship suggested by



Fig. 4. One step growth curve of phage P206 on E. coli K-12 MG1655.



Infection, Genetics and Evolution 64 (2018) 254-261

Fig. 5. Bacterial challenge assay of bacteriophage P206 against E. coli O157:H7 Sakai strain. A dilution of overnight bacterial culture was infected with phage P206 at a MOI of 10. Non-phage treated bacterial culture was used as a control. The graphs show OD at 600 nm for the cultures every hour. The error bars indicate standard deviations from the results of the experiment.

BLAST, with vB EcoM FFH2 showing closest general homology on the nucleotide level to C203/P206, was not reflected by the whole-genome based phylogenetic analysis, which rather depicted a closer relationship of C203/P206 to slur16 and slur12 (Fig. 3).

The host specificity of rV5-like phages is wide, but C203/P206 showed the broadest lysis spectrum of those reported. EOP values of strains other than the K-12 derivative strain MG1655 were generally orders of magnitude lower (Table 1). Strains lysed by C203/P206 include several major pathotypes of intestinal pathogenic E. coli besides EHEC O157:H7, such as enteroinvasive (EIEC), avian pathogenic (APEC) as well as strains with atypical or unknown pathotype, with MDR strains among the latter.

The rV5-like phages also include three of the typing phages used in the phage typing of E. coli O157:H7 strains (Ahmed et al., 1987; Cowley et al., 2015) namely TP 4, 5 and 14, whose whole genomes are also available (Cowley et al., 2015). For comparative reasons, several E. coli O157:H7 strains (Tóth et al., 2009) of various phage types (PT) were also included in testing the host spectrum of C203/P206, and it was found that their lysis pattern was different from those of the typing phages, viz., they produced lysis on two out of three PT21 strains and one PT50d strain (Table 2). Differences in the protein sequence of C203/P206 tail fibers and those of TPs could be responsible for the discrepancies. These differences are summarized in Table 3, with the ORFs of phage C203 as reference. It is noteworthy that ORF5 is identical to the corresponding ORFs of slur12 and slur16 phages, ORF7 to that carried by ApCEC02, and ORF8 to those in ApCEC02 and 2 JES-2013. On the other hand, the longer tail fiber ORFs, 12 and 16 show significant variability and in many cases only partial coverage. These data suggest that the host spectrum of rV5-like phages could be variable, but currently our study is the most comprehensive in this regard, as we aimed to include most of the significant pathotypes and species of Enterobacterial pathogens.

The fact that C203 and P206, besides E. coli, were able to lyse Shigella dysenteriae and S. sonnei strains deserves attention. To our knowledge, no rV5-like bacteriophage has been reported to infect Shigella strains, which, especially S. dysenteriae even today remain the leading agents of bacillary dysentery (Anderson et al., 2016). As MDR Shigella strains emerge frequently (Doyle, 2015), any potential alternative antibacterial agent that could be used against them is an important finding. Noteworthy is the finding that bacteriophage P206 effectively reduced the number of living EHEC O157:H7 Sakai in experimentally inoculated minced meat. Throughout the experiments, we did not observe the appearance of potentially lysogenic colonies. The annotation did not show the presence of any integrase or recombinase genes, suggesting that phages C203 and P206 are incapable of lysogeny.

Schwarzer et al. (2012) characterized an rV5-like phage with a similarly broad host range, capable of infecting several Salmonella serovars besides E. coli K-12 derivatives. They suggested that multiple tail fiber genes could be responsible for the wide host spectrum of phi92 (Schwarzer et al., 2012), which could also be the case with C203/P206,

as the genome harbors several ORFs annotated as tail fibers. It has to be noted however, that there is very low nucleotide-level homology between phi92 and C203/P206. It is also worth mentioning that the truncation of ORF1 annotated as "tail fibers" in P206 (ORF8 in C203) seems to have no effect on the host specificity of the phage when compared to C203.

Lee et al. (2016) have proposed that genes associated with host specificity should be cloned into one bacteriophage genome, which then could be used as a universal agent against several related pathogenic bacteria. Broad host spectrum bacteriophages such as C203/P206 could be a potential source of such host specificity associated genes, or alternatively, they could be used in their native form as biocontrol agent either as a stand-alone or in phage cocktails.

In conclusion, we isolated and characterized two new members of rV5-like bacteriophages, which showed an unusually broad host spectrum spanning more pathotypes of pathogenic E. coli, Salmonella and Shigella than any other similar phage isolated before. To our knowledge, this is also the first rV5-like phage to be isolated from foodstuff. These characteristics, together with its stability, strictly lytic lifestyle and the fact that they do not carry any pathogenicity-associated genes, makes C203 and P206 suitable candidates to be tested as a biocontrol agent against EHEC O157 strains and other foodborne pathogenic enterobacteria.

Supplementary data to this article can be found online at https:// doi.org/10.1016/j.meegid.2018.07.004.

Acknowledgments

The study was supported by the Hungarian National Research, Development and Innovative Office (NKFIH, grant 124335), 7th EU Framework Programme PROMISE project (project n. 265877) and by grants to the German Center of Infection Research (DZIF, site Giessen-Marburg-Langen), through the German Federal Ministry of Education and Research, grant # TI06.001 and 8032808811 to TC. We thank Christina Gerstmann for excellent technical assistance. Domonkos Sváb is holder of a Bolyai János Stipend from the Hungarian Academy of Sciences

Conflicts of interest

The authors declare no conflict of interest.

References

Ahmed, R., Bopp, C., Borczyk, A., Kasatiya, S., 1987. Phage-typing scheme for Escherichia coli O157:H7. J. Infect. Dis. 155, 806-809.

Allué-Guardia, A., García-Aljaro, C., Muniesa, M., 2011. Bacteriophage-encoding cytolethal distending toxin type V gene induced from nonclinical Escherichia coli isolates. Infect. Immun. 79, 3262-3272. https://doi.org/10.1128/IAI.05071-11.

Anderson, M., Sansonetti, P.J., Marteyn, B.S., 2016. Shigella diversity and changing landscape: insights for the twenty-first century. Front. Cell. Infect. Microbiol. 6 (45). https://doi.org/10.3389/fcimb.2016.00045

Bielaszewska, M., Karch, H., 2005. Consequences of enterohaemorrhagic Escherichia coli infection for the vascular endothelium. Thromb. Haemost. 94, 312–318. https://doi. org/10.1160/TH05-04-0265.

- Blattner, F.R., Plunkett, G., Bloch, C.A., Perna, N.T., Burland, V., Riley, M., Collado-Vides, J., Glasner, J.D., Rode, C.K., Mayhew, G.F., Gregor, J., Davis, N.W., Kirkpatrick, H.A., Goeden, M.A., Rose, D.J., Mau, B., Shao, Y., 1997. The complete genome sequence of *Escherichia coli* K-12. Science 277, 1453–1462.
- Cowley, L.A., Beckett, S.J., Chase-Topping, M., Perry, N., Dallman, T.J., Gally, D.L., Jenkins, C., 2015. Analysis of whole genome sequencing for the *Escherichia coli* 0157:H7 typing phages. BMC Genomics 16 (271). https://doi.org/10.1186/s12864-015-1470-z.
- Croxen, M.A., Law, R.J., Scholz, R., Keeney, K.M., Wlodarska, M., Finlay, B.B., 2013. Recent advances in understanding enteric pathogenic *Escherichia coli*. Clin. Microbiol. Rev. 26, 822–880. https://doi.org/10.1128/CMR.00022-13.
- Darling, A.E., Tritt, A., Eisen, J.A., Facciotti, M.T., 2011. Mauve assembly metrics. Bioinforma. Oxf. Engl. 27, 2756–2757. https://doi.org/10.1093/bioinformatics/ btr451.
- Doyle, M.E., 2015. Multidrug-resistant pathogens in the food supply. Foodborne Pathog. Dis. 12, 261–279. https://doi.org/10.1089/fpd.2014.1865.
- Gyles, C.L., 2007. Shiga toxin-producing Escherichia coli: an overview. J. Anim. Sci. 85, E45–E62. https://doi.org/10.2527/jas.2006-508.
- Hagens, S., Loessner, M.J., 2010. Bacteriophage for biocontrol of foodborne pathogens: calculations and considerations. Curr. Pharm. Biotechnol. 11, 58–68.
- Hayashi, T., Makino, K., Ohnishi, M., Kurokawa, K., Ishii, K., Yokoyama, K., Han, C.G., Ohtsubo, E., Nakayama, K., Murata, T., Tanaka, M., Tobe, T., Iida, T., Takami, H., Honda, T., Sasakawa, C., Ogasawara, N., Yasunaga, T., Kuhara, S., Shiba, T., Hattori, M., Shinagawa, H., 2001. Complete genome sequence of enterohemorrhagic *Escherichia coli* O157:H7 and genomic comparison with a laboratory strain K-12. DNA Res. Int. J. Rapid Publ. Rep. Genes Genomes 8, 11–22.
- Hochhut, B., Wilde, C., Balling, G., Middendorf, B., Dobrindt, U., Brzuszkiewicz, E., Gottschalk, G., Carniel, E., Hacker, J., 2006. Role of pathogenicity island-associated integrases in the genome plasticity of uropathogenic *Escherichia coli* strain 536. Mol. Microbiol. 61, 584–595. https://doi.org/10.1111/j.1365-2958.2006.05255.x.
- Iguchi, A., Thomson, N.R., Ogura, Y., Saunders, D., Ooka, T., Henderson, I.R., Harris, D., Asadulghani, M., Kurokawa, K., Dean, P., Kenny, B., Quail, M.A., Thurston, S., Dougan, G., Hayashi, T., Parkhill, J., Frankel, G., 2009. Complete genome sequence and comparative genome analysis of enteropathogenic *Escherichia coli* 0127:H6 strain E2348/69. J. Bacteriol. 191, 347–354. https://doi.org/10.1128/JB.01238-08.
- Kropinski, A.M., Waddell, T., Meng, J., Franklin, K., Ackermann, H.-W., Ahmed, R., Mazzocco, A., Yates, J., Lingohr, E.J., Johnson, R.P., 2013. The host-range, genomics and proteomics of *Escherichia coli* 0157:H7 bacteriophage rV5. Virol. J. 10, 76. https://doi.org/10.1186/1743-422X-10-76.
- Lee, H., Ku, H.-J., Lee, D.-H., Kim, Y.-T., Shin, H., Ryu, S., Lee, J.-H., 2016. Characterization and genomic study of the novel bacteriophage HY01 infecting both *Escherichia coli* 0157:H7 and *Shigella flexneri*: potential as a biocontrol agent in food. PLoS One 11, e0168985. https://doi.org/10.1371/journal.pone.0168985.
- Liu, H., Niu, Y.D., Meng, R., Wang, J., Li, J., Johnson, R.P., McAllister, T.A., Stanford, K., 2015. Control of *Escherichia coli* O157 on beef at 37, 22 and 4 °C by T5-, T1-, T4-and O1-like bacteriophages. Food Microbiol. 51, 69–73. https://doi.org/10.1016/j.fm. 2015.05.001.
- Marchès, O., Ledger, T.N., Boury, M., Ohara, M., Tu, X., Goffaux, F., Mainil, J., Rosenshine, I., Sugai, M., De Rycke, J., Oswald, E., 2003. Enteropathogenic and enterohaemorrhagic *Escherichia coli* deliver a novel effector called Cif, which blocks cell cycle G2/M transition. Mol. Microbiol. 50, 1553–1567.
- Meier-Kolthoff, J.P., Goeker, M., 2017. VICTOR: genome-based phylogeny and classification of prokaryotic viruses. bioRxiv 107862. https://doi.org/10.1101/107862.
- Moriel, D.G., Bertoldi, I., Spagnuolo, A., Marchi, S., Rosini, R., Nesta, B., Pastorello, I., Corea, V.A.M., Torricelli, G., Cartocci, E., Savino, S., Scarselli, M., Dobrindt, U., Hacker, J., Tettelin, H., Tallon, L.J., Sullivan, S., Wieler, L.H., Ewers, C., Pickard, D., Dougan, G., Fontana, M.R., Rappuoli, R., Pizza, M., Serino, L., 2010. Identification of protective and broadly conserved vaccine antigens from the genome of extraintestinal pathogenic Escherichia coli. Proc. Natl. Acad. Sci. U. S. A. 107, 9072–9077. https:// doi.org/10.1073/pnas.0915077107.
- Nagy, B., Szmolka, A., Smole Možina, S., Kovač, J., Strauss, A., Schlager, S., Beutlich, J., Appel, B., Lušicky, M., Aprikian, P., Pászti, J., Tóth, I., Kugler, R., Wagner, M., 2015. Virulence and antimicrobial resistance determinants of verotoxigenic *Escherichia coli* (VTEC) and of multidrug-resistant *E. coli* from foods of animal origin illegally imported to the EU by flight passengers. Int. J. Food Microbiol. 209, 52–59. https://doi.

org/10.1016/j.ijfoodmicro.2015.06.026.

- Overbeek, R., Olson, R., Pusch, G.D., Olsen, G.J., Davis, J.J., Disz, T., Edwards, R.A., Gerdes, S., Parrello, B., Shukla, M., Vonstein, V., Wattam, A.R., Xia, F., Stevens, R., 2014. The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). Nucleic Acids Res. 42, D206–D214. https://doi.org/10.1093/ nar/gkt1226.
- Perna, N.T., Plunkett, G., Burland, V., Mau, B., Glasner, J.D., Rose, D.J., Mayhew, G.F., Evans, P.S., Gregor, J., Kirkpatrick, H.A., Pósfai, G., Hackett, J., Klink, S., Boutin, A., Shao, Y., Miller, L., Grotbeck, E.J., Davis, N.W., Lim, A., Dimalanta, E.T., Potamousis, K.D., Apodaca, J., Anantharaman, T.S., Lin, J., Yen, G., Schwartz, D.C., Welch, R.A., Blattner, F.R., 2001. Genome sequence of enterohaemorrhagic *Escherichia coli* 0157:H7. Nature 409, 529–533. https://doi.org/10.1038/35054089.
- Petty, N.K., Feltwell, T., Pickard, D., Clare, S., Toribio, A.L., Fookes, M., Roberts, K., Monson, R., Nair, S., Kingsley, R.A., Bulgin, R., Wiles, S., Goulding, D., Keane, T., Corton, C., Lennard, N., Harris, D., Willey, D., Rance, R., Yu, L., Choudhary, J.S., Churcher, C., Quail, M.A., Parkhill, J., Frankel, G., Dougan, G., Salmond, G.P., Thomson, N.R., 2011. *Citrobacter rodentium* is an unstable pathogen showing evidence of significant genomic flux. PLoS Pathog. 4https://doi.org/10.1371/journal. ppat.1002018. e1002018.
- Raya, R.R., Oot, R.A., Moore-Maley, B., Wieland, S., Callaway, T.R., Kutter, E.M., Brabban, A.D., 2011. Naturally resident and exogenously applied T4-like and T5-like bacteriophages can reduce *Escherichia coli* O157:H7 levels in sheep guts. Bacteriophage 1, 15–24. https://doi.org/10.4161/bact.11.14175.

Sambrook, J., Maniatis, T., Fritsch, E.F., Laboratory, C.S.H., 1987. Molecular Cloning: A Laboratory Manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.

- Santos, S.B., Kropinski, A.M., Ceyssens, P.-J., Ackermann, H.-W., Villegas, A., Lavigne, R., Krylov, V.N., Carvalho, C.M., Ferreira, E.C., Azeredo, J., 2011. Genomic and proteomic characterization of the broad-host-range Salmonella phage PVP-SE1: creation of a new phage genus. J. Virol. 85, 11265–11273. https://doi.org/10.1128/JVI. 01769-10.
- Schwarzer, D., Buettner, F.F.R., Browning, C., Nazarov, S., Rabsch, W., Bethe, A., Oberbeck, A., Bowman, V.D., Stummeyer, K., Mühlenhoff, M., Leiman, P.G., Gerardy-Schahn, R., 2012. A multivalent adsorption apparatus explains the broad host range of phage phi92: a comprehensive genomic and structural analysis. J. Virol. 86, 10384–10398. https://doi.org/10.1128/JVI.00801-12.
- Smith, R., O'Hara, M., Hobman, J.L., Millard, A.D., 2015. Draft genome sequences of 14 *Escherichia coli* phages isolated from cattle slurry. Genome Announc. 3. https://doi. org/10.1128/genomeA.01364-15.
- Strauch, E., Lurz, R., Beutin, L., 2001. Characterization of a Shiga toxin-encoding temperate bacteriophage of *Shigella sonnei*. Infect. Immun. 69, 7588–7595. https://doi. org/10.1128/IAI.69.12.7588-7595.2001.
- Sváb, D., Bálint, B., Vásárhelyi, B., Maróti, G., Tóth, I., 2017. Comparative genomic and phylogenetic analysis of a Shiga toxin producing *Shigella sonnei* (STSS) strain. Front. Cell. Infect. Microbiol. 7 (229). https://doi.org/10.3389/fcimb.2017.00229.
- Sváb, D., Falgenhauer, L., Rohde, M., Szabó, J., Chakraborty, T., Tóth, I., 2018. Identification and characterization of T5-like bacteriophages representing two novel subgroups from food products. Front. Microbiol. 9. https://doi.org/10.3389/fmicb. 2018.00202.
- Tóth, I., Nougayrède, J.P., Dobrindt, U., Ledger, T.N., Boury, M., Morabito, S., Fujiwara, T., Sugai, M., Hacker, J., Oswald, E., 2009a. Cytolethal distending toxin type I and type IV genes are framed with lambdoid prophage genes in extraintestinal pathogenic *Escherichia coli*. Infect Immun. 77, 492–500. https://doi.org/10.1128/IAI.00962-08.
- Tóth, I., Schmidt, H., Kardos, G., Lancz, Z., Creuzburg, K., Damjanova, I., Pászti, J., Beutin, L., Nagy, B., 2009b. Virulence genes and molecular typing of different groups of *Escherichia coli* 0157 strains in cattle. Appl. Environ. Microbiol. 75, 6282–6291. https://doi.org/10.1128/AEM.00873-09.
- Tóth, I., Sváb, D., Bálint, B., Brown-Jaque, M., Maróti, G., 2016. Comparative analysis of the Shiga toxin converting bacteriophage first detected in *Shigella sonnei*. Infect. Genet. Evol. J. Mol. Epidemiol. Evol. Genet. Infect. Dis. 37, 150–157. https://doi.org/ 10.1016/j.meegid.2015.11.022.
- Truncaite, L., Šimoliūnas, E., Zajančkauskaite, A., Kaliniene, L., Mankevičiūte, R., Staniulis, J., Klausa, V., Meškys, R., 2012. Bacteriophage vB_EcoM_FV3: a new member of "rV5-like viruses". Arch. Virol. 157, 2431–2435. https://doi.org/10. 1007/s00705-012-1449-x.
- Wilder, J.N., Lancaster, J.C., Cahill, J.L., Rasche, E.S., Kuty Everett, G.F., 2015. Complete genome sequence of enterotoxigenic *Escherichia coli* myophage murica. Genome Announc. 3. https://doi.org/10.1128/genomeA.01135-15.