The Optimal Pebbling Number of Staircase Graphs

Ervin Győri*^{1,2}, Gyula Y. Katona^{†3,4}, László F. Papp^{‡3}, and Casey Tompkins^{§1}

¹Alfréd Rényi Institute of Mathematics, Budapest, Hungary

²Department of Mathematics, Central European University, Budapest, Hungary

³Department of Computer Science and Information Theory, Budapest

University of Technology and Economics, Hungary

⁴MTA-ELTE Numerical Analysis and Large Networks Research Group,

Hungary

October 20, 2018

Abstract

Let G be a graph with a distribution of pebbles on its vertices. A pebbling move consists of removing two pebbles from one vertex and placing one pebble on an adjacent vertex. The optimal pebbling number of G is the smallest number of pebbles which can placed on the vertices of G such that, for any vertex v of G, there is a sequence of pebbling moves resulting in at least one pebble on v. We determine the optimal pebbling number for several classes of induced subgraphs of the square grid, which we call staircase graphs.

1 Introduction

Graph pebbling is a game on graphs introduced by Saks and Lagarias. The main framework is the following: A distribution of pebbles is placed on the vertices of a simple graph. A pebbling move removes two pebbles from a vertex and places one pebble on an adjacent vertex. The goal is to reach any specified vertex by a sequence of pebbling moves. We begin with some notation which we will need to state our results.

Let G be a simple graph. We denote the vertex and edge set of G by V(G) and E(G), respectively. A *pebbling distribution* P is a function from V(G) to the nonnegative integers. We say that G has P(v) pebbles placed at the vertex v under the distribution P. We say that a vertex v is occupied if P(v) > 0 and unoccupied otherwise. The *size* of a pebbling distribution P, denoted |P|, is the total number of pebbles placed on the vertices of G.

Let u be a vertex with at least two pebbles under P, and let v be a neighbor of u. A pebbling move from u to v consists of removing two pebbles from u and adding one pebble to v. That is, a pebbling move yields a new pebbling distribution P' with P'(u) = P(u) - 2 and

^{*}gyori.ervin@renyi.mta.hu

[†]kiskat@cs.bme.hu

[‡]lazsa@cs.bme.hu

[§]ctompkins496@gmail.com




Figure 1: Solvable distribution of the square grid.

P'(v) = P(v) + 1. We say that a vertex v is k-reachable under the distribution P if we can obtain, after a sequence of pebbling moves, a distribution with at least k pebbles on v. If k = 1 we say simply that v is reachable under P. More generally, a set of vertices S is k-reachable under the distribution P if, after a sequence of pebbling moves, we can obtain a distribution with at least a total of k pebbles on the vertices in S.

A pebbling distribution P on G is *solvable* if all vertices of G are reachable under P. Similarly, P is k-solvable if all vertices are k-reachable. A pebbling distribution on G is k-optimal if it is k-solvable and its size is minimal among all of the k-solvable distributions of G; when k=1 we simply say optimal. Note that k-optimal distributions are usually not unique.

The *optimal pebbling number* of G, denoted by $\pi_{\text{opt}}(G)$, is the size of an optimal pebbling distribution. In general, the decision problem for this graph parameter is NP-complete [8].

We denote with P_n and C_n the path and cycle on n vertices, respectively. The *Cartesian product* $G \square H$ of graphs G and H is defined in the following way: $V(G \square H) = V(G) \times V(H)$ and $\{(g_1,h_1),(g_2,h_2)\} \in E(G \square H)$ if and only if $\{g_1,g_2\} \in E(G)$ and $h_1=h_2$ or $\{h_1,h_2\} \in E(H)$ and $g_1=g_2$.

The optimal pebbling number is known for several graphs including paths, cycles [1, 2, 7], caterpillars [3] and m-ary trees [4]. The optimal pebbling number of grids has also been investigated. Exact values were proved for $P_n \square P_2$ [1] and $P_n \square P_3$ [9]. The question for bigger grids is still open. We gave a construction, which is restated in the next paragraph, for big square grids in [5]. Furthermore, we are going to improve the lower bound on the optimal pebbling number of these grids in a forthcoming paper [6].

The distribution P which gives the current best upper bound for the optimal pebbling number of the square grid takes groups of seven consecutive diagonals and places pebbles on the middle one (see Figure 1). Using these pebbles, it is possible to reach any vertex on any diagonal in the group.

We conjecture that P is optimal, however we do not know a proof for this. Can we at least show that the induced distributions on these smaller graphs are optimal? If this was not the case, it would refute the conjecture. These considerations provide the main motivation for the present paper.

We investigate a family of graphs which we call *staircase graphs*. These graphs are connected induced subgraphs of the square grid. The width seven instances correspond to the



Figure 2: Slashes in the three-wide staircase graph.

groups of seven diagonals discussed above.

Let $SG_{n,n} = P_n \square P_n$ and let $SG = P_{\infty} \square P_{\infty}$ be the infinite square grid where P_{∞} is the doubly infinite path with vertex set \mathbb{Z} and edge set $\{\{i, i+1\} : i \in \mathbb{Z}\}$.

For any $k \in \mathbb{Z}$, we say that $D_k^+ = \{\{i,j\} \in V(SG) : i-j=k\}$ is a positive diagonal of SG. Similarly we define the negative diagonal: $D_k^- = \{\{i,j\} \in V(SG) : i+j=k\}$. A staircase graph will be defined in terms of the intersection of a set of consecutive positive diagonals in SG with a set of consecutive negative diagonals. When the number of diagonals taken in each direction is odd, there will be two nonisomorphic graphs to consider.

If m is odd, let $S'_{m,n}$ be the graph induced by the vertex set $\left(\bigcup_{j=1}^m D_j^-\right) \cap \left(\bigcup_{i=1}^n D_i^+\right)$, and let $S_{m,n}$ be the graph induced by $\left(\bigcup_{j=1}^m D_j^-\right) \cap \left(\bigcup_{i=0}^{n-1} D_i^+\right)$.

If m is even, let $S_{m,n}$ be the graph induced by the vertex set $(\bigcup_{j=1}^m D_j^-) \cap (\bigcup_{i=1}^n D_i^+)$. In this case we have only one isomorphism class.

Note that $S'_{m,n} \cong S_{m,n}$ in case when n is even. We also remark that $S_{m,n} \cong S_{n,m}$; nevertheless, we say that the first and the second parameters are the *width* and the *length* of the staircase graph, respectively, and generally assume that $n \geq m$. We will refer to the graphs $S_{m,n}$ and $S'_{m,n}$ as *m-wide staircase graphs*.

For simplicity, we call a nonempty intersection of a staircase graph and a positive diagonal a *slash* (See Figure 1 where each dashed ellipse is a slash).

The optimal pebbling number of ladders is proved by a technique based on induction and cutting. We use this technique for narrow staircase graphs and extend it for wider ones.

2 Results

2.1 Three-wide staircases

The results in the 3 and 4-wide cases depend on the value of n modulo 4. Therefore we write 4k + r instead of n, where $r \in \{0, 1, 2, 3\}$.

Theorem 2.1 When $4k + r \ge 2$, then

$$\pi_{\text{opt}}(S_{3,4k+r}) = 3k + r,$$

$$\pi_{\text{opt}}(S'_{3,4k+r}) = \begin{cases} 3k+2 & \text{if } r = 3\\ 3k+r & \text{otherwise.} \end{cases}$$

We also note that $\pi_{\mathrm{opt}}(S_{3,1})=1$ and $\pi_{\mathrm{opt}}(S_{3,1}')=2$, since these cases may appear in our induction. Before we present the proof we establish some lemmas. Each of them utilizes the one preceding it, and the first one strongly relies on the following theorem from [1].

Theorem 2.2 ([1]) A 2-optimal distribution of the n-vertex path contains n+1 pebbles.




Figure 3: Optimal distributions of $S_{3,n}$ and $S'_{3,n}$ when $n \geq 5$.

Lemma 2.3 Let G be $S_{m,n}$ or $S'_{m,n}$ and suppose there is a solvable distribution P on G such that |P| < n + 1. Then there is a slash in G which is not 2-reachable under P.

We are going to use the *collapsing technique* introduced in [1]. Let G and H be simple graphs. We say that H is a *quotient* of G, if there is a surjective mapping $\phi:V(G)\to V(H)$ such that $\{h_1,h_2\}\in E(H)$ if and only if there are $g_1,g_2\in V(G)$, where $\{g_1,g_2\}\in E(G)$, $h_1=\phi(g_1)$ and $h_2=\phi(g_2)$. We say that ϕ collapses G to H, and if P is a pebbling distribution on G, then the *collapsed distribution* P_{ϕ} on H is defined in the following way: $P_{\phi}(h)=\sum_{g\in V(G)|\phi(g)=h}P(g)$.

PROOF: Let ϕ be a mapping which maps each slash in G to a vertex in P_n in such a way that consecutive slashes are mapped to adjacent vertices. It is easy to see that if a slash was 2-reachable under P, then its collapsed image is also 2-reachable under P_{ϕ} . P and P_{ϕ} have less than n+1 pebbles. Therefore, Theorem 2.2 yields that P_{ϕ} is not 2-solvable, therefore there is a vertex in P_n which is not 2-reachable under P_{ϕ} and the corresponding slash is not 2-reachable under P. \square

Lemma 2.4 Let G be $S_{m,n}$ or $S'_{m,n}$ and suppose there is a solvable distribution P on G such that |P| < n - 1. Then there is slash in G which is neither the first nor the last slash and is not 2-reachable under P.

PROOF: If there are only two slashes which are not 2-reachable under P, then we put a pebble on a vertex in both slashes and all slashes are 2-reachable with less than n+1 pebbles. This contradicts Lemma 2.3. Therefore, there are at least three slashes which are not 2-reachable under P. One of them is neither the first nor the last slash. \square

Lemma 2.5 If slash k is not 2-reachable under a distribution P and it is not the first or the last slash, then there is no possible pebbling move between either slash k-1 and k or slash k and k+1.

PROOF: Since slash k is not 2-reachable, it is impossible to make a pebbling move starting at a vertex from slash k. It follows that the pebbling distributions which can be obtained on the slashes numbered less than k do not depend on the pebbling distribution on slashes numbered greater than k and vice versa. If slash k could be reached from a pebbling move from both slashes k-1 and k+1, then these moves could be performed independently and it would follow that slash k is 2-reachable, a contradiction. \square




Figure 4: Optimal distributions of small $S_{3,n}$ and $S'_{3,n}$ graphs.

If there is no possible pebbling move between slashes k and k+1 then, after the deletion of the edges between them, each vertex of the resulting graph is still reachable under P. Hence P induces a solvable distribution on each connected component. We will use this fact during later proofs.

Claim 2.6 Let G be $S_{m,n}$ or $S'_{m,n}$, and assume that there is a solvable distribution on G with less than n-1 pebbles. Then, there is a c such that $1 \le c < n$ and:

$$\pi_{\text{opt}}(S_{m,n}) \geq \pi_{\text{opt}}(S_{m,c}) + \pi_{\text{opt}}(S_{m,n-c}) \text{ if } c \text{ is even}$$

$$\pi_{\text{opt}}(S_{m,n}) \geq \pi_{\text{opt}}(S_{m,c}) + \pi_{\text{opt}}(S'_{m,n-c}) \text{ if } c \text{ is odd}$$

$$\pi_{\text{opt}}(S'_{m,n}) \geq \pi_{\text{opt}}(S'_{m,c}) + \pi_{\text{opt}}(S'_{m,n-c}) \text{ if } c \text{ is even}$$

$$\pi_{\text{opt}}(S'_{m,n}) \geq \pi_{\text{opt}}(S'_{m,c}) + \pi_{\text{opt}}(S_{m,n-c}) \text{ if } c \text{ is odd}.$$

If m *is even, we have only one inequality:*

$$\pi_{\text{opt}}(S_{m,n}) \ge \pi_{\text{opt}}(S_{m,c}) + \pi_{\text{opt}}(S_{m,n-c}).$$

PROOF: An optimal pebbling distribution P has at most n-1 pebbles and so we can apply Lemma 2.4 and Lemma 2.5. Hence there is a slash k, which is neither the first nor the last, such that we cannot move a pebble between slashes k and k+1 or k and k-1. We choose c to be k in the first scenario and k-1 in the second. Then, we delete the edges between these slashes and obtain two smaller graphs with solvable distributions. The sum of the sizes of these two distributions is the same as |P|. The size of a solvable distribution is always at least as big as the optimal pebbling number. This gives us a lower bound. The types of the two resulting graphs depend on the type of the original graph and the parity of c. \square

PROOF OF THEOREM 2.1: The upper bound comes from solvable distributions shown in Figure 3

We prove the lower bounds for $S_{3,n}$ and $S'_{3,n}$ simultaneously by induction on n. The cases where $n \leq 7$ are shown in Figure 4. It can be checked by hand or computer that these are optimal distributions.

Therefore, we will assume that $n \geq 8$ and the lower bound is true for values smaller than n. In this case we can apply Claim 2.6. Unfortunately, we do not know the value of c, hence we have to check all the possible values to prove a lower bound. The formulas which we want to prove and use as an inductive hypothesis are quite complicated and depend on the value of n modulo 4. Therefore, we have several cases. We will see that some of these cases are not possible if P is optimal.

First, consider $S_{3,n}$. We remind the reader that n=4k+r, where $r \in \{0,1,2,3\}$. We cut between the slashes c and c+1, where c=4l+q and $q \in \{0,1,2,3\}$. Similarly, assume that

n-c=4j+p where $p\in\{0,1,2,3\}$. Note that l+j equals k when q+p<4 and k-1 otherwise. Furthermore, the parity of c and q are the same. When q is even:

$$\pi_{\text{opt}}(S_{3,n}) \ge \pi_{\text{opt}}(S_{3,c}) + \pi_{\text{opt}}(S_{3,n-c}) = 3l + q + 3j + p = 3(l+j) + q + p$$

$$= \begin{cases} 3k + r & \text{if } q + p < 4 \\ 3(k-1) + 4 + r & \text{if } q + p \ge 4. \end{cases}$$

Note that the second case in the inequality is not possible because it would give a bigger lower bound on $\pi_{\text{opt}}(S_{3,n})$ than the known upper bound.

When q is odd and $p \neq 3$, we obtain the same estimate on $\pi_{\text{opt}}(S_{3,n})$ except in the case when j=0 and p=1 where the right-hand side is larger by 1 which is not possible for an optimal P. If q is odd and p=3, then p+q=4+r, therefore:

$$\pi_{\text{opt}}(S_{3,n}) \ge \pi_{\text{opt}}(S_{3,c}) + \pi_{\text{opt}}(S'_{3,n-c}) = 3l + q + 3j + p - 1 = 3(l+j) + 3 + r = 3k + r.$$

Now we consider $S'_{3,n}$. We may assume n is odd because $S'_{3,n} \cong S_{3,n}$ when n is even. If r=3, then:

$$\pi_{\text{opt}}(S'_{3,4k+3}) \ge \pi_{\text{opt}}(S'_{3,4l}) + \pi_{\text{opt}}(S'_{3,4j+3}) \ge 3l + 3j + 2 = 3k + 2 \qquad \text{if } q = 0$$

$$\pi_{\text{opt}}(S'_{3,4k+3}) \ge \pi_{\text{opt}}(S'_{3,4l+1}) + \pi_{\text{opt}}(S_{3,4j+2}) \ge 3l + 1 + 3j + 2 = 3k + 3 \qquad \text{if } q = 1$$

$$\pi_{\text{opt}}(S'_{3,4k+3}) \ge \pi_{\text{opt}}(S'_{3,4l+2}) + \pi_{\text{opt}}(S'_{3,4j+1}) \ge 3l + 2 + 3j + 1 = 3k + 3 \qquad \text{if } q = 2$$

$$\pi_{\text{opt}}(S'_{3,4k+3}) \ge \pi_{\text{opt}}(S'_{3,4l+3}) + \pi_{\text{opt}}(S_{3,4j}) \ge 3l + 2 + 3j = 3k + 2 \qquad \text{if } q = 3.$$

Finally, when r = 1:

$$\begin{split} \pi_{\mathrm{opt}}(S'_{3,4k+1}) &\geq \pi_{\mathrm{opt}}(S'_{3,4l}) + \pi_{\mathrm{opt}}(S'_{3,4j+1}) \geq 3l + 3j + 1 = 3k + 1 & \text{if } q = 0 \\ \pi_{\mathrm{opt}}(S'_{3,4k+1}) &\geq \pi_{\mathrm{opt}}(S'_{3,4l+1}) + \pi_{\mathrm{opt}}(S_{3,4j}) \geq 3l + 1 + 3j = 3k + 1 & \text{if } q = 1 \\ \pi_{\mathrm{opt}}(S'_{3,4k+1}) &\geq \pi_{\mathrm{opt}}(S'_{3,4l+2}) + \pi_{\mathrm{opt}}(S'_{3,4j+3}) \geq 3l + 2 + 3j + 2 = 3k + 1 & \text{if } q = 2 \\ \pi_{\mathrm{opt}}(S'_{3,4k+1}) &\geq \pi_{\mathrm{opt}}(S'_{3,4l+3}) + \pi_{\mathrm{opt}}(S_{3,4j+2}) \geq 3l + 2 + 3j + 2 = 3k + 1 & \text{if } q = 3. \end{split}$$

We have checked each case and the resulting lower bound is at least as strong as the desired one, hence the proof is complete. \Box

2.2 Four-wide staircases

In this case the proof is simpler because $S'_{4,n}$ is isomorphic to $S_{4,n}$.

Theorem 2.7

$$\pi_{\text{opt}}(S_{4,4k+r}) = 3k + r$$

except in the following small cases: $\pi_{opt}(S_{4,1}) = 2$, $\pi_{opt}(S_{4,2}) = 3$.




Figure 5: Optimal distributions of small $S_{4,n}$ graphs. For the $n \leq 3$ cases refer to the previous figures.

PROOF: The optimal distributions of small graphs are shown in Figure 5. For larger n, we obtain a solvable distribution of $S_{4,n}$ in the following way: We partition $S_{4,n}$ into k-1 copies of $S_{4,4}$ and one copy of $S_{4,4+r}$ for some r, and we use the optimal distributions shown in Figure 5 in each of the parts. This gives the required upper bound.

To obtain the lower bound for $n \geq 8$, we assume that the statement is true for all values below n. We have $\pi_{\text{opt}}(S_{4,n}) \leq n-2$ by the upper bound so Claim 2.6 can be applied yielding

$$\pi_{\text{opt}}(S_{4,n}) \ge \pi_{\text{opt}}(S_{4,c}) + \pi_{opt}(S_{4,n-c}) \ge 3l + q + 3j + p \ge 3k + r,$$

where
$$n = 4k + r$$
, $c = 4l + q$, $n - c = 4j + p$ and $r, q, p \in \{0, 1, 2, 3\}$.

2.3 Five-wide staircases

In this case we again must distinguish between the graphs $S_{5,n}$ and $S'_{5,n}$ when n is odd. Fortunately, we are in a simpler situation than in the three-wide case because the formula optimal pebbling number of both these graphs turns out to be the same. The formula depends on the value of n modulo 5, so in this section let n = 5k + r where $r \in \{0, 1, 2, 3, 4\}$.

Theorem 2.8

$$\pi_{\mathrm{opt}}(S_{5,5k+r}) = \pi_{\mathrm{opt}}(S'_{5,5k+r}) = 4k + r,$$
except for $n \in \{1, 2, 3, 7\}$. $\pi_{\mathrm{opt}}(S'_{5,3}) = \pi_{\mathrm{opt}}(S'_{5,3}) = 4$ and $\pi_{\mathrm{opt}}(S'_{5,7}) = 7$.

PROOF: The smaller cases can be seen in Figure 6. Solvable distributions with the given sizes can be constructed with the proper concatenation of the optimal distributions of $S_{5,4}$, $S_{5,5}$, $S_{5,6}$, $S_{5,7}$, $S_{5,8}$, $S_{5,9}$, $S'_{5,5}$ and $S'_{5,9}$. For large n use many copies of $S_{5,5}$ and $S'_{5,5}$ in the middle and suitably extend it on the ends with the other distributions.

We can still apply Claim 2.6 when $n \ge 10$. The formula is the same for $S_{5,n}$ and $S'_{5,n}$, therefore we introduce the notation $S^*_{5,n}$ to denote either $S_{5,n}$ or $S'_{5,n}$. We can formalize the statement of Claim 2.6 in one inequality:

$$\pi_{\text{opt}}(S_{5,n}^*) \ge \pi_{\text{opt}}(S_{5,c}^*) + \pi_{\text{opt}}(S_{5,n-c}^*) = 4l + q + 4j + p \ge 4k + r,$$

where
$$n = 5k + r$$
, $c = 5l + q$, $n - c = 5j + p$ and $r, q, p \in \{0, 1, 2, 3, 4\}$.

2.4 Six-wide staircases

To handle this case and the seven-wide case we need some additional tools. We start by recalling some known results about the path.

Lemma 2.9 ([1]) A 2-optimal distribution of P_n consists of prime segments separated by single unoccupied vertices, where a prime segment is a subpath of one of two possible types. Either all but one of the vertices have one pebble and one vertex has two pebbles or there are three consecutive vertices with zero, four and zero pebbles, in that order, and the remaining vertices having one pebble.

A corollary of this lemma is that under a 2-optimal distribution no vertex is 5-reachable, and if a vertex is 3-reachable, then it has to contain 4 pebbles. The neighbours of a vertex v having 4 pebbles can get pebbles only from v. Similarly, each of the neighbours of a vertex u containing exactly 2 pebbles can get only one pebble and only from u. Using these facts we obtain the following statement for staircases.




Figure 6: Optimal distributions of small $S_{5,n}$ graphs. For the cases where $n \leq 4$ refer to the previous figures.

Lemma 2.10 Let G be a staircase graph with n slashes. If P is a pebbling distribution on G with n+1 pebbles, such that each slash is 2-reachable under P, then:

- Each slash contains at most 4 pebbles.
- If a slash is 3-reachable, then it contains 4 pebbles.
- if a slash has at least one pebble, then at most one pebble can be moved to that slash with a pebbling move.
- If a slash contains 4 pebbles, then the adjacent slashes have no pebbles, furthermore they can get pebbles only from this slash.
- If a slash contains 2 pebbles, then a pebbling move can only be performed ending in that slash if a pebbling move was first performed beginning on that slash.

PROOF: We simply apply the collapsing function defined in the proof of Lemma 2.3. The collapsed distribution on P_n is 2-optimal so Lemma 2.9 applies. All of the statements then follow from easily observable facts about prime segments. \square

Theorem 2.11

10.

$$\pi_{\rm opt}(S_{6,n}) = n,$$
 except for $n \in \{1,2,3,4,8,9\}$. $\pi_{\rm opt}(S_{6,3}) = \pi_{\rm opt}(S_{6,4}) = 5$, $\pi_{\rm opt}(S_{6,8}) = 9$ and $\pi_{\rm opt}(S_{6,9}) = 1$

To prove the lower bound we cannot use exactly the same method which we used for the narrower staircases because Lemma 2.4 does not apply when $|P| \ge n - 1$. We can overcome this difficulty with the next lemma.

Lemma 2.12 There is no solvable distribution P on $S_{6,n}$ with size at most n, such that each inner slash is 2-reachable.




Figure 7: Optimal distributions of small $S_{6,n}$ graphs. For $n \leq 5$ cases check the previous figures.




Figure 8: Labelling of the vertices

PROOF: Indirectly assume that such a P exists. The size of P can be either n-1 or n, otherwise an inner slash must be not 2-reachable according to Lemma 2.4. Without the loss of generality, we assume that the first slash is not 2-reachable.

Since P is solvable, we can reach all vertices of the first slash. If the first slash has exactly one pebble, then we need to move another pebble to reach its unoccupied vertices. This means that it is 2-reachable, therefore the first slash cannot have a pebble.

Put a pebble on the first slash and, in the case of |P|=n-1, another one at the last slash. This results a new distribution P' and each slash is 2-reachable under it. Therefore |P'|=n+1 and, if we collapse the graph into P_n with ϕ , then the collapsed distribution P'_{ϕ} is a 2-optimal distribution.

We will require some additional notation. Vertex names are shown in Figure 8. Now we exploit Lemma 2.10.

The first slash contains exactly one pebble under P'. By Lemma 2.10 (points 2 and 4) the second slash cannot be 3-reachable under P'. So it has at most two pebbles. v_3 has to be 2-reachable under P and P' because u_3 can be reachable only from this vertex. The reachability of u_1 requires that v_1 or v_2 is also 2-reachable.

If the second slash has two pebbles, then it is clear from the last statement of Lemma 2.10 that at most one vertex in that slash is 2-reachable. This contradicts that v_3 and one of v_1 or v_2 are 2-reachable.

When the second slash has exactly one pebble, then the third statement of Lemma 2.10 implies at most one vertex is 2-reachable in the second slash, which is not enough.

So neither the first nor the second slash contain a pebble under P, but the reachability of the first slash requires that the third slash is 4-reachable under P'. According to Lemma 2.10 the third slash must contain exactly 4 pebbles. Every vertex in the first three slashes must be reachable using only the 4 pebbles on slash three, but this is impossible (in particular we cannot reach both u_3 and v_1). \square

PROOF OF THEOREM 2.11: For small optimal distributions see Figure 7. Solvable distribution with n pebbles in cases when $n \ge 10$ can be created by combining the optimal distributions of $S_{6.5}$, $S_{6.6}$ and $S_{6.7}$.

Assume that the lower bound is proved for each integer less than n. Let P be an optimal distribution. We know that the size of P is at most n. By Lemma 2.12 we have that some inner slash is not 2-reachable. Then we can apply Claim 2.6 and the induction hypothesis, which gives that $|P| \ge c + n - c = n$. This completes the proof. \square

2.5 Seven-wide staircase graphs

Lemma 2.13 There is no solvable distribution P on $S_{7,n}$ with size at most n, such that each inner slash is 2-reachable.

PROOF: Assume the contrary. We have a solvable distribution P on $S_{7,n}$ with size n, such that each inner slash is 2-reachable.

Now we collapse $S_{7,n}$ to $S_{6,n}$ by mapping the first and the third negative diagonal of $S_{7,n}$ to one diagonal of $S_{6,n}$. For an example see Figure 9.




Figure 9: Collapsing $S_{7,6}$ to $S_{6,6}$.

Each pebbling sequence which is executable under P determines an executable pebbling sequence in P_{ϕ} which moves the same amount of pebbles to the slashes and moves at least as many pebbles to the same vertices except for the deleted ones. Therefore, P_{ϕ} is solvable and each inner slash is 2-reachable. This contradicts Lemma 2.12.

Theorem 2.14 Let $S_{7,n}^*$ be $S_{7,n}$ or $S_{7,n}'$, then

$$n+1 \le \pi_{\text{opt}}(S_{7,n}^*) \le n+3.$$

The lower bound is sharp for graphs $S_{7,5}$, $S_{7,6}$, $S_{7,7}$, $S_{7,8}$ and every $S'_{7,n}$ where $n \equiv 3 \mod 4$.

The proof of the lower bound is slightly different compared to the previous proofs. The main reason is that we do not have a tool for handling distributions with n+1 pebbles. n+1 pebbles is enough to construct a 2-solvable distribution on the n-vertex path, so we can not guarantee a cut which creates a smaller instance of the problem.

We overcome this difficulty with an indirect assumption. We assume that there is a minimal counterexample with at most n pebbles. In contrast, we show that it is not minimal.

Unfortunately we cannot apply this idea for values larger than n+1. Therefore we cannot prove an exact formula for $\pi_{\text{opt}}(S_7^*)$.




Figure 10: General pattern to make solvable distributions for large graphs with n+O(1) pebbles.




Figure 11: The optimal distributions of small $S_{7,n}$ and $S'_{7,n}$ graphs are pictured. For the $n \le 6$ cases, see the previous figures.

PROOF: We prove the upper bound first. For solvable small distributions see Figure 11, and for a general pattern see Figure 10.

The optimal distribution for $S_{7,4k+3}'$ is obtained by putting 1 pebble on the vertices neighboring the degree 1 vertices, and 4 pebbles at k vertices as shown in the general pattern. The solvable distribution of $S_{7,4k+1}'$ is the concatenation of $S_{7,6}'$ and $S_{7,4(k-2)+3}'$, which has 4k+3 pebbles. Similarly the solvable distribution of $S_{7,4k}$ is the concatenation of $S_{7,5}$ and $S_{7,4(k-2)+3}'$, which has 4k+2 pebbles. The solvable distribution of $S_{7,4k+1}$ is given by the general pattern with 4k+4 pebbles. We add two more slashes to this such that the first has two pebbles to obtain a distribution for $S_{7,4k+3}$ with 4k+6 pebbles. For $S_{7,4k+2}$ the optimal distribution of $S_{7,4k+3}'$ can be used with 4k+4 pebbles.

To prove the lower bound, assume that the statement is not true, and we let $G = S_{7,n}^*$ be the smallest counterexample. This means that there is a solvable distribution on G with n pebbles. By Lemma 2.13 and Lemma 2.5 we can break the graph into two smaller parts, and P induces a solvable distribution on both. One of the graphs has at most as many pebbles as slashes, but this means that we found a smaller counterexample, which is a contradiction. \square

Conjecture 2.15 *Each solvable distribution which was mentioned in the proof is optimal, which implies the following when* $n \ge 10$:

$$\pi_{\text{opt}}(S_{7,n}) = \begin{cases} n+2 & \text{if } n \equiv 0 \mod 2\\ n+3 & \text{if } n \equiv 1 \mod 2 \end{cases}$$

$$\pi_{\text{opt}}(S'_{7,n}) = \begin{cases} n+2 & \text{if } n \equiv 1 \mod 4\\ n+1 & \text{if } n \equiv 3 \mod 4. \end{cases}$$

3 Wider staircases

Question 1 What is the optimal pebbling number of $S_{8,n}$?

We determined the asymptotic behavior when $n \leq 7$, but we think that the general behavior of the eight-wide case differs from the seven-wide case. We obtained $\pi_{\rm opt}(S_{8,8})=11$ by computer search. Unfortunately, even the n=9 case requires more computational power than an average PC has. We have some partial results:

Let P be an optimal distribution of $S_{m,n}$. Expand $S_{m,n}$ with an additional m+1th negative diagonal to obtain $S_{m+1,n}$. Construct P' from P by placing additional pebbles at intersections of the mth negative diagonal and every fourth slash starting from the second one. Put a pebble to the last vertex of the mth negative diagonal if it has not obtained one yet. For an example see Figure 12.




Figure 12: A solvable distribution of $S_{8,8}$. Note that it is not optimal.

Claim 3.1 P' is a solvable distribution of $S_{m+1,n}$.

PROOF: The vertices where we placed additional pebbles are 2-reachable under P'. Each vertex of the 8th negative diagonal is adjacent to such a vertex, thus it is reachable.

We believe that if for each optimal pebbling distribution P for $S_{7,n}$ we form the distribution P', then these distributions will have size equal to the optimal pebbling number of $S_{8,n}$ asymptotically.

Conjecture 3.2

$$\pi_{\text{opt}}(S_{8,n}) = \frac{5}{4}n + O(1).$$

Finally, it seems that by simply duplicating an optimal distribution for $S_{7,n}$ k times, we can obtain a distribution of asymptotically optimal size for $S_{7k,n}$.

Conjecture 3.3 *For all* $k \ge 1$ *, we have*

$$\pi_{\text{opt}}(S_{7k,n}) = kn + O(1).$$

Acknowledgment

The researach of Ervin Győri and Gyula Y. Katona is partially suported by National Research, Development and Innovation Office NKFIH, grant K116769. The research of Gyula Y. Katona and László F. Papp is partially suported by National Research, Development and Innovation Office NKFIH, grant K108947.

References

- [1] D.P. Bunde, E. W. Chambers, D. Cranston, K. Milans, D. B. West, *Pebbling and optimal pebbling in graphs* J. Graph Theory 57 no. 3. (2008) pp. 215–238.
- [2] T. Friedman, C.Wyels. *Optimal pebbling of paths and cycles* arXiv:math/0506076 [math.CO]
- [3] H. Fu, C. Shiue, *The optimal pebbling number of the caterpillar*, Taiwanese Journal of Mathematics, **13** no. 2A (2009) pp. 419–429
- [4] H. Fu, C. Shiue, *The optimal pebbling number of the complete m-ary tree*, Discrete Mathematics, **222** 1–3 (2000) pp. 89–100
- [5] E. Győri, G. Y. Katona, L. F. Papp *Constructions for the optimal pebbling of grids* arXiv:1601.02229 [math.CO]
- [6] E. Győri, G. Y. Katona, L. F. Papp, C. Tompkins *A lower bound on the optimal pebbling number of grids* in preparation
- [7] L. Pachter, H.S. Snevily, B. Voxman *On pebbling graphs* Congressus Numerantium, **107** (1995) pp. 65–80
- [8] K. Milans, B. Clark, *The complexity of graph pebbling* SIAM J. Discrete Math. **20** no. 3 (2006) pp. 769–798.
- [9] C. Xue, C. Yerger *Optimal Pebbling on Grids* Graphs and Combinatorics, **32** no. 3 (2016) pp. 1229–1247