

## **GrassPlot – a database of multi-scale plant diversity in Palaearctic grasslands**

Jürgen Dengler\*, Viktoria Wagner, Iwona Dembicz, Itziar García-Mijangos, Alireza Naqinezhad, Steffen Boch, Alessandro Chiarucci, Timo Conradi, Goffredo Filibeck, Riccardo Guarino, Monika Janišová, Manuel J. Steinbauer, Svetlana Aćić, Alicia T.R. Acosta, Munemitsu Akasaka, Marc-Andre Allers, Iva Apostolova, Irena Axmanová, Branko Bakan, Alina Baranova, Manfred Bardy-Durchhalter, Sándor Bartha, Esther Baumann, Thomas Becker, Ute Becker, Elena Belonovskaya, Karin Bengtsson, José Luis Benito Alonso, Asun Berastegi, Ariel Bergamini, Ilaria Bonini, Hans Henrik Bruun, Vasyl Budzhak, Alvaro Bueno, Juan Antonio Campos, Laura Cancellieri, Marta Carboni, Cristina Chocarro, Luisa Conti, Marta Czarniecka-Wiera, Pieter De Frenne, Balázs Deák, Yakiv P. Didukh, Martin Diekmann, Christian Dolnik, Cecilia Dupré, Klaus Ecker, Nikolai Ermakov, Brigitta Erschbamer, Adrián Escudero, Javier Etayo, Zuzana Fajmonová, Vivian A. Felde, Maria Rosa Fernández Calzado, Manfred Finckh, Georgios Fotiadis, Mariano Fracchiolla, Anna Ganeva, Daniel García-Magro, Rosario G. Gavilán, Markus Germany, Itamar Giladi, François Gillet, Gian Pietro Giusso del Galdo, Jose M. González, John-Arvid Grytnes, Michal Hájek, Petra Hájková, Aveliina Helm, Mercedes Herrera, Eva Hettenbergerová, Carsten Hobohm, Elisabeth M. Hüllbusch, Nele Ingerpuu, Ute Jandt, Florian Jeltsch, Kai Jensen, Anke Jentsch, Michael Jeschke, Borja Jiménez-Alfaro, Zygmunt Kącki, Kaoru Kakinuma, Jutta Kapfer, Ali Kavgacı, András Kelemen, Kathrin Kiehl, Asuka Koyama, Tomoyo F. Koyanagi, Łukasz Kozub, Anna Kuzemko, Magni Olsen Kyrkjeeide, Sara Landi, Nancy Langer, Lorenzo Lastrucci, Lorenzo Lazzaro, Chiara Lelli, Jan Lepš, Swantje Löbel, Arantzazu L. Luzuriaga, Simona Maccherini, Martin Magnes, Marek Malicki, Corrado Marcenò, Constantin Mardari, Leslie Mauchamp, Felix May, Ottar Michelsen, Joaquín Molero Mesa, Zsolt Molnár, Ivan Y. Moysiyenko, Yuko K. Nakaga, Rayna Natcheva, Jalil Noroozi, Robin J. Pakeman, Salza Palpurina, Meelis Pärtel, Ricarda Pätsch, Harald Pauli, Hristo Pedashenko, Robert K. Peet, Remigiusz Pielech, Nataša Pipenbaher, Chrisoula Pirini, Zuzana Plesková, Mariya A. Polyakova, Honor C. Prentice, Jennifer Reinecke, Triin Reitalu, Maria

Pilar Rodríguez-Rojo, Jan Roleček, Vladimir Ronkin, Leonardo Rosati, Ejvind Rosén, Eszter Ruprecht, Solvita Rusina, Marko Sabovljević, Ana María Sánchez, Galina Savchenko, Oliver Schuhmacher, Sonja Škornik, Marta Gaia Sperandii, Monika Staniaszek-Kik, Zora Stevanović-Dajić, Marin Stock, Sigrid Suchrow, Laura M.E. Sutcliffe, Grzegorz Swacha, Martin Sykes, Anna Szabó, Amir Talebi, Cătălin Tănase, Massimo Terzi, Csaba Tölgyesi, Marta Torca, Péter Török, Béla Tóthmérész, Nadezda Tsarevskaya, Ioannis Tsiripidis, Rossen Tzonev, Atushi Ushimaru, Orsolya Valkó, Eddy van der Maarel, Thomas Vanneste, Iuliia Vashenyak, Kiril Vassilev, Daniele Viciani, Luis Villar, Risto Virtanen, Ivana Vitasović Kosić, Yun Wang, Frank Weiser, Julia Went, Karsten Wesche, Hannah White, Manuela Winkler, Piotr T. Zaniwski, Hui Zhang, Yaron Ziv, Sergey Znamenskiy & Idoia Biurrun

\*Corresponding author's address: Vegetation Ecology Group, Institute of Natural Resource Sciences (IUNR), Zurich University of Applied Sciences (ZHAW), Grüentalstr. 14, 8820 Wädenswil, Switzerland; [juergen.dengler@zhaw.ch](mailto:juergen.dengler@zhaw.ch). Complete addresses of all authors can be found at the bottom of the paper.

**Running title:** GrassPlot – Long Database Report

**Abstract:** GrassPlot is a collaborative vegetation-plot database organised by the Eurasian Dry Grassland Group (EDGG) and listed in the Global Index of Vegetation-Plot Databases (GIVD ID EU-00-003). GrassPlot collects plot records (relevés) from grasslands and other open habitats of the Palaearctic biogeographic realm. It focuses on precisely delimited plots of eight standard grain sizes (0.0001; 0.001; ... 1,000 m<sup>2</sup>) and on nested-plot series with at least four different grain sizes. The usage of GrassPlot is regulated through bylaws that intend to balance the interests of data contributors and data users. The current version (v. 1.00) contains data for approximately 170,000 plots of different sizes and 2,800 nested-plot series. The key components are richness data and metadata. However, most included datasets also encompass compositional data. About 14,000 plots have near-complete records of terricolous bryophytes and lichens in addition to vascular plants. At present, GrassPlot contains data from 36 countries throughout the Palaearctic, spread across elevational gradients and major grassland types. GrassPlot with its multi-scale and multi-taxon focus complements the larger international vegetation-plot databases, such as the European Vegetation Archive (EVA) and the global database “sPlot”. Its main aim is to facilitate studies on the scale- and taxon-dependency of biodiversity patterns and drivers along macroecological gradients. GrassPlot is a dynamic database and will expand through new data collection coordinated by the elected Coordinating Board. We invite researchers with suitable data to join GrassPlot. Researchers with project ideas addressable with GrassPlot data are welcome to submit proposals to the Governing Board.

**Keywords:** biodiversity; European Vegetation Archive (EVA); Eurasian Dry Grassland Group (EDGG); grassland vegetation; GrassPlot; macroecology; multi-taxon; nested plot, scale-dependence; species-area relationship (SAR); sPlot; vegetation-plot database.

**Abbreviations:** EDGG = Eurasian Dry Grassland Group; EVA = European Vegetation Archive; GrassPlot = Database of Scale-Dependent Phytodiversity Patterns in Palaearctic Grasslands; SAR = species-area relationship.

Submitted: 15 January 2018

Co-ordinating Editor: Florian Jansen

## GIVD Fact Sheet

#Separate file, needs to be first updated in GIVD#

### Introduction

The Palaearctic is the largest biogeographic realm of the world (Olson et al. 2001). It contains large areas of grasslands (9.7 million km<sup>2</sup> or 22% of the Palaearctic realm), of both natural and secondary origin (Török & Dengler in press). These grasslands harbour a high diversity of many taxonomic groups and encompass contrasting local diversity. While some grassland types contain the majority of global vascular plant diversity records surveyed at the small-scale (Wilson et al. 2012), others can be very species poor (Dengler et al. 2016a). The high variation in local diversity and wide environmental gradients occupied (different biomes, elevational zones from the sea level to the alpine, diverse soil types, etc.) make Palaearctic grasslands an ideal study object for understanding patterns and drivers of local plant diversity. Moreover, since many Palaearctic grasslands contain significant numbers of bryophytes and lichens, they allow testing of biodiversity patterns across taxa with contrasting biological traits (e.g. Löbel et al. 2006).

Plant community ecology is aimed at describing and understanding patterns of species composition and diversity recorded in small plots (“relevés” in phytosociology) in order to infer patterns and processes at local or regional scales. Macroecology, by contrast, analyses and explains patterns of diversity and its components across large regions, such as continents or the planet. The latter so far has typically relied on single species distribution data derived from sources such as the Global Biodiversity Information Facility (GBIF; <https://www.gbif.org/>) and gridded to coarse spatial grains, such as cells of 10,000 km<sup>2</sup> (Beck et al. 2012). This is far from the grain sizes at which relevant processes as the interaction among species and with their abiotic environment occur (Beck et al. 2012). In Europe, local studies on plant community composition, typically using the phytosociological method (Dengler et al. 2008; Guarino et al. 2018), surged in the last century (Schaminée et al. 2009). However, their grain sizes (e.g. Chytrý & Otýpková 2003) are still significantly larger than those at which some local processes, such as biotic interactions and edaphic filters (Siefert et al. 2012; Turtureanu et al. 2014), might act, which could be distances of centimetres or decimetres. Moreover, local studies have been criticized as being idiosyncratic and failing to derive general trends across regions (Chiarucci 2007; Dengler et al. 2011; Beck et al. 2012). A way to overcome this shortcoming, and to link community ecology to macroecology, is to

unite individual vegetation-plot datasets into big databases that cover large geographic areas (Dengler et al. 2011; Wiser 2016).

The European Vegetation Archive (EVA; Chytrý et al. 2016) and the global vegetation-plot database “sPlot” (Dengler & sPlot Core Team 2014), each with more than one million plots, are examples for recently assembled large vegetation-plot databases (Appendix 1). The first pilot biodiversity studies of fine-grain plot data across large biogeographic extents (e.g. Wagner et al. 2017) demonstrated the opportunities of large vegetation-plot databases. However, analyses based on large databases face methodological difficulties. First, plot sizes can vary considerably among different schools, regions, decades and vegetation types (Chytrý & Otýpková 2003). In some phytosociological schools, plots might not even be delimited in the field, have rather vague boundaries or irregular shapes to ensure so-called “floristic homogeneity” (e.g. Géhu 2010). Second, the degree of completeness of the species list recorded within each plot can vary due to sampling effort or taxonomic skills. Moreover, in certain phytosociological traditions, species or even whole life forms that were perceived as not belonging to an “ideal” community were (and sometimes still are) not recorded even when present in the plot (e.g. Géhu 1980).

While it is generally accepted that patterns and drivers of biodiversity are scale-dependent, this idea is based largely on theoretical considerations (Shmida & Wilson 1985) and insights from meta-analyses (Field et al. 2009; Siefert et al. 2012). By contrast, this hypothesis was rarely investigated in the field, using nested multi-scale studies from the same location and plant community (e.g. Podani et al. 1993; Reed et al. 1993; Turtureanu et al. 2014). Moreover, notwithstanding that terrestrial vegetation is made up of taxa with contrasting biological traits, including vascular plants, bryophytes and lichens, large vegetation databases to date have been focusing on vascular plants (see Appendix 1).

The outlined aspects inspired us to set up GrassPlot, the “Database of Scale-Dependent Phytodiversity Patterns in Palaeartic Grasslands”. The aim was to complement EVA and sPlot with a specialised and selective database of multi-scale (and often multi-taxon) data from Palaeartic grasslands exhaustively sampled on precisely delimited plots. We use this Long Database Report to introduce GrassPlot to the scientific community, summarise its current content and demonstrate arising opportunities in the concert of existing databases.

## History and governance of GrassPlot

The interest of some co-authors in small-scale species-area relationships (SARs) (Dengler 2009a; Wilson et al. 2012) motivated several regional studies in various dry grasslands in Europe (Dengler et al. 2004; Dengler & Boch 2008) and led then to the launch of the annual Research Expeditions (now: Field Workshops) of the European Dry Grassland Group (EDGG; now: Eurasian Dry Grassland Group; Vrahnakis et al. 2013; <http://www.edgg.org>). The first expedition took place in 2009 in Transylvania, Romania. It revealed grasslands that scored several global records of small-scale vascular plant diversity (Wilson et al. 2012). With the aim of facilitating overarching studies of SARs, Dengler et al. (2012) compiled available data in the “Database Species-Area Relationships in Palaearctic Grasslands” with 727 nested-plot series comprising a total of 7,202 individual plot observations. The EDGG Field Workshops continued to record standardised multi-scale vegetation data of grasslands across the Palaearctic, from Spain to Siberia (Vrahnakis et al. 2013). This effort resulted in several regional analyses of biodiversity patterns (e.g. Turtureanu et al. 2014; Polyakova et al. 2016). By 2016, the accumulation of data from the EDGG Field Workshops and from other researchers who had started to adopt the EDGG sampling methodology (Madari & Tănase 2016; Cancellieri et al. 2017) prompted the EDGG to create a comprehensive database. Initial steps included the compilation of an overview of existing datasets (Dengler et al. 2016a) and a description of the sampling approach (Dengler et al. 2016b), based on earlier suggestions by Dengler (2009b).

During an international workshop in Bayreuth in March 2017, the database was formally established with the name “GrassPlot” as a collaborative initiative within the EDGG (see <http://bit.ly/2BIHmnq>; logo in Fig. 1). The Data Property and Governance Rules (Bylaws) of GrassPlot (Supplement S1) have been set up to balance the interests of data providers and data users in a fair and transparent manner. In particular, data contributors remain owners of their data, are informed about any plans to use their data and can opt-in as active co-authors of papers. Depending on the size and complexity, a dataset in GrassPlot can have one or several owners. The GrassPlot Consortium is made up of these data owners and the 17 participants of the initial GrassPlot workshop. The Consortium elects the Governing Board every two years. The current Governing Board consists of J. Dengler (as Custodian), I. Biurrun (as Deputy Custodian) as well as T. Conradi, I. Dembicz, R. Guarino and A. Naqinezhad (as other members). It is responsible for managing GrassPlot and for handling data requests as well as offering co-authorship under the Bylaws. Paper

proposals can be submitted only by members of the GrassPlot Consortium or by author teams at least comprising one Consortium member.

GrassPlot is registered in the Global Index of Vegetation-Plot Databases (GIVD; <http://www.givd.info/>; Dengler et al. 2011) under the ID EU-00-003 and has its own website with regularly updated information on the current content (<http://bit.ly/2qKTQt2>). Moreover, the Governing Board actively approached researchers worldwide whose publications were based on data that potentially met the GrassPlot criteria. This has maintained a constant inflow of datasets, accompanied by a substantial growth of the Consortium to currently 198 members from 35 countries.

## Technical implementation

Since GrassPlot focuses on species richness and species-area relationships, its header data are stored in a single large spread sheet, with every row representing a (sub-) plot and storing information on species richness, the locality, vegetation structure and ecological parameters, plus an indication of nesting within larger plots. We adopted this solution because the nested nature of many plots is something that could not be easily accustomed in the common software for vegetation management (Turboveg 2; Hennekens & Schaminée 2001). Two additional spreadsheets list the metadata for the correspondent datasets and contact information of the Consortium members. As such, GrassPlot is organised differently from EVA and its contributing databases (Chytrý et al. 2016; see Appendix 1).

Compositional data, i.e. species composition and cover values, were not the original focus of GrassPlot and are not required parameters for new data (see Appendix 1). However, since they were widely available for most individual datasets, they were also incorporated. GrassPlot stores these data in long format .txt files. The latter were created semi-automatically based on the original, wide-format tables, provided by the data owners. Species names are taxonomically and nomenclaturally harmonized by a series of documented and repeatable R scripts (R Core Team 2017), similar to those used in sPlot (Purschke 2017). By this circumstance we are not able to resolve identical names that refer to different taxonomic concepts (Jansen & Dengler 2010; see Appendix 1). This way, the data do not lend themselves for syntaxonomic analyses but they are a solid ground to analyse local diversity patterns and assembly rules.

The simple structure of the richness- and metadata in GrassPlot allows updates with little delay when new data are submitted. By contrast, compositional data are usually integrated with a time lag as they can come in many different formats, and the harmonisation of their taxonomies is

challenging. GrassPlot data are stored in the .xlsx and .txt formats, which can be directly fed into different analytical software. While GrassPlot is updated continuously, each version is numbered and stored, enabling analyses with older versions.

## Content of GrassPlot v. 1.00

GrassPlot collects vegetation-plot data of grasslands in the widest sense (i.e. everything except forests, tall shrublands, aquatic and segetal communities) from the Palaearctic biogeographic realm (i.e. Europe, North Africa, West, Central, North and Northeast Asia). With respect to sampling methodology, GrassPlot is more restrictive than typical vegetation-plot databases. It only includes data of plots with one of our eight standard grain sizes: 0.0001, 0.001 (or 0.0009), 0.01, 0.1 (or 0.09), 1, 10 (or 9) 100, 1,000 (or 900 or 1,024) m<sup>2</sup>. However, we also allow deviations up to 10% from these grain sizes, e.g. 9 m<sup>2</sup> instead of 10 m<sup>2</sup>. Nested-plot series with at least four different grain sizes are also included; for the latter, any grain size is allowed. Plots must have been precisely delimited in the field (e.g. with a tape around the perimeter or with frames for smaller sizes) and thoroughly been sampled at least for vascular plants, but preferentially also for terricolous bryophytes and lichens. GrassPlot accepts (i) pure richness data (together with the required metadata) or (ii) complete vegetation plots (compositional data), i.e. species identities with presence-absence, cover, abundance or any other measure of dominance.

The first publicly released GrassPlot version 1.00 of 14 January 2018 contains data from 126 contributing datasets (Supplements S2 and S3). In total, the database comprises 168,997 plots of different grain sizes and 2,797 nested-plot series with at least four grain sizes (often consisting of several subseries). Most contributors have assigned their plots to the semi-restricted access regime, few in “restricted access” and currently none in free access (Table 1). For the majority of plots (98%), the owners also provided compositional data although these are not fully integrated yet (Table 1).

Geographically, the plots range from Morocco in the west (9.2° W) to Japan in the east (161.6° E) and from Tibet (China) in the south (28.6° N) to Svalbard (Norway) in the north (77.9° N). The highest density of plots was recorded in temperate Europe (Fig. 2). In total, the plots originate from 36 countries, with Spain having the highest number (54,608 plots) and Austria the highest density (15.62 plots per 100 km<sup>2</sup>) of plots (Table 2). However, GrassPlot also contains relatively high densities of plots in countries that were hitherto only poorly represented in EVA (Chytrý et al. 2016) and sPlot (Dengler & sPlot Core Team 2014), namely Iran, Israel, Norway and Sweden. Plot



elevation ranges from sea level (0 m a.s.l.) to 5,197 m a.s.l., with the largest fraction encompassing 2001–3000 m a.s.l. (Table 1). In total, data were sampled during the period of 1948 to 2017, with 79% of all plots surveyed in the decade of 2000–2009 (Table 1). Currently, 74% of all plots are syntaxonomically assigned to a class or a more precise level (Table 3). The temperate dry grasslands of the *Festuco-Brometea* (21%) and the Oromediterranean *Festucetea indigestae* (18%) are the best represented classes.

The most frequent standard plot sizes are 0.01 m<sup>2</sup>, followed by 1 m<sup>2</sup> and 9–10 m<sup>2</sup> (Table 2). Data for the complete terricolous vegetation (vascular plants, terricolous bryophytes and lichens) are available for 14,064 of all plots (8.3%) (Table 4, Fig. 2). Methodologically, the majority of contributors used shoot sampling rather than rooted sampling (Table 1), which can make a big difference for the assessment of vascular plant richness at small spatial grains (Dengler 2008; Güler et al. 2016; Cancellieri et al. 2017). Among plot shapes, squares were most frequently employed (75%), followed by rectangles with 1:2 edge length ratio (23%). Circles are the most compact shape, but difficult to delimit (see Güler et al. 2016), and were used in less than 2% of the records. The geographic coordinates stored in GrassPlot are nearly always more accurate than 1 km and in 3.4% of plots have an accuracy of 1 m or less (Table 1). Many structural (e.g. cover and height of vegetation layers; biomass) and ecological (e.g. topography, soil, land use) parameters are stored by GrassPlot in header data fields with harmonized terminology and units of measurement (see Supplement S4).

## GrassPlot in the context of other large vegetation-plot databases

With EVA (Chytrý et al. 2016) and sPlot (Dengler & sPlot Core Team 2014) providing huge amounts of vegetation-plot data of any vegetation type across Europe and the world (see Appendix 1), respectively, the need of an additional supra-national database like GrassPlot could be questioned. Actually, EVA and sPlot are unprecedented in spatial coverage (see Appendix 1). Being set up as all-purpose databases, however, they are not always suited optimally for certain specific questions. For this reason, specialised smaller databases have emerged e.g. with special focus on provision of plots with extensive and standardised soil data measured in the plot (e.g. Wamelink et al. 2012), for comparison of ecological impacts (e.g. PREDICTS, not only vegetation: Hudson et al. 2014) or for time-series in permanent plots (e.g. GLORIA: Pauli et al. 2012; forestREplot: Verheyen et al. 2017).

GrassPlot was set up with the aim to assemble data from Palaearctic grasslands by focusing on a multi-scale and multi-taxon approach. Multi-scale data are either not covered by the other large international vegetation-plot databases such as EVA (Chytrý et al. 2016) and sPlot (Dengler & sPlot Core Team 2014) or, if covered not clearly labelled as such, reducing accessibility (see Appendix 1). While one might think that alternatively one could just use the huge amount of plots of different sizes found in “normal” vegetation-plot databases, tests have shown that with this approach not even the most simple scaling law in ecology, the species-area relationship (SAR), is realistically depicted (see Chytrý 2001; Dengler et al. 2006). Therefore, GrassPlot complements the existing databases by specifically filling the gap of multi-scale plot data. This enables analyses of scale-dependent patterns and processes across distant regions, which so far have been impossible. By contrast, EVA and sPlot are better suited for any type of analyses that requires high spatial coverage (see Appendix 1). GrassPlot is not suited for purposes of vegetation classification due to the low spatial coverage/high spatial autocorrelation and the fact that plant names are only matched by synonymy but not by concepts (taxonyms) (see Appendix 1). Certain types of analyses could benefit from conducting them parallel in EVA/sPlot and in GrassPlot. For example, patterns of plot-scale species richness in European grasslands could be captured with high spatial resolution through the data contained in EVA, but the results might be considerably biased by regional differences in the sampling methodology (e.g. the completeness of species records). The same study done with GrassPlot would suffer much less from differences in sampling quality, but hardly could produce an alpha-richness map of Europe, simply because the available data are much sparser (see Fig. 2). A combination of both data sources might thus allow taking advantage of both “approaches”.

While the majority of plots either are suited for EVA/sPlot or for GrassPlot, a rather small fraction is meeting the requirements of both (see Appendix 1): These are Palaearctic grassland plots on precisely delimited areas of 1, 9, 10 or 100 m<sup>2</sup> with thoroughly sampled species composition, including “importance values” (i.e. cover, abundance, biomass,...). It makes sense to include this limited amount of data in both EVA/sPlot and GrassPlot because they are stored in different formats that are readily prepared for different analyses. Good coordination between GrassPlot, EVA and sPlot is ensured because J. Dengler and I. Biurrun from the GrassPlot Governing Board are also involved in the EVA Coordinating Board and J. Dengler additionally in the sPlot Steering Committee. That way, redundant work is reduced and the effective inclusion of data whose qualities meet the criteria of several of these huge supranational databases in all of these is ensured (if data providers agree). Moreover, GrassPlot is also accepting small, local datasets that are in number of

plots far below the thresholds of EVA/sPlot. Several such small datasets together could then be provided to EVA or sPlot.

## **Resumé and outlook**

Despite being relatively small for an international vegetation-plot database, we believe that GrassPlot can become a valuable tool in “community macroecology”. While the big databases EVA and sPlot are better suited for the majority of purposes, GrassPlot can be advantageous for specific questions that require highly standardised data. Potential users are advised to select the most suitable database for a certain purpose based on the particular characteristics of these three (Appendix 1) and other databases.

Beyond that we hope that GrassPlot with its focus on methodological aspects of sampling and the prevalence for a few “standard” plot sizes, will encourage many vegetation scientists to consider these issues and thus promote the collection of highly comparable data sets. Noteworthy, the same plot sizes (or a subset of these), each separated from the next by one order of magnitude, had previously been proposed in various frameworks (Shmida 1984; Peet et al. 1998; Chiarucci et al. 2001; Dengler 2009b).

GrassPlot is a dynamic database that will continue to integrate suitable datasets in the future. Researchers in possession of data that meet the GrassPlot specification and who wish to join our Consortium are welcome to contact our database manager (I. Biurrun). We particularly seek data from underrepresented regions (most of Asia, North Africa and some parts of Europe; see Fig. 2) and vegetation types (e.g. mesic, wet and Mediterranean grasslands; see Table 3) as well as generally plots with recording of bryophytes and lichens. Readers who wish to address a research idea with GrassPlot data are welcome to submit a project proposal jointly with a Consortium member of their choice to the Governing Board.

## **Author contributions**

J.D. managed the predecessor databases of GrassPlot, while I.B. served as database manager from the start of GrassPlot onwards and V.W. handled the compositional data. J.D. led the writing of this report, with major contributions from V.W. as well as I.B., S.B., A.C., T.C., I.D., G.F., I.G.-M., R.G., M.J., A.N. and M.J.S. The figures were prepared by I.D. and the supplements by J.D., A.N. and I.G.-M. All other authors contributed data to GrassPlot, checked and approved the manuscript.

## Acknowledgements

We thank the BayIntAn program of Bavarian Research Alliance (<https://www.bayfor.org/en/research-funding/bayintan.php>; grant no. UBT\_2017\_58) as well as the Bayreuth Centre of Ecology and Environmental Research (BayCEER; <https://www.bayceer.uni-bayreuth.de/>) for funding the GrassPlot workshop in Bayreuth. Furthermore, we are grateful to the International Association of Vegetation Science (IAVS; <http://iavs.org/>), the Eurasian Dry Grassland Group (EDGG; <http://www.edgg.org/>) and the Förderkreis für Allgemeine Naturkunde (Biologie) (FAN(B); <http://www.fan-b.de/>) for supporting the EDGG Expeditions/Field Workshops and all the colleagues who contributed to the high quality data in GrassPlot without being listed as co-authors. Two anonymous reviewers helped to improve the manuscript with their suggestions.

## References

- Beck, J., Ballesteros-Mejia, L., Buchmann, C.M., Dengler, J., Fritz, S., Gruber, B., Hof, C., Jansen, F., Knapp, S., (...) & Dormann, C.F. 2012. What's on the horizon for macroecology? *Ecography* 35: 673–683.
- Cancellieri, L., Mancini, L.D., Sperandii, M.G. & Filibeck, G. 2017. In and out: Effects of shoot- vs. rooted-presence sampling methods on plant diversity measures in mountain grasslands. *Ecological Indicators* 72: 315–321.
- Chiarucci, A. 2007. To sample or not to sample? That is the question ... for the vegetation scientist. *Folia Geobotanica* 42: 209–216.
- Chiarucci, A., De Dominicis, V. & Wilson, J.B. 2001. Structure and floristic diversity in permanent monitoring plots in forest ecosystems of Tuscany. *Forest Ecology and Management* 141: 201–210.
- Chytrý, M. 2001. Phytosociological data give biased estimates of species richness. *Journal of Vegetation Science* 12: 439–444.
- Chytrý, M. & Otýpková, Z. 2003. Plot sizes used for phytosociological sampling of European vegetation. *Journal of Vegetation Science* 14: 563–570.
- Chytrý, M., Hennekens, S.M., Jiménez-Alfaro, B., Knollová, I., Dengler, J., Jansen, F., Landucci, F., Schaminée, J.H.G., Aćić, S., (...) & Yamalov, S. 2016. European Vegetation Archive (EVA): an integrated database of European vegetation plots. *Applied Vegetation Science* 19: 173–180.
- Dengler, J. 2008. Pitfalls in small-scale species-area sampling and analysis. *Folia Geobotanica* 43: 269–287.
- Dengler, J. 2009a. Which function describes the species-area relationship best? – A review and empirical evaluation. *Journal of Biogeography* 36: 728–744.
- Dengler, J. 2009b. A flexible multi-scale approach for standardised recording of plant species richness patterns. *Ecological Indicators* 9: 1169–1178.
- Dengler, J. & Boch, S. 2008. Sampling-design effects on properties of species-area curves – A case study from Estonian dry grassland communities. *Folia Geobotanica* 43: 289–304.

- Dengler, J. & sPlot Core Team. 2014. sPlot: the first global vegetation-plot database and opportunities to contribute. *IAVS Bulletin* 2014(2): 34–37.
- Dengler, J., Bedall, P., Bruchmann, I., Hoeft, I. & Lang, A. 2004. Artenzahl-Areal-Beziehungen in uckermärkischen Trockenrasen unter Berücksichtigung von Kleinstflächen – eine neue Methode und erste Ergebnisse. *Kieler Notizen zur Pflanzenkunde in Schleswig-Holstein und Hamburg* 32: 20–25.
- Dengler, J., Růsina, S., Boch, S., Bruun, H.H., Diekmann, M., Dierßen, K., Dolnik, C., Dupré, C., Golub, V.B., (...) & Zobel, M. 2006. Working group on dry grasslands in the Nordic and Baltic region – Outline of the project and first results for the class *Festuco-Brometea*. *Annali di Botanica, Nuova Serie*, 6: 1–28.
- Dengler, J., Chytrý, M. & Ewald, J. 2008. Phytosociology. In: Jørgensen, S.E. & Fath, B.D. (eds.) *Encyclopedia of ecology*, pp. 2767–2779. Elsevier, Oxford, UK.
- Dengler, J., Jansen, F., Glöckler, F., Peet, R.K., De Cáceres, M., Chytrý, M., Ewald, J., Oldeland, J., Finckh, M., (...) & Spencer, N. 2011. The Global Index of Vegetation-Plot Databases (GIVD): a new resource for vegetation science. *Journal of Vegetation Science* 22: 582–597.
- Dengler, J., Todorova, S., Becker, T., Boch, S., Chytrý, M., Diekmann, M., Dolnik, C., Dupré, C., Giusso del Galdo, G.P., (...) & Vassilev, K. 2012. Database Species-Area Relationships in Palaearctic Grasslands. *Biodiversity & Ecology* 4: 321–322.
- Dengler, J., Biurrun, I., Apostolova, I., Baumann, E., Becker, T., Berastegi, A., Boch, S., Dembicz, I., Dolnik, C., (...) & Weiser, F. 2016a. Scale-dependent plant diversity in Palaearctic grasslands: a comparative overview. *Bulletin of the Eurasian Dry Grassland Group* 31: 12–26.
- Dengler, J., Boch, S., Filibeck, G., Chiarucci, A., Dembicz, I., Guarino, R., Henneberg, B., Janišová, M., Marcenò, C., (...) & Biurrun, I. 2016b. Assessing plant diversity and composition in grasslands across spatial scales: the standardised EDGG sampling methodology. *Bulletin of the Eurasian Grassland Group* 32: 13–30.
- Field, R., Hawkins, B.A., Cornell, H.V., Currie, D.J., Diniz-Filho, A.F., Guégan, J.-F., Kaufman, D.M., Kerr, J.T., Mittelbach, G.G., (...) & Turner, J.R.G. 2009. Spatial species-richness gradients across scales: a meta-analysis. *Journal of Biogeography* 36: 132–147.
- Géhu, J.-M. 1980. La phytosociologie d'aujourd'hui. Méthodes et orientations. *Notiziario della Società Italiana di Fitosociologia* 16: 1–16.
- Géhu, J.-M. 2010. Problématique de l'étude phytosociologique des forêts. *Revue Forestière Française* 62: 219–228.
- Guarino, R., Willner, W., Pignatti, S., Attorre, F. & Loidi, J.J. 2018. Spatio-temporal variations in the application of the Braun-Blanquet approach in Europe. *Phytocoenologia* 48 (in press). DOI: 10.1127/phyto/2017/0181.
- Güler, B., Jentsch, A., Bartha, S., Bloor, J.M.G., Campetella, G., Canullo, R., Házi, J., Kreyling, J., Pottier, J., (...) & Dengler, J. 2016. How plot shape and dispersion affect plant species richness counts: implications for sampling design and rarefaction analyses. *Journal of Vegetation Science* 27: 692–703.
- Hennekens, S.M. & Schaminée, J.H.J. 2001. TURBOVEG, a comprehensive data base management system for vegetation data. *Journal of Vegetation Science* 12: 589–591.

- Hudson, L.W., Newbold, T., Contu, S., Hill, S.L.L., Lysenko, I., De Palma, A., Phillips, H.R.P., Senior, R.A., Bennett, D.J., (...) & Purvis, A. 2014. The PREDICTS database: a global database of how local terrestrial biodiversity responds to human impacts. *Ecology and Evolution* 4: 4701–4735.
- Jansen, F. & Dengler, J. 2010. Plant names in vegetation databases – a neglected source of bias. *Journal of Vegetation Science* 21: 1179–1186.
- Löbel, S., Dengler, J. & Hobohm, C. 2006. Species richness of vascular plants, bryophytes and lichens in dry grasslands: The effects of environment, landscape structure and competition. *Folia Geobotanica* 41: 377–393.
- Madari, C. & Tănase, C. 2016. Plant diversity-environment relationships in xeric grasslands in North-eastern Romania. *Applied Ecology and Environmental Research* 14: 111–127.
- Mucina, L., Bültmann, H., Dierßen, K., Theurillat, J.-P., Raus, T., Čarni, A., Šumberová, K., Willner, W., Dengler, J., (...) & Tichý, L. 2016. Vegetation of Europe: Hierarchical floristic classification system of vascular plant, bryophyte, lichen, and algal communities. *Applied Vegetation Science* 19, Suppl. 1: 3–264.
- Olson, D.M., Dinerstein, E., Wikramanayake, E.D., Burgess, N.D., Powell, G.V.N., Underwood, E.C., D’Amico, J.A., Itoua, I., Strand, H.E., (...) & Kassem, K.R. 2001. Terrestrial ecoregions of the world: a new map of life on Earth. *BioScience* 51: 933–938.
- Pauli, H., Gottfried, M., Dullinger, S., Abdaladze, O., Akhalkatsi, M., Benito, J.L., Coldea, G., Dick, J., Erschbamer, B., (...) & Grabherr, G. 2012. Recent plant diversity changes on Europe’s mountain summits. *Science* 336: 353–355.
- Peet, R.K., Wentworth, T.R. & White, P.S. 1998. A flexible, multipurpose method for recording vegetation composition and structure. *Castanea* 63: 262–274.
- Podani, J., Czárán, T. & Bartha, S. 1993. Pattern, area and diversity: the importance of spatial scale in species assemblages. *Abstracta Botanica* 17: 37–51.
- Polyakova, M.A., Dembiczy, I., Becker, T., Becker, U., Demina, O.N., Ermakov, N., Filibeck, G., Guarino, R., Janišová, M., (...) & Dengler, J. 2016. Scale- and taxon-dependent patterns of plant diversity in steppes of Khakassia, South Siberia (Russia). *Biodiversity and Conservation* 25: 2251–2273.
- Purschke, O. 2017. *oliverpurschke/Taxonomic\_Backbone: First release of the workflow to generate the taxonomic backbone for sPlot v.2.1 and TRY v.3.0*. DOI: [10.5281/zenodo.845445](https://doi.org/10.5281/zenodo.845445).
- R Core Team 2017. *R: A language and environment for statistical computing*. Vienna, AT. URL: <https://www.R-project.org/>.
- Reed, R.A., Peet, R.K., Palmer, M.W. & White, P.S. 1993. Scale dependence of vegetation-environment correlations: A case study of a North Carolina piedmont woodland. *Journal of Vegetation Science* 4: 329–340.
- Schaminée, J.H.J., Hennekens, S.M., Chytrý, M. & Rodwell, J.S. 2009. Vegetation-plot data and databases in Europe: an overview. *Preslia* 81: 173–185.
- Shmida, A. 1984. Whittaker’s plant diversity sampling method. *Israel Journal of Botany* 33: 41–46.
- Shmida, A. & Wilson, M.V. 1985. Biological determinants of species diversity. *Journal of Biogeography* 12: 1–20.



- Siefert, A., Ravenscroft, C., Althoff, D., Alvarez-Yépiz, J.C., Carter, B.E., Glennon, K.L., Heberling, J.M., Jo, I.S., Pontes, A., (...) & Fridley, J.D. 2012. Scale dependence of vegetation-environment relationships: a meta-analysis of multivariate data. *Journal of Vegetation Science* 23: 942–951.
- Török, P. & Dengler, J. in press. Palaearctic grasslands in transition: overarching patterns and future prospects. In: Squires, V.R., Dengler, J., Feng, H. & Hua, L. (eds.) *Grasslands of the world: diversity, management and conservation*. CRC Press, Boca Raton, US.
- Turtureanu, P.D., Palpurina, S., Becker, T., Dolnik, C., Ruprecht, E., Sutcliffe, L.M.E., Szabó, A. & Dengler, J. 2014. Scale- and taxon-dependent biodiversity patterns of dry grassland vegetation in Transylvania (Romania). *Agriculture, Ecosystems & Environment* 182: 15–24.
- Verheyen, K., De Frenne, P., Baeten, L., Waller, D.M., Hédli, R., Perring, M.P., Blondeel, H., Brunet, J., Chudomelová, M., ... & Bernhardt-Römermann, M. 2017. Combining biodiversity resurveys across regions to advance global change research. *BioScience* 67: 73–83.
- Vrahnakis, M.S., Janišová, M., Rūsiņa, S., Török, P., Venn, S. & Dengler, J. 2013. The European Dry Grassland Group (EDGG): stewarding Europe's most diverse habitat type. In: Baumbach, H. & Pfützenreuter, S. (eds.) *Steppenlebensräume Europas – Gefährdung, Erhaltungsmaßnahmen und Schutz*, pp. 417–434. Thüringer Ministerium für Landwirtschaft, Forsten, Umwelt und Naturschutz, Erfurt, DE.
- Wagner, V., Chytrý, M., Jiménez-Alfaro, B., Pergl, J., Hennekens, S., Biurrun, I., Knollová, I., Berg, C., Vassilev, K., (...) & Pyšek, P. 2017. Alien plant invasions in European woodlands. *Diversity and Distributions* 23: 969–981.
- Wamelink, G.W.W., van Adrichem, M.H.C., van Dobben, H.F., Frissel, J.Y., den Held, M., Joosten, V., Malinowska, A.H., Slim, P.A. & Wegman, R.M.A. 2012. Vegetation relevés and soil measurements in the Netherlands: the Ecological Conditions Database (EC). *Biodiversity & Ecology* 4: 125–132.
- Wilson, J.B., Peet, R.K., Dengler, J. & Pärtel, M. 2012. Plant species richness: the world records. *Journal of Vegetation Science* 23: 796–802.
- Wiser, S.K. 2016. Achievements and challenges in the integration, reuse and synthesis of vegetation plot data. *Journal of Vegetation Science* 27: 868–879.

## Author addresses

**Dengler, J.** (Corresponding author, juergen.dengler@zhaw.ch)<sup>1,2,3</sup>, **Wagner, V.** (viktor.wagner@ualberta.ca)<sup>4</sup>, **Dembicz, I.** (i.dembicz@biol.uw.edu.pl)<sup>5</sup>, **Garcia-Mijangos, I.** (itziar.garcia@ehu.es)<sup>6</sup>, **Naqinezhad, A.** (a.naqinezhad@umz.ac.ir)<sup>7</sup>, **Boch, S.** (steffen.boch@wsl.ch)<sup>8</sup>, **Chiarucci, A.** (alessandro.chiarucci@unibo.it)<sup>9</sup>, **Conradi, T.** (timo.conradi@uni-bayreuth.de)<sup>2,10</sup>, **Filibeck, G.** (filibeck@unitus.it)<sup>11</sup>, **Guarino, R.** (guarintrotto@hotmail.com)<sup>12</sup>, **Janišová, M.** (monika.janisova@gmail.com)<sup>13</sup>, **Steinbauer, M.J.** (manuel.steinbauer@fau.de)<sup>14</sup>, **Ačić, S.** (acic@agrif.bg.ac.rs)<sup>15</sup>, **Acosta, A.T.R.** (acosta@uniroma3.it)<sup>16</sup>, **Akasaka, M.** (muuak@cc.tuat.ac.jp)<sup>17</sup>, **Allers, M.-A.** (marc-

andre@posteo.de)<sup>18</sup>, **Apostolova, I.** (iva.apostolova@gmail.com)<sup>19</sup>, **Axmanová, I.** (axmanova@sci.muni.cz)<sup>20</sup>, **Bakan, B.** (branko.bakan@um.si)<sup>21</sup>, **Baranova, A.** (alina.baranova@uni-hamburg.de)<sup>22</sup>, **Bardy-Durchhalter, M.** (manfred.bardy-durchhalter@oeaw.ac.at)<sup>23</sup>, **Bartha, S.** (bartha.sandor@okologia.mta.hu)<sup>24</sup>, **Baumann, E.** (esther.baumann@uni-bayreuth.de)<sup>25</sup>, **Becker, T.** (beckerth@uni-trier.de)<sup>26</sup>, **Becker, U.** (beckeru@uni-mainz.de)<sup>27</sup>, **Belonovskaya, E.** (ebelonovskaya.0709@gmail.com)<sup>28</sup>, **Bengtsson, K.** (karin.bengtsson@ebc.uu.se)<sup>29</sup>, **Benito Alonso, J.L.** (jolube@jolube.net)<sup>30</sup>, **Berastegi, A.** (aberastg@gan-nik.es)<sup>31</sup>, **Bergamini, A.** (ariel.bergamini@wsl.ch)<sup>8</sup>, **Bonini, I.** (ilaria.bonini@unisi.it)<sup>32</sup>, **Bruun, H.H.** (hhbruun@bio.ku.dk)<sup>33</sup>, **Budzhak, V.** (budzhakv@gmail.com)<sup>34</sup>, **Bueno, A.** (abueno@uniovi.es)<sup>35</sup>, **Campos, J.A.** (juan.campos@ehu.es)<sup>6</sup>, **Cancellieri, L.** (cancellieri@unitus.it)<sup>11</sup>, **Carboni, M.** (marta.carboni@gmx.net)<sup>36</sup>, **Chocarro, C.** (chocarro@pvcf.udl.es)<sup>37</sup>, **Conti, L.** (luisa.conti@gmail.com)<sup>16</sup>, **Czarniecka-Wiera, M.** (m.czarniecka86@gmail.com)<sup>38</sup>, **De Frenne, P.** (pieter.defrenne@UGent.be)<sup>39</sup>, **Deák, B.** (debalazs@gmail.com)<sup>40</sup>, **Didukh, Y.P.** (ya.didukh@gmail.com)<sup>41</sup>, **Diekmann, M.** (mdiekman@uni-bremen.de)<sup>42</sup>, **Dolnik, C.** (cdolnik@ecology.uni-kiel.de)<sup>43</sup>, **Dupré, C.** (dupre@uni-bremen.de)<sup>42</sup>, **Ecker, K.** (klaus.ecker@wsl.ch)<sup>8</sup>, **Ermakov, N.** (brunnera@mail.ru)<sup>44</sup>, **Erschbamer, B.** (brigitta.erschbamer@uibk.ac.at)<sup>45</sup>, **Escudero, A.** (adrian.escudero@urjc.es)<sup>46</sup>, **Etayo, J.** (jetayosa@educacion.navarra.es)<sup>47</sup>, **Fajmonová, Z.** (zuzana.fajmonova@ibot.cas.cz)<sup>48</sup>, **Felde, V.A.** (vivian.felde@uib.no)<sup>49</sup>, **Fernández Calzado, M.R.** (rosafcalzado@gmail.com)<sup>50</sup>, **Finckh, M.** (mfinckh@googlemail.com)<sup>51</sup>, **Fotiadis, G.** (gfotiad95@gmail.com)<sup>52</sup>, **Fracchiolla, M.** (mariano.fracchiolla@uniba.it)<sup>53</sup>, **Ganeva, A.** (animoss@bio.bas.bg)<sup>19</sup>, **García-Magro, D.** (danigarcia1985@hotmail.com)<sup>6</sup>, **Gavilán, R.G.** (rgavilan@ucm.es)<sup>54</sup>, **Germany, M.** (mgermany@ecology.uni-kiel.de)<sup>55</sup>, **Giladi, I.** (itushgi@bgu.ac.il)<sup>56</sup>, **Gillet, F.** (francois.gillet@univ-fcomte.fr)<sup>57</sup>, **Giusso del Galdo, G.P.** (g.giusso@unict.it)<sup>58</sup>, **González, J.M.** (jose.gonzalez@urjc.es)<sup>46</sup>, **Grytnes, J.-A.** (jon.grytnes@uib.no)<sup>49</sup>, **Hájek, M.** (hajek@sci.muni.cz)<sup>20</sup>, **Hájková, P.** (buriana@sci.muni.cz)<sup>20</sup>, **Helm, A.** (aveliina.helm@ut.ee)<sup>59</sup>, **Herrera, M.** (meme.herrera@ehu.eus)<sup>6</sup>, **Hettenbergerová, E.** (eva.hette@centrum.cz)<sup>20</sup>, **Hobohm, C.** (hobohm@uni-flensburg.de)<sup>60</sup>, **Hüllbusch, E.M.** (elisabeth.huellbusch@uni-bayreuth.de)<sup>2</sup>, **Ingerpuu, N.** (nele.ingerpuu@ut.ee)<sup>59</sup>, **Jandt, U.** (ute.jandt@botanik.uni-halle.de)<sup>61</sup>, **Jeltsch, F.** (jeltsch@uni-potsdam.de)<sup>62</sup>, **Jensen, K.** (kai.jensen@uni-hamburg.de)<sup>63</sup>, **Jentsch, A.** (anke.jentsch@uni-bayreuth.de)<sup>64</sup>, **Jeschke, M.** (michael\_jeschke@hotmail.com)<sup>65</sup>, **Jiménez-Alfaro, B.** (borja.jimenez-alfaro@botanik.uni-halle.de)<sup>66</sup>, **Kącki, Z.**



(zygmunt.kacki@uwr.edu.pl)<sup>38</sup>, **Kakinuma, K.** (k.kakinuma0214@gmail.com)<sup>67</sup>, **Kapfer, J.** (jutta.kapfer@nibio.no)<sup>68</sup>, **Kavgaci, A.** (alickavgaci1977@yahoo.com)<sup>69</sup>, **Kelemen, A.** (kelemen.andras12@gmail.com)<sup>40</sup>, **Kiehl, K.** (k.kiehl@hs-osnabrueck.de)<sup>70</sup>, **Koyama, A.** (asukoyama@gmail.com)<sup>71</sup>, **Koyanagi, T.F.** (koya23jp@u-gakugei.ac.jp)<sup>72</sup>, **Kozub, Ł.** (kozub.lukasz@gmail.com)<sup>5</sup>, **Kuzemko, A.** (anyameadow.ak@gmail.com)<sup>41</sup>, **Kyrkjeeide, M.O.** (magni.kyrkjeeide@nina.no)<sup>73</sup>, **Landi, S.** (landsara@gmail.com)<sup>74</sup>, **Langer, N.** (cavalry@web.de)<sup>75</sup>, **Lastrucci, L.** (lastruccilorenzo73@gmail.com)<sup>53</sup>, **Lazzaro, L.** (lorenzo.lazzaro@unifi.it)<sup>53</sup>, **Lelli, C.** (chiara.elli6@gmail.com)<sup>9</sup>, **Lepš, J.** (suspa@prf.jcu.cz)<sup>76</sup>, **Löbel, S.** (s.loebel@tu-braunschweig.de)<sup>77</sup>, **Luzuriaga, A.L.** (arantzazu.lopezdeluzuriaga@urjc.es)<sup>46</sup>, **Maccherini, S.** (simona.maccherini@unisi.it)<sup>32</sup>, **Magnes, M.** (martin.magnes@uni-graz.at)<sup>78</sup>, **Malicki, M.** (malickimarek@interia.pl)<sup>79</sup>, **Marcenò, C.** (marceno.corrado@ehu.eus)<sup>6</sup>, **Mardari, C.** (constantin.mardari@uaic.ro)<sup>80</sup>, **Mauchamp, L.** (lesliemauchamp@gmail.com)<sup>57</sup>, **May, F.** (felix.may@ufz.de)<sup>81</sup>, **Michelsen, O.** (ottar.michelsen@ntnu.no)<sup>82</sup>, **Molero Mesa, J.** (jmolero@ugr.es)<sup>50</sup>, **Molnár, Z.** (molnar.zsolt@okologia.mta.hu)<sup>24</sup>, **Moysiyenko, I.Y.** (ivan.moysiyenko@gmail.com)<sup>83</sup>, **Nakaga, Y.K.** (y.nagata621@gmail.com)<sup>84</sup>, **Natcheva, R.** (raynanatcheva@yahoo.com)<sup>19</sup>, **Noroozi, J.** (noroozi.jalil@gmail.com)<sup>85</sup>, **Pakeman, R.J.** (robin.pakeman@hutton.ac.uk)<sup>86</sup>, **Palpurina, S.** (salza.palpurina@gmail.com)<sup>20</sup>, **Pärtel, M.** (meelis.partel@ut.ee)<sup>59</sup>, **Pätsch, R.** (ricarda.paetsch@gmail.com)<sup>87</sup>, **Pauli, H.** (harald.pauli@boku.ac.at)<sup>23</sup>, **Pedashenko, H.** (hristo\_pedashenko@yahoo.com)<sup>19</sup>, **Peet, R.K.** (peet@unc.edu)<sup>88</sup>, **Pielech, R.** (remekpielech@gmail.com)<sup>89</sup>, **Pipenbaher, N.** (natasia.pipenbaher@um.si)<sup>21</sup>, **Pirini, C.** (chpirini@bio.auth.gr)<sup>90</sup>, **Plesková, Z.** (pleskovicova@gmail.com)<sup>20</sup>, **Polyakova, M.A.** (galatella@mail.ru)<sup>44</sup>, **Prentice, H.C.** (honor\_c.prentice@biol.lu.se)<sup>91</sup>, **Reinecke, J.** (jennifer.reinecke@senckenberg.de)<sup>92</sup>, **Reitalu, T.** (triinreitalu@gmail.com)<sup>93</sup>, **Rodríguez-Rojo, M.P.** (mpilar.rodriguez@uclm.es)<sup>94</sup>, **Roleček, J.** (honza.rolecek@centrum.cz)<sup>48,20</sup>, **Ronkin, V.** (ronkinvl@discover-ua.com)<sup>95</sup>, **Rosati, L.** (leonardo.rosati@unibas.it)<sup>96</sup>, **Rosén, E.** (eje.rosen@gmail.com)<sup>97</sup>, **Ruprecht, E.** (eszter.ruprecht@ubbcluj.ro)<sup>98</sup>, **Rusina, S.** (rusina@lu.lv)<sup>99</sup>, **Sabovljević, M.** (marko@bio.bg.ac.rs)<sup>100</sup>, **Sánchez, A.M.** (ana.sanchez@urjc.es)<sup>46</sup>, **Savchenko, G.** (savchgala5@gmail.com)<sup>95</sup>, **Schuhmacher, O.** (schuhmacher@nabu-hamburg.de)<sup>101</sup>, **Škornik, S.** (sonja.skornik@um.si)<sup>21</sup>, **Sperandii, M.G.** (martagaia.sperandii@uniroma3.it)<sup>16</sup>, **Staniaszek-Kik, M.** (kik@biol.uni.lodz.pl)<sup>102</sup>, **Stevanović-Dajić, Z.** (dajic@agrif.bg.ac.rs)<sup>15</sup>, **Stock, M.** (martin.stock@lkn.landsh.de)<sup>103</sup>, **Suchrow, S.** (ssuchrow@web.de)<sup>63</sup>, **Sutcliffe, L.M.E.** (sutcliffe.laura@gmail.com)<sup>104</sup>, **Swacha, G.** (gswacha@gmail.com)<sup>38</sup>, **Sykes, M.**

(mtsnykes999@gmail.com)<sup>105</sup>, **Szabó, A.** (annuc19@gmail.com)<sup>106</sup>, **Talebi, A.** (amirtalebi@khayam.ut.ac.ir)<sup>107</sup>, **Tănase, C.** (tanase@uaic.ro)<sup>108</sup>, **Terzi, M.** (massimo.terzi@ibbr.cnr.it)<sup>109</sup>, **Tölgyesi, C.** (festuca7@yahoo.com)<sup>110</sup>, **Torca, M.** (marta.torca@ehu.es)<sup>6</sup>, **Török, P.** (molinia@gmail.com)<sup>111</sup>, **Tóthmérész, B.** (tothmerb@gmail.com)<sup>111</sup>, **Tsarevskaya, N.** (ngtsar@yandex.ru)<sup>28</sup>, **Tsiripidis, I.** (tsiripid@bio.auth.gr)<sup>90</sup>, **Tzonev, R.** (rossentzonev@abv.bg)<sup>112</sup>, **Ushimaru, A.** (ushimaru@kobe-u.ac.jp)<sup>84</sup>, **Valkó, O.** (valkoorsi@gmail.com)<sup>40</sup>, **van der Maarel, E.** (eddy.arteco@planet.nl)<sup>113</sup>, **Vanneste, T.** (thomas.vanneste@ugent.be)<sup>39</sup>, **Vashenyak, I.** (arrhenatherum@gmail.com)<sup>114</sup>, **Vassilev, K.** (kiril5914@abv.bg)<sup>19</sup>, **Viciani, D.** (daniele.viciani@unifi.it)<sup>53</sup>, **Villar, L.** (lvillar@ipe.csic.es)<sup>115</sup>, **Virtanen, R.** (risto.virtanen@oulu.fi)<sup>116</sup>, **Vitasović Kosić, I.** (ivitasovic@agr.hr)<sup>117</sup>, **Wang, Y.** (yunwang.hh@gmail.com)<sup>92</sup>, **Weiser, F.** (frank.weiser@uni-bayreuth.de)<sup>25</sup>, **Went, J.** (juliawent@gmx.de)<sup>2</sup>, **Wesche, K.** (karsten.wesche@senckenberg.de)<sup>92,3</sup>, **White, H.** (hannah.white@ucd.ie)<sup>118</sup>, **Winkler, M.** (manuela.winkler@boku.ac.at)<sup>23</sup>, **Zaniewski, P.T.** (piotr.zaniewski@wl.sggw.pl)<sup>119</sup>, **Zhang, H.** (zhanghuitianxia@163.com)<sup>120</sup>, **Ziv, Y.** (yziv@bgu.ac.il)<sup>121</sup>, **Znamenskiy, S.** (seznam@krc.karelia.ru)<sup>122</sup> & **Biurrun, I.** (idoia.biurrun@ehu.es)<sup>6</sup>

<sup>1</sup> Research Group Vegetation Ecology, Institute of Natural Resource Sciences (IUNR), Zurich University of Applied Sciences (ZHAW), Grüentalstr. 14, Postfach, 8820 Wädenswil, Switzerland

<sup>2</sup> Plant Ecology, Bayreuth Center of Ecology and Environmental Research (BayCEER), University of Bayreuth, Universitätsstr. 30, 95447 Bayreuth, Germany

<sup>3</sup> German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany

<sup>4</sup> Department of Biological Sciences, University of Alberta, Edmonton, AB, T6G 2R3, Canada

<sup>5</sup> Department of Plant Ecology and Environmental Conservation, Faculty of Biology, University of Warsaw, ul. Żwirki i Wigury 101, 02-089 Warsaw, Poland

<sup>6</sup> Department of Plant Biology and Ecology, University of the Basque Country UPV/EHU, P.O. Box 644, 48080 Bilbao, Spain

<sup>7</sup> Department of Biology, Faculty of Basic Sciences, University of Mazandaran, Babolsar, P.O. Box 47416-95447, Mazandaran, Iran.

<sup>8</sup> Swiss Federal Research Institute WSL, Zürcherstrasse 111, 8903 Birmensdorf, Switzerland

<sup>9</sup> Department of Biological, Geological and Environmental Sciences, University of Bologna, via Irnerio 42, 40126 Bologna, Italy

- <sup>10</sup> Ecoinformatics and Biodiversity, Department of Bioscience, Aarhus University, 8000 Aarhus, Denmark
- <sup>11</sup> Department of Agricultural and Forestry Science (DAFNE), University of Tuscia, 01100 Viterbo, Italy
- <sup>12</sup> Department STEBICEF – Botanical Unit, University of Palermo, via Archiarafi 38, 90123 Palermo, Italy
- <sup>13</sup> Institute of Botany, Plant Science and Biodiversity Center, Slovak Academy of Sciences, Ďumbierska 1, 974 11 Banská Bystrica, Slovakia
- <sup>14</sup> GeoZentrum Nordbayern, Department of Geography and Geosciences, Friedrich-Alexander University Erlangen-Nürnberg (FAU), Loewenichstr. 28, 91054 Erlangen, Germany
- <sup>15</sup> Department of Botany, Faculty of Agriculture, University of Belgrade, Nemanjina 6, 11080 Belgrade-Zemun, Serbia
- <sup>16</sup> Dipartimento di Scienze, Università degli Studi di Roma Tre, Viale Marconi, 446, 00146 Roma, Italy
- <sup>17</sup> Institute of Agriculture, Tokyo University of Agriculture and Technology, Fuchu, 183-8509 Tokyo, Japan
- <sup>18</sup> Albert-Einstein-Str. 11a, 14473 Potsdam, Germany
- <sup>19</sup> Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, 23 Acad. Georgi Bonchev str., 1113 Sofia, Bulgaria
- <sup>20</sup> Department of Botany and Zoology, Masaryk University, Kotlářská 2, 61137 Brno, Czech Republic
- <sup>21</sup> Biology Department, Faculty of Natural Sciences and Mathematics, University of Maribor, Koroska cesta 160, 2000 Maribor, Slovenia
- <sup>22</sup> CEN Center for Earth System Research and Sustainability, University of Hamburg, Bundesstr. 55, 20146 Hamburg, Germany
- <sup>23</sup> GLORIA co-ordination, Institute for Interdisciplinary Mountain Research, Austrian Academy of Sciences, Silbergasse 30/3, 1190 Vienna, Austria
- <sup>24</sup> Institute of Ecology and Botany, MTA Centre for Ecological Research, Alkotmány u. 2., 2163 Vácrátót, Hungary
- <sup>25</sup> Biogeography, University of Bayreuth, Universitätsstr. 30, 95447 Bayreuth, Germany
- <sup>26</sup> Geobotany, Regional and Environmental Science, University of Trier, Behringstr. 21, 54296 Trier, Germany

- <sup>27</sup> Grüne Schule im Botanischen Garten, Johannes Gutenberg Universität Mainz, Anselm-Franz-von-Bentzel-Weg 9b, 55128 Mainz, Germany
- <sup>28</sup> Institute of Geography, Russian Academy of Sciences, Staromonetny per., 29, 119017 Moscow, Russia
- <sup>29</sup> Department of Ecology and Genetics (EBC), Uppsala University, Campus Gotland, 62167 Visby, Sweden
- <sup>30</sup> JOLUBE Consultor Botánico, C/ Mariano Rguez. De Ledesma 4-3ºA, 22700 Jaca, Spain
- <sup>31</sup> Biodiversity Team, Gestión Ambiental de Navarra, S.A., Padre Adoain 219, Bajo, 31015 Pamplona, Spain
- <sup>32</sup> Life Sciences, University of Siena, P.A. Mattioli, 4, 53100 Siena, Italy
- <sup>33</sup> Section for Ecology and Evolution, Department of Biology, University of Copenhagen, Universitetsparken 15, 2100 Copenhagen, Denmark
- <sup>34</sup> Botany and Natural Protection Department, Chernivtsi National University, Fedkovich Street 11, 58022 Chernivtsy, Ukraine
- <sup>35</sup> Instituto de Recursos Naturales y Ordenación del Territorio (INDUROT), Universidad de Oviedo, Campus de Mieres. Edificio de Investigación, 33600 Mieres, Spain
- <sup>36</sup> Laboratoire d'Ecologie Alpine, UMR-CNRS. 5553, Université Grenoble Alpes, BP 53, 38041 Grenoble Cedex 9, France
- <sup>37</sup> Crop and Forest Science, Universitat de Lleida, Rovira Roure 177, 25110 Lleida, Spain
- <sup>38</sup> Botanical Garden, University of Wrocław, Sienkiewicza 23, 50-335 Wrocław, Poland
- <sup>39</sup> Forest & Nature Lab, Ghent University, Geraardsbergsesteenweg 267, 9090 Gontrode, Belgium
- <sup>40</sup> Department of Ecology, University of Debrecen, Egyetem tér 1, 4032 Debrecen, Hungary
- <sup>41</sup> Geobotany and Ecology Department, M.G. Kholodny Institute of Botany NAS of Ukraine, Tereschenkivska str. 2, 1601 Kyiv, Ukraine
- <sup>42</sup> Vegetationsökologie und Naturschutzbiologie, FB 2, Universität Bremen, Leobener Str. 5, 28359 Bremen, Germany
- <sup>43</sup> Ecology Centre Kiel, Kiel University, Olshausenstr. 40, 24098 Kiel, Germany
- <sup>44</sup> Central Siberian Botanical Garden, Russian Academy of Science, Zolotodolinskaya 101, 630090 Novosibirsk, Russia
- <sup>45</sup> Department of Botany, University of Innsbruck, Sternwartestr. 15, 6020 Innsbruck, Austria
- <sup>46</sup> Department of Biology, Geology, Physics and Inorganic Chemistry, Universidad Rey Juan Carlos, Tulipán s/n, 28933 Móstoles, Spain
- <sup>47</sup> Navarro Villoslada 16-3ºdcha, 31003 Pamplona, Spain

- <sup>48</sup> Department of Vegetation Ecology, Institute of Botany, Czech Academy of Sciences, Lidická 25/27, 60200 Brno, Czech Republic
- <sup>49</sup> Department of Biology, University of Bergen, Postbox 7803, 5020 Bergen, Norway
- <sup>50</sup> Department of Botany, Faculty of Pharmacy, University of Granada, Campus de Cartuja s/n, 18071 Granada, Spain
- <sup>51</sup> Biodiversity, Evolution and Ecology of Plants (BEE), Biocentre Klein Flottbek and Botanical Garden, University of Hamburg, Ohnhorststr. 18, 22609 Hamburg, Germany
- <sup>52</sup> Department of Forestry and Natural Environment Management, TEI (Technological Education Institute) of Sterea Ellada, Dimokratias 3, 36100 Karpenisi, Greece
- <sup>53</sup> Department of Agricultural and Environmental Science, University of Bari, Via Orabona 4, 70126 Bari, Italy
- <sup>54</sup> Departamento de Biología Vegetal II, Facultad de Farmacia, Universidad Complutense de Madrid, 28040 Madrid, Spain
- <sup>55</sup> Institute for Ecosystem Research, Kiel University, Olshausenstr. 75, 24118 Kiel, Germany
- <sup>56</sup> Mitrani Department of Desert Ecology, Swiss Institute for Dryland Environmental and Energy Research, Jacob Blaustein Institutes for Desert Research, Ben-Gurion University of the Negev, Sede Boqer Campus, 84990 Midreshet Ben Gurion, Israel
- <sup>57</sup> UMR Chrono-environnement, Université de Franche-Comté, 16 route de Gray, 25030 Besançon, France
- <sup>58</sup> Department of Biological, Geological and Environmental Sciences (BiGeA), University of Catania, via A. Longo 19, 95125 Catania, Italy
- <sup>59</sup> Institute of Ecology and Earth Sciences, University of Tartu, Lai 40, 51005 Tartu, Estonia
- <sup>60</sup> Interdisciplinary Institute of Environmental, Social and Human Sciences, University of Flensburg, Auf dem Campus 1, 24943 Flensburg, Germany
- <sup>61</sup> Geobotany and Botanical Garden, Institute of Biology, Martin Luther University, Am Kirchtor 1, 6108 Halle (Saale), Germany
- <sup>62</sup> Plant Ecology and Nature Conservation, University of Potsdam, Am Mühlenberg 3, 14476 Potsdam, Germany
- <sup>63</sup> Applied Plant Ecology, Biocentre Klein Flottbek, University of Hamburg, Ohnhorststr. 18, 22609 Hamburg, Germany
- <sup>64</sup> Disturbance Ecology, Bayreuth Center of Ecology and Environmental Research (BayCEER), University of Bayreuth, Universitätsstr. 30, 95447 Bayreuth, Germany
- <sup>65</sup> Friesenstr. 47, 82223 Eichenau, Germany

- <sup>66</sup> Research Unit of Biodiversity (CSIC, UO, PA), Oviedo University, Campus de Mieres, 33600 Mieres, Spain
- <sup>67</sup> NASA Goddard Institute for Space Studies, Columbia University, 2880 Broadway, NY 10025 New York, United States
- <sup>68</sup> Division for Geography and Statistics, Department of Landscape Monitoring, Norwegian Institute of Bioeconomy Research, Holtveien 66, 9269 Tromsø, Norway
- <sup>69</sup> Southwest Anatolia Forest Research Institute, POB 264, 07002 Antalya, Turkey
- <sup>70</sup> Vegetation Ecology and Botany, Faculty of Agricultural Sciences and Landscape Architecture, University of Applied Sciences Osnabrück, Oldenburger Landstr. 24, 49090 Osnabrück, Germany
- <sup>71</sup> Institute for Sustainable Agro-ecosystem Services, University of Tokyo, 1-1-1, Midori-cho, Nishi-Tokyo, 188-0002 Tokyo, Japan
- <sup>72</sup> Field Studies Institute for Environmental Education, Tokyo Gakugei University, 4-1-1 Koganei, 184-8501 Tokyo, Japan
- <sup>73</sup> Norwegian Institute for Nature Research, 7485 Trondheim, Norway
- <sup>74</sup> Department of Sciences of Nature and Territory, University of Sassari, Via Piandanna, 07100 Sassari, Italy
- <sup>75</sup> Zum Schwärzensee 27, 16227 Eberswalde, Germany
- <sup>76</sup> Department of Botany, Faculty of Science, University of South Bohemia, Branišovská 31, 370 05 Ceske Budejovice, Czech Republic
- <sup>77</sup> Landscape Ecology and Environmental Systems Analysis, Institute of Geoecology, Langer Kamp 19c, 38106 Braunschweig, Germany
- <sup>78</sup> Institute of Plant Sciences, University of Graz, Holteigasse 6, 8010 Graz, Austria
- <sup>79</sup> Department of Botany, University of Wrocław, ul. Kanonia 6/8, 50-328 Wrocław, Poland
- <sup>80</sup> Anastasie Fatu Botanical Garden, Alexandru Ioan Cuza University from Iași, Dumbrava Roșie 7-9, 700487 Iași, Romania
- <sup>81</sup> Biodiversity Synthesis Research Group, German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany
- <sup>82</sup> NTNU Sustainability, Norwegian University of Science and Technology, 7491 Trondheim, Norway
- <sup>83</sup> Department of Botany, Kherson State University, ul. Universytetska 27, 73000 Kherson, Ukraine
- <sup>84</sup> Graduate School of Human Development and Environment, Kobe University, 3-11 Tsurukabuto, 657-8501 Kobe, Japan

- <sup>85</sup> Department of Botany and Biodiversity Research, University of Vienna, Rennweg 14, 1030 Vienna, Austria
- <sup>86</sup> Ecological Sciences, The James Hutton Institute, Craigiebuckler, AB15 8QH Aberdeen, United Kingdom
- <sup>87</sup> Vegetation and Phytodiversity Analysis, Georg-August University of Göttingen, Untere Karspüle 2, 37073 Göttingen, Germany
- <sup>88</sup> Department of Biology, University of North Carolina, 27599-3280 Chapel Hill, NC, United States
- <sup>89</sup> Department of Forest Biodiversity, Faculty of Forestry, University of Agriculture in Kraków, al. 29 Listopada 46, 31-425 Kraków, Poland
- <sup>90</sup> Department of Botany, School of Biology, Aristotle University of Thessaloniki, 54124 Thessaloniki, Greece
- <sup>91</sup> Department of Biology, Lund University, Ecology Building, 22362 Lund, Sweden
- <sup>92</sup> Botany Department, Senckenberg Museum of Natural History Görlitz, Am Museum 1, 2826 Görlitz, Germany
- <sup>93</sup> Institute of Geology, Tallinn University of Technology, Ehitajate tee 5, 19086 Tallinn, Estonia
- <sup>94</sup> Institute of Environmental Sciences, University of Castilla-La Mancha, Avenida Carlos III, s/n, 45071 Toledo, Spain
- <sup>95</sup> Dept of Zoology and Ecology, Kharkiv National University, 4 Svobody Sq, 61022 Kharkiv, Ukraine
- <sup>96</sup> School of Agricultural, Forest, Food and Environmental Sciences, University of Basilicata, Via Ateneo Lucano 10, 85100 Potenza, Italy
- <sup>97</sup> Department of Ecology and Genetics (EBC), Uppsala University, Norbyvägen 18 D, 75236 Uppsala, Sweden
- <sup>98</sup> Hungarian Department of Biology and Ecology, Babes-Bolyai University, Republici str. 42, 400015 Cluj-Napoca, Romania
- <sup>99</sup> Faculty of Geography and Earth Sciences, University of Latvia, 1 Jelgavas Street, 1004 Riga, Latvia
- <sup>100</sup> Institute for Botany and Botanical Garden "Jevremovac", Faculty of Biology, University of Belgrade, Takovska 43, 11000 Belgrade, Serbia
- <sup>101</sup> NABU Hamburg, Am Stadtbad 45, 29451 Dannenberg, Germany
- <sup>102</sup> Department of Geobotany and Plant Ecology, University of Łódź, Banacha 12/16, 90- 237 Łódź, Poland



- <sup>103</sup> Schleswig-Holstein Agency for Coastal Defence, National Park and Marine Conservation, National Park Authority, Schlossgarten 1, 25832 Tönning, Germany
- <sup>104</sup> Plant Ecology and Ecosystem Research, Georg-August University of Göttingen, Untere Karspüle 2, 37073 Göttingen, Germany
- <sup>105</sup> Department of Physical Geography and Ecosystem Science, Lund University, Sölvegatan 12, 22362 Lund, Sweden
- <sup>106</sup> Romanian Ornithological Society, Gh. Dima street 49, 400342 Cluj-Napoca, Romania
- <sup>107</sup> School of Biology and Center of Excellence in Phylogeny of Living Organisms, College of Science, University of Tehran, Enghelab, 14155-6455 Tehran, Iran
- <sup>108</sup> Faculty of Biology, Alexandru Ioan Cuza University, Carol I 20A, 700505 Iași, Romania
- <sup>109</sup> Institute of Biosciences and Bioresources (IBBR), Italian National Council of Research (CNR), Via Amendola 165/A, 70126 Bari, Italy
- <sup>110</sup> Department of Ecology, University of Szeged, Közép fasor 52, 6726 Szeged, Hungary
- <sup>111</sup> MTA-DE Lendület Functional and Restoration Ecology Research Group, Egyetem tér 1, 4032 Debrecen, Hungary
- <sup>112</sup> Department of Ecology and Environmental Protection, Sofia University ‘St. Kliment Ohridski’, 8 Dragan Tzankov Blvd., 1164 Sofia, Bulgaria
- <sup>113</sup> Centre for Ecological and Evolutionary Studies, University of Groningen, Linnaeusborg - Nijenborgh 7 Building U, 9747 AG Groningen, Netherlands
- <sup>114</sup> Khmelnytskyi Institute of Interregional Academy of Personnel Management, Prospect Myru Str., 101A, 29015 Khmelnytskyi, Ukraine
- <sup>115</sup> Instituto Pirenaico de Ecología, Consejo Superior de Investigaciones Científicas, 22700 Jaca, Spain
- <sup>116</sup> Ecology & Genetics, University of Oulu, P.O Box 3000, 90014 Oulu, Finland
- <sup>117</sup> Faculty of Agriculture, Department of Agricultural Botany, University of Zagreb, Svetošimunska cesta 25, 10000 Zagreb, Croatia
- <sup>118</sup> School of Biology and Environmental Science, Earth Institute, University College Dublin, Belfield, Dublin 4, Ireland
- <sup>119</sup> Department of Forest Botany, Faculty of Forestry, Warsaw University of Life Sciences – SGGW, Nowoursynowska 159, 02-776 Warsaw, Poland
- <sup>120</sup> Key Laboratory of Vegetation Restoration and Management of Degraded Ecosystems, South China Botanical Garden, Chinese Academy of Sciences, Xingke Road 723, 510650 Guangzhou, China



<sup>121</sup> Department of Life Sciences, Ben-Gurion University of the Negev, 84105 Negev, Israel

<sup>122</sup> Institute of Biology, Karelian Research Centre RAS, Pushkinskaya 11, 185910 Petrozavodsk, Russia

## Electronic Supplements


Supplementary material associated with this article is embedded in the article's pdf. The online version of Phytocoenologia is hosted at [www.ingentaconnect.com/content/schweiz/phyt](http://www.ingentaconnect.com/content/schweiz/phyt) and the journal's website [www.schweizerbart.com/journals/phyto](http://www.schweizerbart.com/journals/phyto). The publisher does not bear any liability for the lack of usability or correctness of supplementary material.

Supplement S1. GrassPlot Bylaws.

Supplement S2. Overview of the datasets in GrassPlot 1.00.


Supplement S3. Bibliographic references to the datasets contained in GrassPlot 1.00.

Supplement S4. Overview of the content of the header data fields other than those in Tables 1–4 and Fig. 2.



**Fig. 1.** GrassPlot logo developed by Iwona Dembicz. It links the *Stipa* awns (reminiscent of the EDGG logo) to the multi-scale sampling approach of precisely delimited plots.





**Fig. 2.** Maps showing the spatial distribution of the plots contained in GrassPlot v. 1.00. Grey dots refer to plots of any size, while black dots indicate nested-plot series with at least four different grain sizes.

**Table. 1.** Overview of some key parameters of GrassPlot v. 1.00 in terms of access regime, quality of the data, methodological aspects as well as temporal and elevational distribution. The column “NA” indicates the fraction of plots in GrassPlot for which the respective field is currently not filled.

Parameter	NA	Frequency distribution of parameter values
<b>Availability of data</b>		
– Access regime	–	1 – restricted access (1.7%); 2 – semi-restricted access (98.3%); 3 – free access (0.0%)
– Availability of compositional data	–	Yes (97.7%); to be provided later (0.2%); no (2.1%)
<b>Methodological aspects</b>		
– Recording method	<0.1%	Shoot presence (87%); rooted presence (11.2%)
– Plot shape	–	Squares (75.3%); rectangles 1:2 (22.5%); rectangles 1:1.6 (0.5%); rectangles more elongated than 1:2 (< 0.1%); circles (1.6%)
– Accuracy of coordinates	0.4%	≤ 1 m (3.4%); 1.1–10 m (30.1%); 11–100 m (6.2%); 101–1,000 m (59.1%); > 1,000 m (0.7%)
<b>Distribution of plots</b>		
– Year of recording	–	Before 1980 (< 0.1%); 1980–1989 (2.4%); 1990–1999 (2.7%); 2000–2009 (79.1%); 2010 and later (15.7%)
– Elevation	3.9%	≤ 10 m a.s.l. (8.4%); 11–100 m a.s.l. (17.2%); 101–1,000 m a.s.l. (12.1%); 1,001–2,000 m a.s.l. (12.0%); 2,001–3,000 m a.s.l. (34.2%); 3,001–4,000 m a.s.l. (16.0%); > 4,000 m a.s.l. (< 0.1%)

**Table. 2.** Number of plots ( $N$ ) and the mean ( $S_{\text{mean}}$ ) and maximum ( $S_{\text{max}}$ ) richness in GrassPlot (v. 1.00) across different plot sizes, and for vascular plants and the complete terricolous vegetation (vascular plants, bryophytes and lichens), respectively. Non-standard plot sizes include all other plot sizes (which are collected only in case of nested-plot series). Note that due to different samples, maxima of bigger plot sizes could sometimes be lower than for smaller plot sizes or that maxima for complete terricolous vegetation could sometimes be lower than for vascular plants only. Information on plot size pairs, such as 10 m<sup>2</sup> and 9 m<sup>2</sup>, is combined in one line because based on species-area relationships with typical  $z$ -values between 0.15 and 0.30, the relative difference in richness would only be about 1.6–3.2%, i.e. negligible given the overall variability of the data.

Plot size	Vascular plants			Complete terricolous vegetation		
	$N$	$S_{\text{mean}}$	$S_{\text{max}}$	$N$	$S_{\text{mean}}$	$S_{\text{max}}$
0.0001 m <sup>2</sup>	2,206	1.9	11	1,540	2.0	10
0.001 or 0.0009 m <sup>2</sup>	3,344	3.3	19	1,481	3.3	19
0.01 m <sup>2</sup>	66,000	3.8	24	2,224	6.5	29
0.1 or 0.09 m <sup>2</sup>	3,737	11.7	43	1,496	10.3	46
1 m <sup>2</sup>	17,206	13.8	79	2,008	18.2	82
10 or 9 m <sup>2</sup>	5,520	31.0	98	2,016	34.1	101
100 m <sup>2</sup>	2,545	31.9	127	824	46.8	134
1,000 or 900 or 1,024 m <sup>2</sup>	181	47.2	134	45	59.1	123
Non-standard plot sizes	68,207			2,430		
Total	168,946			14,064		

**Table. 3.** The ten most represented phytosociological classes (according to Mucina et al. 2016) in GrassPlot 1.00, based on the numbers (*N*) and percentages of plots (%) in the total dataset.

Class	Group	<i>N</i>	%
<i>Festuco-Brometea</i>	Temperate dry grasslands	36,242	21.5%
<i>Festucetea indigestae</i>	Alpine grasslands	31,086	18.4%
<i>Juncetea trifidi</i>	Alpine grasslands	13,947	8.3%
<i>Carici rupestris-Kobresietea bellardii</i>	Alpine grasslands	10,958	6.5%
<i>Stipo-Trachynietea distachyae</i>	Mediterranean grasslands	6,697	4.0%
<i>Molinio-Arrhenatheretea</i>	Temperate mesic and wet grasslands	6,078	3.6%
<i>Koelerio-Corynephoretea canescentis</i>	Temperate dry grasslands	3,410	2.0%
<i>Ammophiletea</i>	Coastal grasslands	3,390	2.0%
<i>Juncetea maritimi</i>	Coastal grasslands	3,347	2.0%
<i>Helichryso-Crucianelletea maritimae</i>	Coastal grasslands	3,259	1.9%
Other classes		6,638	3.9%
Not yet assigned to a class		42,458	25.7%

**Table. 4.** Numbers ( $N$ ) and densities of plots per country (or dependent territory), sorted by decreasing density of plots per 100 km<sup>2</sup>. The twenty countries with the highest densities are given in the table. The remaining 16 countries can be found in the GIVD Fact Sheet. Area [km<sup>2</sup>] refers to the size of the respective territory.

Code	Country	Area [km <sup>2</sup> ]	$N$	$N / 100 \text{ km}^2$
AT	Austria	83,855	13,099	15.62
ES	Spain	504,790	54,608	10.82
IL	Israel	20,724	1,795	8.66
SE	Sweden	440,940	26,149	5.93
CH	Switzerland	41,285	2,307	5.59
IT	Italy	301,245	14,943	4.96
NO	Norway	323,758	12,717	3.93
HU	Hungary	93,030	3,648	3.92
EE	Estonia	45,100	1,578	3.50
DE	Germany	356,840	7,311	2.05
CZ	Czech Republic	78,864	1,111	1.41
UK	United Kingdom	244,587	2,886	1.18
PL	Poland	312,685	2,778	0.89
NL	Netherlands	41,160	354	0.86
SK	Slovakia	49,035	405	0.83
IR	Iran	1,648,000	12,992	0.79
RS	Serbia	77,453	493	0.64
BG	Bulgaria	110,910	572	0.52
SJ	Svalbard and Jan Mayen	61,397	280	0.46
RO	Romania	237,500	1,025	0.43

**Appendix 1.** Comparison of the three large supra-national databases of vegetation-plot data: EVA, sPlot and GrassPlot, indicating their similarities and differences (information as of 14 January 2018).

Aspect	EVA	sPlot	GrassPlot
<b>Scope</b>			
Geographic scope	Europe (+ Canary Islands, Turkey, Caucasus countries)	World	Palearctic biogeographic realm
Vegetation types included	All	All	Grasslands and other open habitats
Plot sizes	Any in the range 1–1,000 m <sup>2</sup> and also plots without reported size	Any in the range 1–10,000 m <sup>2</sup>	Eight standard grain sizes from 0.0001 to 1,000 m <sup>2</sup> (other sizes only if part of nested plot series)
Nested plots	Not supported	Not supported	Specialised in nested plots; information on hierarchy of nesting is stored
Delimitation of plots and comprehensiveness of sampling	No requirements	No requirements; even plots are included where only dominant species have been sampled (but this information is available)	Only plots that have been precisely delimited in the field and sampled comprehensively
<b>Data types and formats</b>			
Information contained in the database	Plots with compositional data	Plots with compositional data	Plots with compositional data or just richness data + metadata
Format in which the data are stored and provided	Turboveg 2 databases combined in a Turboveg 3 database	Turboveg 2 databases combined in a Turboveg 3 database; data provision as R Data.table with harmonized information	Spread sheet for richness, methodological and environmental data; long table format in R for compositional data
Matching with plant trait and phylogenetic data available	No (but in the future possible via collaboration with sPlot/TRY)	Yes	No
<b>Available information per plot</b>			
Recording of non-vascular plants	Rare and if available often not comprehensive; plots with comprehensive data cannot be extracted	Rare and if available often not comprehensive; plots with comprehensive data cannot be extracted	Often included and then comprehensive
Importance values of	Normally required (Br.-	Multitude of quantitative	Importance values (often



species	Bl., % or similar)	scales, but also presence-absence	%) or just presence-absence
Precision of plot coordinates	High to very low; field often not filled	High to very low	Mostly high
Environmental data measured in the plot	Not standardised	Not standardised	Standardised and thus directly usable
Names of plants provided	Standardised to an internal taxonomic backbone for Europe (SynBioSys Taxon Database), also taking into account different meanings of the same name in different floras	Harmonized with online tools, taking into account synonymy, but not different meanings of the same name in different floras	Harmonized with online tools, taking into account synonymy, but not different meanings of the same name in different floras
<b>Current content</b>			
Plot number	1,474,590	1,121,244	168,997
Countries covered	57	160	36
Spatial density of available plots	High	High in Europe, medium in parts of North America and Australia, sparse elsewhere	Relatively sparse
Overlap with the other databases in the table	The majority of EVA plots are also in sPlot	sPlot accepts European plots only via EVA	Overlap with EVA and sPlot is small and documented; it is recommended that plots that are suitable for EVA/sPlot and GrassPlot should be contributed twice
<b>Responsible working groups and their rules</b>			
Affiliated with	European Vegetation Survey (EVS)	German Centre for Integrative Biodiversity Research (iDiv)	Eurasian Dry Grassland Group (EDGG)
Website	<a href="http://euroveg.org/eva-database">http://euroveg.org/eva-database</a>	<a href="https://www.idiv.de/splot">https://www.idiv.de/splot</a>	<a href="http://bit.ly/2qKTQt2">http://bit.ly/2qKTQt2</a>
Governed by	7-head Coordinating Board	5-head Steering Committee	7-head Governing Board
Members	72 supranational, national and regional databases	110 supranational, national and regional databases, 2 continental data aggregators	192 owners of 126 regional datasets
Required offers of opt-in authorships for analytical papers	No requirement, usually one co-author for each database that contributed at least (5%) 10% of the final dataset	One opt-in co-author for each database used in the study	One opt-in co-author for each dataset that contributed at least 2% of the final dataset