

A characterization of n -associative, monotone, idempotent functions on an interval that have neutral elements

Gergely Kiss*

University of Luxembourg, Faculty of Science, Mathematical
Research Unit, rue Richard Coudenhove-Kalergi 6, L-1359
Luxembourg, gergely.kiss@uni.lu

Gábor Somlai †

Eötvös Loránd University, Faculty of Science, Institute of
Mathematics, Pázmány Péter sétány. 1/c, Budapest, H-1117
Hungary, gsomlai@cs.elte.hu

November 5, 2018

Abstract

We investigate monotone idempotent n -ary semigroups and provide a generalization of the Czogala–Drewniak Theorem, which describes the idempotent monotone associative functions having a neutral element. We also present a complete characterization of idempotent monotone n -associative functions on an interval that have neutral elements.

Keywords: quasitrivial, ordered semigroups, n -associativity, idempotency, monotonicity, neutral element

MSR 2010 classification: 06F05, 20M99

1 Introduction

A function $F: X^n \rightarrow X$ is called *n -associative* if for every $x_1, \dots, x_{2n-1} \in X$ and every $i = 1, \dots, n-1$, we have

$$\begin{aligned} F(F(x_1, \dots, x_n), x_{n+1}, \dots, x_{2n-1}) = \\ F(x_1, \dots, x_i, F(x_{i+1}, \dots, x_{i+n}), x_{i+n+1}, \dots, x_{2n-1}). \end{aligned} \quad (1)$$

*The first author was supported by the internal research project R-AGR-0500-MR03 of the University of Luxembourg and the Hungarian Scientific Research Fund (OTKA) K104178.

†The second author was supported by the Hungarian Scientific Research Fund (OTKA) K115799.

Throughout this paper we assume that the underlying sets of the algebraic structures under consideration are partially ordered sets (poset). Some of our results only work for totally ordered sets. In our main results we investigate n -ary semigroups on an arbitrary nonempty subinterval of the real numbers.

A set X endowed with an n -associative function $F: X^n \rightarrow X$ is called an *n -ary semigroup* and is denoted by (X, F_n) . Clearly, we obtain a generalization of associative functions, which are the 2-associative functions using our terminology.

The main purpose of this paper is to describe a class of n -ary semigroups. An n -ary semigroup is called *idempotent* if $F(a, \dots, a) = a$ for all $a \in X$. Another important property is the monotonicity. An n -associative function is called *monotone in the i -th variable* if for all fixed $a_1, \dots, a_{i-1}, a_{i+1}, \dots, a_n \in X$, the 1-variable functions $f_i(x) := F(a_1, \dots, a_{i-1}, x, a_{i+1}, \dots, a_n)$ is order-preserving or order-reversing. An n -associative function is called *monotone* if it is monotone in each of its variables. Further, we say that $e \in X$ is a *neutral element* for an n -associative function F if for every $x \in X$ and every $i = 1, \dots, n$, we have $F(e, \dots, e, x, e, \dots, e) = x$, where x is substituted for the i -th coordinate.

An important construction of n -ary semigroups is the following. Let (X, F_2) be a binary semigroup. Let $F_n := \underbrace{F_2 \circ F_2 \circ \dots \circ F_2}_{n-1}$, where

$$\begin{aligned} F_n(x_1, \dots, x_n) &= \underbrace{F_2 \circ F_2 \circ \dots \circ F_2}_{n-1}(x_1, \dots, x_n) \\ &= F_2(x_1, F_2(x_2, \dots, F_2(x_{n-1}, x_n))). \end{aligned}$$

We get an n -associative function $F_n: X^n \rightarrow X$ and an n -ary semigroup (X, F_n) . In this case we say that (X, F_n) is *derived* from the binary semigroup (X, F_2) or, simply, that F_n is derived from F_2 . We also say that (X, F_n) is a *totally (partially) ordered n -ary semigroup* for emphasizing that X is totally (partially) ordered.

It is easy to show (see Lemma 3.1 below) that if F_n is derived from F_2 and F_2 is either monotone or idempotent or has a neutral element, then so is F_n .

An n -ary semigroup (X, F_n) is called an *n -ary group* if for each $i \in \{1, \dots, n\}$, every $n-1$ elements $x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n$ in X and every $a \in X$, there exists a unique $b \in X$ with $F_n(x_1, \dots, x_{i-1}, b, x_{i+1}, \dots, x_n) = a$. It is easy to see from the definition that ordinary groups are exactly the 2-ary groups.

Clearly, a function F_n derived from a semigroup F_2 is n -associative but not every n -ary semigroup can be obtained in this way. Dudek and Mukhin [3] (see also Proposition 2.5) proved that an n -ary semigroup (X, F_n) is derived from a binary one if and only if (X, F_n) contains a neutral element or one can adjoin a neutral element to it. As a special case of this theorem they obtained that an n -ary group is derived from a group if and only if it contains a neutral element.

This result allows one to construct n -ary groups that are not derived from binary groups if n is odd. Indeed, let $(X, +)$ be a group and $n = 2k - 1$. Define $G_n(x_1, \dots, x_n) := \sum_{i=1}^n (-1)^i x_i$. It is easy to verify that G_n is n -associative

and we obtain an n -ary group. Moreover G_n is clearly monotone. It is also easy to check that there is no neutral element for G_n .

Finally, we say that an n -ary semigroup (X, F_n) is *quasitrivial* (or it is said to be *conservative*) if for every $x_1, \dots, x_n \in X$, we have $F_n(x_1, \dots, x_n) \in \{x_1, \dots, x_n\}$. Such an n -variable function F_n is called a *choice function*. One might also say that F_n preserves all subsets of X . Ackerman (see [1]) investigated quasitrivial semigroups and also gave a characterization of them.

Our paper is organized as follows. In Section 2 we collect the main results proved in the paper. In Section 3 we establish connections between n -ary semigroups and binary semigroups and prove Theorem 2.3. Section 4 is devoted to the proof of Theorems 2.4 and 2.6. Section 5 contains a few concluding remarks.

2 Main results

Let $I \subset \mathbb{R}$ be a not necessarily bounded, nonempty interval. We denote by \bar{I} the compact linear closure of I .¹ Let $g: \bar{I} \rightarrow \bar{I}$ be a decreasing function. For every $x \in I$, let $g(x-0)$ and $g(x+0)$ denote the limit of g at x from the left and from the right, respectively.² We denote by Γ_g the *completed graph* of g , which is a subset of \bar{I}^2 obtained by modifying the graph of the function g in the following way. If x is a discontinuity point of g , then we add a vertical line segment between the points $(x, g(x-0))$ and $(x, g(x+0))$ to extend the graph of g . Formally,

$$\Gamma_g = \{(x, y) \in \bar{I}^2 : g(x+0) \leq y \leq g(x-0)\}.$$

We call Γ_g (*id*)-*symmetric* if Γ_g is symmetric to the line $x = y$.

The following theorem gives a description of idempotent monotone (2-ary) semigroups with neutral elements. These semigroups were first investigated by Czogala and Drewniak [2], where the authors only dealt with closed subintervals of \mathbb{R} but the statement holds for any non-empty interval. On the other hand, instead of monotonicity it was assumed that the binary function is monotone increasing. However, Lemma 3.10 shows that monotonicity implies monotone increasingness in this case.

Theorem 2.1. *Let I be an arbitrary nonempty real interval. If a function $F_2: I^2 \rightarrow I$ is associative idempotent monotone and has a neutral element $e \in I$, then there exists a monotone decreasing function $g: \bar{I} \rightarrow \bar{I}$ with $g(e) = e$ such that*

$$F_2(x, y) = \begin{cases} \min(x, y) & \text{if } y < g(x), \\ \max(x, y) & \text{if } y > g(x), \\ \min(x, y) \text{ or } \max(x, y) & \text{if } y = g(x). \end{cases}$$

¹If I is bounded and we denote the end-points of I by m and M ($m \leq M$), then $\bar{I} = [m, M]$. If I is not bounded from below (or above), then we let $m = -\infty$ ($M = +\infty$, respectively). For instance, $\mathbb{R} = \mathbb{R} \cup \{\pm\infty\}$.

²Let m and M be the boundary points of \bar{I} . We use the convention that $g(m-0) = M$ and $g(M+0) = m$.

Now we present a complete characterization of idempotent monotone increasing (2-ary) semigroups with neutral elements. First this was proved by Martin, Mayor, and Torrens [7] for $I = [0, 1]$. Their theorem contained a small error in the description, but essentially it was correct. In the original paper [7] the following condition for g was given instead of the symmetry of Γ_g . The function $g: [0, 1] \rightarrow [0, 1]$ satisfies

$$\inf\{y : g(y) = g(x)\} \leq g^2(x) \leq \sup\{y : g(y) = g(x)\} \text{ for all } x \in [0, 1]. \quad (2)$$

Here (and below) $g^2(x)$ stands for $(g \circ g)(x)$.

The authors of [8] proved that Theorem 2.2 holds if F_2 is commutative also and shown that condition (2) is not equivalent to the symmetry of Γ_g . Recently, Theorem 2.2 was reproved in an alternative way in [5] for any subinterval of \mathbb{R} .

Theorem 2.2. *Let $I \subseteq \mathbb{R}$ be an arbitrary nonempty interval. A function $F_2: I^2 \rightarrow I$ is associative idempotent monotone and has a neutral element $e \in I$ if and only if there exists a decreasing function $g: \bar{I} \rightarrow \bar{I}$ with $g(e) = e \in I$ such that the completed graph Γ_g is symmetric and*

$$F_2(x, y) = \begin{cases} \min(x, y) & \text{if } y < g(x) \text{ or } y = g(x) \text{ and } x < g^2(x), \\ \max(x, y) & \text{if } y > g(x) \text{ or } y = g(x) \text{ and } x > g^2(x), \\ \min(x, y) \text{ or } \max(x, y) & \text{if } y = g(x) \text{ and } x = g^2(x). \end{cases} \quad (3)$$

Moreover, $F_2(x, y) = F_2(y, x)$ except perhaps the set of points $(x, y) \in I^2$ satisfying $y = g(x)$ and $x = g^2(x) = g(y)$.

If (X, F_n) is an n -semigroup having a neutral element e , then one can assign a semigroup (X, F_2) to it by letting $F_2(a, b) := F_n(a, e, \dots, e, b)$ for every $a, b \in X$. This map $F_n \mapsto F_2$ will be denoted by \mathcal{F} . Our main result in Section 3 is the following:

Theorem 2.3. *For any totally ordered set X , the map \mathcal{F} is a bijection between the set of associative idempotent monotone functions on X having neutral elements and the set of n -associative idempotent monotone functions on X having neutral elements.*

We will get the following result as an easy consequence of our investigation.

Theorem 2.4. *Let $I \subset \mathbb{R}$ be a nonempty interval, $n \geq 2$, and $F_n: I^n \rightarrow I$ an n -associative monotone increasing idempotent function with a neutral element. Then F_n is quasitrivial.*

Applying Theorems 2.3 and 2.2, we can obtain a practical method to calculate the value of $F_n(a_1, \dots, a_n)$ for any $a_1, \dots, a_n \in I$, where $I \subset \mathbb{R}$ is an interval.

For every decreasing function $g: \bar{I} \rightarrow \bar{I}$ a pair $(a, b) \in I^2$ is called *critical* if $g(a) = b$ and $g(b) = a$. By Theorem 2.2 and Lemma 3.10, for every idempotent monotone semigroup (X, F_2) with a neutral element, there exists a unique decreasing function g satisfying (3). Theorem 2.2 shows also that F_2 commutes on

every non-critical pair $(x, y) \in I^2$ (i.e., $F_2(x, y) = F_2(y, x)$). Since for a critical pair (a, b) the value of $F_2(a, b)$ and $F_2(b, a)$ can be independently chosen from g , we have two cases. We might have that F_2 commutes on a, b or not. A pair (a, b) is called *extra-critical* if $F_2(a, b) \neq F_2(b, a)$. We note that being critical or extra-critical are both symmetric relations.

Finally, in order to simplify notation and give a compact way to express the value of F_n at some n -tuple (a_1, \dots, a_n) of the elements from a totally ordered set, we introduce the following. The smallest and the largest elements of the set $\{a_1, \dots, a_n\}$ are denoted by c and d , respectively. There exist i, j with $1 \leq i \leq j \leq n$ such that $a_i = c$ or d , $a_j = c$ or d and $a_k \neq c$ and d for every $k < i$ and $k > j$. We write $e_1 := a_i$ and $e_2 := a_j$.

The following statement was proved in [3]:

Proposition 2.5 (Dudek, Mukhin). *If (X, F_n) is an n -ary semigroup with a neutral element e , then F_n is derived from a binary function F_2 , where*

$$F_2(a, b) := F_n(a, e, \dots, e, b). \quad (4)$$

Theorem 2.6. *Let $F_n: I^n \rightarrow I$ be an n -associative idempotent function with a neutral element that is monotone in its first and last coordinates. If (c, d) is a not an extra-critical pair, then $F_n(a_1, \dots, a_n) = F_2(c, d)$. If (c, d) is an extra-critical pair, then $F_n(a_1, \dots, a_n) = F_2(e_1, e_2)$.*

Now we point out three important consequences of Theorem 2.6. First we generalize Czogala–Drewniak’s theorem (Theorem 2.1) as follows.

Theorem 2.7. *Let I be an arbitrary nonempty real interval. If a function $F_n: I^n \rightarrow I$ is n -associative idempotent monotone and has a neutral element $e \in I$, then there exists a monotone decreasing function $g: \bar{I} \rightarrow \bar{I}$ with $g(e) = e$ such that Γ_g is symmetric and*

$$F_n(a_1, \dots, a_n) = \begin{cases} c & \text{if } c < g(d), \\ d & \text{if } c > g(d), \\ c \text{ or } d & \text{if } c = g(d), \end{cases}$$

where c and d denote the minimum and the maximum of the set $\{a_1, \dots, a_n\}$, respectively.

We note that a generalization of Theorem 2.2 is essentially stated in Theorem 2.6. In [8] the authors investigated idempotent uninorms, which are associative, commutative, monotone functions with a neutral element and idempotent also. We introduce *n -uninorms*, which are n -associative, commutative, monotone functions with neutral element. Here we show a generalization of [8, Theorem 3] for n -ary operations.

Theorem 2.8. *An n -ary operator U_n is an idempotent n -uninorm on $[0, 1]$ with a neutral element $e \in [0, 1]$ if and only if there exists a decreasing function*

$g: [0, 1] \rightarrow [0, 1]$ with $g(e) = e$ and with symmetric graph Γ_g such that

$$U_n(a_1, \dots, a_n) = \begin{cases} c & \text{if } c < g(d) \text{ or } d < g(c), \\ d & \text{if } c > g(d) \text{ or } d > g(c), \\ c \text{ or } d & \text{if } c = g(d) \text{ and } d = g(c), \end{cases} \quad (5)$$

where c and d are as in Theorem 2.7. Moreover, if (c, d) is a critical pair ($c = g(d), d = g(c)$), then the value of $U_n(a_1, \dots, a_n)$ can be chosen to be c or d arbitrarily and independently from other critical pairs.

We may generalize our concept in the following way. Let $X^* = \bigcup_{n \in \mathbb{N}} X^n$ be the set of finite length words over the alphabet X . A multivariate function $F: X^* \rightarrow X$ is *associative* if it satisfies

$$F(\mathbf{x}, \mathbf{x}') = F(F(\mathbf{x}), F(\mathbf{x}'))$$

for all $\mathbf{x}, \mathbf{x}' \in X^*$. It is easy to check that $F|_{X^n}$ is n -associative for every $n \in \mathbb{N}$. We say that F is idempotent or monotone or that it has a neutral element if so are the functions $F|_{X^n}$ for every $n \in \mathbb{N}$.

Theorem 2.9. *Let I be a nonempty real interval. Then $F: I^* \rightarrow I$ is associative idempotent monotone and has a neutral element if and only if there is a decreasing function $g: \bar{I} \rightarrow \bar{I}$ with symmetric completed graph Γ_g such that $F|_{X^2}$ satisfies (3). Furthermore F must be monotone increasing in each variable.*

Concerning to associativity of multivariate functions the interested reader is referred to [4, 6].

3 From n -ary to binary semigroups

In this section we prove Theorem 2.3. Therefore the main purpose of this section is to transfer properties from an n -ary semigroup to the corresponding binary semigroup. We start with the converse. We have already mentioned that, given a semigroup (X, F_2) , one can easily construct the n -ary semigroup (X, F_n) , where $F_n = \underbrace{F_2 \circ \dots \circ F_2}_{n-1}$. The following lemma is an easy consequence of the definitions.

Lemma 3.1. *Let (X, F_2) be a partially ordered semigroup. If F_2 has any of the following properties*

- (i) *monotone*
- (ii) *idempotent*
- (iii) *has a neutral element*

then so does the function F_n .

Observation 3.2. If F_2 is defined by (4), the element e is also a neutral element of F_2 since $F_2(e, a) = F_n(e, \dots, e, a) = a = F_n(a, e, \dots, e) = F_2(a, e)$ for every $a \in X$.

Lemma 3.3. Let $F_n: X^n \rightarrow X$ be an n -associative function on the partially ordered set X . Assume F_n is idempotent and monotone in the first and the last coordinates and is derived from an associative function F_2 . Then F_2 is monotone.

Proof. We show that if F_n is monotone in its last coordinate then so is F_2 . Take an arbitrary $a \in X$ and let $b = F_{n-1}(a, \dots, a)$. In this case $F_2(b, a) = F_n(a, \dots, a) = a$. Substituting $a = F_2(b, a)$, we obtain $a = F_2(b, F_2(b, a))$. Using the same substitution $n - 2$ times, we get that a can be expressed as $F_{n-1}(c_1, \dots, c_{n-1})$ for some c_1, \dots, c_{n-1} . Then $F_2(a, x) = F_n(c_1, \dots, c_{n-1}, x)$ is clearly monotone in its last coordinate.

Similarly, F_2 is monotone in its first coordinate if F_n is. \square

Remark 3.4. If F_n is n -associative idempotent and monotone in the first and the last variables on a poset X , then, by Lemma 3.3, F_2 is also monotone. It is easy to show that $F_k := \underbrace{F_2 \circ \dots \circ F_2}_{k-1}$ is k -associative and monotone in each

variable. In particular, F_n is monotone in each of its variables.

Lemma 3.5. Let $F_n: X^n \rightarrow X$ be an n -associative function on a **totally ordered** set. Assume F_n is idempotent and monotone in each variable and F_n has a neutral element or is derived from an associative function F_2 . Then F_2 is idempotent as well.

Proof. We prove that $F_k = F_2 \circ \dots \circ F_2$ is idempotent for every $2 \leq k \leq n$. We use backward induction. Arguing by contradiction, assume that for some k with $3 \leq k \leq n$ there exists $a \in X$ such that has $F_{k-1}(a, \dots, a) = b \neq a$ and by the inductive hypothesis $F_k(x, \dots, x) = x$ for every $x \in X$. We note that the second condition holds for $k = n$, since F_n is idempotent. Clearly, we may assume without any loss that $a < b$. We compare the following terms:

Table 1:

$F_k(a, \dots, a, b)$	$F_k(a, \dots, a, b, b)$	$F_k(a, \dots, a, b, b, b)$	\dots	$F_k(a, b, \dots, b, b)$
$F_k(a, \dots, a, a)$	$F_k(a, \dots, a, b, a)$	$F_k(a, \dots, a, b, b, a)$	\dots	$F_k(a, b, \dots, b, a)$

The function F_k is monotone in each variable by Remark 3.4. Observe that in Table 1 the elements in each column only differ in the last coordinate. Hence each of the elements in the lower row is not greater than the element above it by the monotonicity of F_k .

Now we calculate expressions in Table 1. It is clear that $F_k(a, \dots, a) = a$ by the inductive assumption. Before we continue, we present two useful lemmas.

Lemma 3.6. *Let a and b be as above. Further, let $x_1 = \dots = x_l = a$ and $x_{l+1} = \dots = x_k = b$. Then for every $\pi \in \text{Sym}(k)$ we have*

$$F_k(x_1, \dots, x_k) = F_k(x_{\pi(1)}, \dots, x_{\pi(k)}).$$

Proof. Substituting $b = F_{k-1}(a, \dots, a)$ in the expression above, it is easy to see that we may rearrange a 's and b 's arbitrarily. \square

Lemma 3.7. *Let l and m be fixed and $l + m = k$. If $1 \leq m \leq k - 2$, then*

$$F_k(\underbrace{a, \dots, a}_l, \underbrace{b, \dots, b}_m) = F_l(\underbrace{a, \dots, a}_l).$$

In particular, if $m = k - 1$, then

$$F_k(a, \underbrace{b, \dots, b}_{k-1}) = a.$$

Proof. A direct calculation shows that the statement holds. Indeed,

$$F_k(\underbrace{a, \dots, a}_l, \underbrace{b, \dots, b}_m) = F_k(\underbrace{a, \dots, a}_l, \underbrace{F_{k-1}(a, \dots, a), \dots, F_{k-1}(a, \dots, a)}_m).$$

Now using associativity of F_2 and idempotency of F_k , we obtain

$$F_2(a, F_{k-1}(a, \dots, a)) = F_k(a, \dots, a) = a.$$

Applying Lemma 3.6 and the previous observation m times, we obtain

$$F_k(\underbrace{a, \dots, a}_l, \underbrace{F_{k-1}(a, \dots, a), \dots, F_{k-1}(a, \dots, a)}_m) = F_l(a, \dots, a).$$

\square

Using Lemma 3.6 and Lemma 3.7, we get that

$$\begin{aligned} F_k(a, \dots, a) &= a \\ F_k(a, \dots, a, a, b, a) &= F_k(a, \dots, a, a, a, b) = F_{k-1}(a, \dots, a), \\ F_k(a, \dots, a, b, b, a) &= F_k(a, \dots, a, a, b, b) = F_{k-2}(a, \dots, a), \\ &\vdots \\ F_k(a, b, b, \dots, b, a) &= F_k(a, a, b, \dots, b, b) = F_2(a, a), \\ F_k(a, b, \dots, b, b) &= a. \end{aligned} \tag{6}$$

Since in each column of the table we just change the last coordinate, we can use monotonicity. We note that F_k is increasing (order-preserving) in the last

variable since $F_k(a \dots, a, b) = b > a = F_k(a, \dots, a, a)$. Substituting the results of (6) into the table, we get the following:

$$\begin{array}{c|c|c|c|c} F_{k-1}(a, \dots, a) & F_{k-2}(a, \dots, a) & F_{k-3}(a, \dots, a) & \dots & a \\ \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \\ a & F_{k-1}(a, \dots, a) & F_{k-2}(a, \dots, a) & \dots & F_2(a, a) \end{array}$$

Here the notation \uparrow means that an element in the lower row is less than or equal to the corresponding element in the upper row. Thus,

$$a \leq F_{k-1}(a, \dots, a) \leq F_{k-2}(a, \dots, a) \leq \dots \leq F_2(a, a) \leq a.$$

This gives

$$a = F_{k-1}(a, \dots, a) = F_{k-2}(a, \dots, a) = \dots = F_2(a, a),$$

a contradiction, since $F_{k-1}(a, \dots, a) = b \neq a$ by our assumption. This shows that F_k is idempotent for every $k \geq 2$, finishing the proof of Lemma 3.5. \square

The underlying set of the n -associative function in Lemma 3.5 is totally ordered. The following example shows that this requirement is essential.

Example 3.8. For $k \geq 3$ we construct a k -ary semigroup (X, F_k) , which is derived from a non-idempotent semigroup (X, F_2) , where F_2 is monotone in both of its variables and has a neutral element. Thus, F_{k-1} and F_k are also monotone having neutral element. We show that F_k is idempotent and F_{k-1} is not idempotent, thus F_2 cannot be idempotent by Lemma 3.1 (ii). This example shows that the condition that X is a totally ordered set is crucial in Lemma 3.5.

Let $X = \{m, M\} \cup Z_{k-1}$, where Z_{k-1} is the cyclic group of order $k-1$. We define a partial ordering on X in the following way. M and m are the largest and smallest elements of X , respectively. The elements of Z_{k-1} are mutually incomparable but they are all larger than m and smaller than M . The set X endowed with this partial ordering is a modular lattice. Further we build an associative function F_2 as follows:

$$F_2(x, y) = \begin{cases} M & \text{if } x = M \text{ or } y = M, \\ m & \text{if } x = m \text{ or } y = m \text{ and } x, y < M, \\ xy & \text{if } x, y \in Z_{k-1}. \end{cases}$$

It is easy to verify that F_2 is associative and monotone increasing in both of its variables. The identity element e of Z_{k-1} is the neutral element of (X, F_2) . One can define F_{k-1} and F_k as before. By Lemma 3.1 the functions F_{k-1} and F_k are $(k-1)$ - and k -associative functions, respectively. Both of them are monotone having neutral element. Finally, it is easy to check that F_{k-1} is not idempotent since $F_{k-1}(a, \dots, a) = e$ for every $a \in Z_{k-1}$ while $F_k(x, \dots, x) = x$ for every $x \in X$. Note that the cyclic group Z_{k-1} might have been substituted by any nontrivial group whose exponent divides $k-1$.

Remark 3.9. For distributive lattices the statement of Lemma 3.5 seems true, but a potential proof would be basically different from the proof of the lemma. Thus, it goes beyond the topic of the current paper. (See also Question 5.2 in Section 5.)

The following lemma provides extra information about monotone, associative and idempotent semigroups.

Lemma 3.10. *Let X be a partially ordered set. If $F_2: X^2 \rightarrow X$ is associative idempotent and monotone in each variable, then F_2 is monotone increasing in each variable.*

Proof. Assume that F_2 is not monotone increasing in each variable. Let us assume that F_2 is decreasing in the second variable. We also exclude the case when F_2 is both increasing and decreasing in the second variable (i.e., $F_2(x, \cdot)$ is constant for any fixed $x \in X$), so that we may assume that there exist $x, y, z \in X$ such that $y < z$ and $F_2(x, y) > F_2(x, z)$.

Now by the idempotency of F_2 we have $F_2(F_2(x, x), y) = F_2(x, y)$ and $F_2(F_2(x, x), z) = F_2(x, z)$. Our assumption then gives

$$F_2(F_2(x, x), y) > F_2(F_2(x, x), z).$$

Using the associativity of F_2 we get $F_2(x, F_2(x, y)) > F_2(x, F_2(x, z))$.

On the other hand, since $F_2(x, y) > F_2(x, z)$ and F_2 is decreasing in the second variable we get $F_2(x, F_2(x, y)) \leq F_2(x, F_2(x, z))$, which contradicts our assumption.

One can get the same type of contradiction if we switch the role of the coordinates. Thus, F_2 is monotone increasing in both variables. \square

Remark 3.11. The following examples demonstrate that if we omit any of the conditions of Lemma 3.10, the conclusion of the lemma fails.

1. Let $F_2(x, x) = x$ for $x \in \mathbb{R}$ and $F_2(x, y) = 0$ if $x, y \in \mathbb{R}$, $x \neq y$. Then F_2 is associative and idempotent, but not monotone in each variable.
2. Let $F_2(x, y) = 2x - y$ for $x, y \in \mathbb{R}$. Then F_2 is idempotent and monotone in each variable, but not associative and clearly not monotone increasing.
3. Let $F_2(x, y) = -x$, if $x, y > 0$, and $F_2(x, y) = 0$ otherwise. Then F_2 is associative, since $F_2(x, F_2(y, z)) = F_2(F_2(x, y), z) = 0$ and F_2 is monotone decreasing in each variable and F_2 is not idempotent.

Corollary 3.12. *If (X, F_n) is a totally ordered n -ary semigroup, where $F_n = \underbrace{F_2 \circ F_2 \circ \dots \circ F_2}_{n-1}$ is idempotent and monotone in the first and the last variables, then F_n is monotone increasing in each variable. Moreover, $F_k = \underbrace{F_2 \circ \dots \circ F_2}_{k-1}$ is monotone increasing for every $k \geq 2$.*

Proof. By definition, F_2 is associative. Since F_n is monotone in each variable, so is F_2 by Lemma 3.3. By Lemma 3.5, F_2 is idempotent. Thus by Lemma 3.10, it is monotone increasing. Thus $F_k = F_2 \circ F_2 \circ \cdots \circ F_2$ is also monotone increasing for every $k \geq 2$. \square

If F_n is n -associative and has a neutral element, then there exists F_2 such that $F_n = F_2 \circ F_2 \circ \cdots \circ F_2$. Using the results of this section, we prove the following proposition.

Proposition 3.13. *Let (X, F_n) be a totally ordered n -ary semigroup, which is monotone idempotent and has a neutral element. Then F_n is derived from a binary semigroup (X, F_2) , where F_2 is also monotone idempotent and it also has a neutral element. Moreover F_n is monotone increasing in each variables.*

Proof. Since F_n is idempotent n -associative and has a neutral element, it follows from Proposition 2.5 that $F_n = F_2 \circ \cdots \circ F_2$, where $F_2: X^2 \rightarrow X$ is associative. By Lemmas 3.3, 3.5, and 3.10, F_2 is monotone increasing and idempotent. By Observation 3.2 that in this case F_2 has a neutral element, as well. \square

Proof of Theorem 2.3. By Proposition 3.13, every n -associative function F_n which is monotone idempotent and has a neutral element, is derived from an associative function F_2 determined by $F_2(a, b) := F_n(a, e, \dots, e, b)$ which is monotone idempotent and has a neutral element. Then we have

$$F_n(a, \dots, a, b) = F_n(a, e, \dots, e, b) = F_n(a, b, \dots, b) = F_2(a, b). \quad (7)$$

Recall that the map that assigns F_2 to F_n was denoted by \mathcal{F} . By Lemma 3.1 and Proposition 3.13, for every F_2 , there exists F_n satisfying (7) whence \mathcal{F} is surjective. The map \mathcal{F} is injective since F_2 uniquely defines F_n . This finishes the proof of Theorem 2.3. \square

Remark 3.14. Using Corollary 3.12 we may weaken the assumptions of Theorem 2.3, where F_n is assumed to be monotone in each variable. Instead, we might have assumed that F_n is monotone in the first and last variables.

Lemma 3.15. *Let (X, F_n) be a totally ordered n -ary semigroup derived from (X, F_2) , where F_2 is idempotent, associative, monotone increasing and has a neutral element. Then*

$$F_n(a, y_1, \dots, y_{n-2}, b) = F_2(a, b) \quad (8)$$

whenever $a \leq y_1, \dots, y_{n-2} \leq b$.

Proof. By Theorem 2.3, the function F_n is monotone, and therefore, the claim directly comes from (7). \square

4 Proof of the main results

Proof of Theorem 2.4: It follows from Proposition 3.13 that F_n is derived from an associative function F_2 . Moreover F_2 is monotone, idempotent and has a neutral element. Therefore, we may apply Theorem 2.2 in a special form which we obtain when F_2 is a choice function (i.e., when (X, F_2) is quasitrivial). Since F_n is obtained as the composition of $n - 1$ copies of F_2 , we get that F_n is also a choice function. \square

Proof of Theorem 2.6. First assume that c and d commute with every element of the set $A := \{a_1, \dots, a_n\} \subset I$. In this case we may assume, using idempotency of F_2 , that there exists $k \leq n$ such that $F_n(a_1, \dots, a_n) = F_k(c, a'_2, \dots, a'_{k-1}, d)$ and $c < a'_i < d$ for all $i = 2, \dots, k-1$. By Proposition 3.13 we can apply Lemma 3.15 that gives $F_n(a_1, \dots, a_n) = F_2(c, d)$.

Now assume that d does not commute with an element of A but c commutes with all of them. In this case $g(d) \in A$ is the one not commuting with d . Since c is the smallest element of A we get $c < g(d)$. Further, d is the largest element of A and g is decreasing so $g(a_i) > c$ for all $i = 1, \dots, n$. Theorem 2.2 gives $F_2(c, a_i) = F_2(a_i, c) = c$ for all $i = 1, \dots, n$. Therefore $F_n(a_1, \dots, a_n) = c$. Since $F_2(c, a_i) = c$ for every i , we get $F_n(a_1, \dots, a_n) = F_2(c, d) = c$. A similar argument shows that $F_n(a_1, \dots, a_n) = d = F_2(c, d)$ if c and d switch the roles.

Finally, assume that neither c nor d commutes with every element of A . In this case the set A contains $g(c)$ and $g(d)$ and $g(g(c)) = c$, $g(g(d)) = d$. We claim that $g(c) = d$ and $g(d) = c$. Indeed, if $g(c) \in A$, then $g(c) \leq d$ since d is the largest element of A , and similarly $g(d) \geq c$. Since g is monotone and $g(g(d)) = d$, we get $d = g(g(d)) \leq g(c)$. Therefore $g(c) = d$. Similarly using $c = g(g(c)) \geq g(d)$ we get $g(d) = c$. What we obtained is that (c, d) is an extra-critical pair in this case.

Now c and d are the elements in A that do not commute. This also implies that c and d commute with all other element of A .

By definition of e_1 and e_2 (see Section 2), e_1 and e_2 are the value of the first and respectively the last appearance of c or d . Since e_1 commutes with its left neighbours and e_2 commutes with its right neighbours, we may assume that $a_1 = e_1$ and $a_2 = e_2$. We get the following cases:

- (i) If $e_1 \neq e_2$, then by Lemma 3.15 $F_n(e_1, \dots, e_2) = F_2(e_1, e_2)$.
- (ii) If $e_1 = e_2$, then we show that $F_n(e_1, \dots, e_2) = F_2(e_1, e_2) = e_1$.

Using Lemma 3.15 for arbitrary number of variables, we get that every subsequence of a_1, \dots, a_n consisting of elements lying strictly between c and d can be eliminated. Thus, one can write $F_n(a_1, \dots, a_n) = F_k(b_1, \dots, b_k)$, where $k \leq n$ and $b_i = c$ or d for every $1 \leq i \leq k$ and, in our case, $b_1 = b_k = e_1$. Since F_2 is idempotent, we may assume $b_i \neq b_{i+1}$ for $1 \leq i \leq k-1$. Using idempotency and associativity of F_2 again, we have $F_2(F_2(c, d), F_2(c, d)) = F_2(c, d)$ and $F_2(F_2(d, c), F_2(d, c)) = F_2(d, c)$. Therefore $F_k(b_1, \dots, b_k)$ can be reduced to either $F_3(c, d, c)$ or $F_3(d, c, d)$.

If $F_2(c, d) = c$, then $F_3(c, d, c) = F_2(F_2(c, d), c) = F_2(c, c) = c = e_1$.

If $F_2(c, d) = d$, then $F_3(c, d, c) = F_2(F_2(c, d), c) = F_2(d, c)$. Since c and d do not commute we get $F_2(d, c) = c = e_1$.

Similarly, one can verify that $F_3(d, c, d) = d = e_1$. This finishes the proof of Theorem 2.6. \square

5 Concluding remarks

In this paper we have investigated the n -ary associative, idempotent, monotone functions $F_n: X^n \rightarrow X$ that have neutral elements. We have shown that such an F_n in general setting when the underlying set X is totally ordered implies the existence of binary functions $F_2: X^2 \rightarrow X$ with similar properties such that F_n is derived from F_2 . However many of the properties of F_n are inherited by F_2 if X is only a partially ordered set. We summarize the results of Section 3 (if X is a totally ordered set) in the following table.

Properties of F_n		Properties of F_2
n -assoc. with a neutral element	\implies	assoc. with a neutral element
Now we assume $F_n = F_2 \circ \dots \circ F_2$:		
n -assoc., idempotent, monotone	\implies	monotone
n -assoc., idempotent, monotone	\implies	idempotent
n -assoc., idempotent, monotone	\implies	monotone increasing
Some easy observations show:		
n -associative	\Leftarrow	associative
monotone increasing	\Leftarrow	monotone increasing
idempotent	\Leftarrow	idempotent
has a neutral element	\Leftarrow	has a neutral element
Thus:		
n -assoc., idempotent, mon. incr. with a neutral element	\iff	assoc., idemp., mon. incr. with a neutral element

In the main results we have obtained a characterization of n -associative, idempotent, monotone functions on any (not necessarily bounded) subinterval of \mathbb{R} in the spirit of the characterization of the binary case. We also generalize the classical Czogala–Drewniak theorem. In addition, we get that every n -associative, idempotent and monotone function with a neutral element must be quasitrivial (conservative).

Further improvement would be based on the elimination of any of the properties of F_n . The most crucial property seems to be that F_n has a neutral element since all of our results based on this condition as otherwise F_n is not necessarily derived from F_2 . On the other hand, in [7] one can be found a characterization of associative, conservative, monotone increasing, idempotent binary functions defined on $[0, 1]$ without assumption of having a neutral element. Therefore, we suggest the following questions for further investigation.

Question 5.1. *How can we characterize n -associative, monotone, idempotent functions on a subinterval of \mathbb{R} ?*

Question 5.2. *How can we characterize n -associative functions on distributive lattices provided that the functions are monotone idempotent and have neutral elements?*

References

- [1] N. L. Ackerman, A characterization of quasitrivial n -semigroups, preprint.
- [2] E. Czogala, J. Drewniak, Associative monotonic operations in fuzzy set theory, *Fuzzy Sets and Systems* **12** (1984) 249–269.
- [3] W. A. Dudek, V. V. Mukhin, On n -ary semigroups with adjoint neutral element, *Quasigroups and Related Systems* **14** (2006) 163–168.
- [4] M. Grabisch, J.-L. Marichal, R. Mesiar, E. Pap. *Aggregation functions*. Encyclopedia of Mathematics and its Applications, vol. 127. Cambridge University Press, Cambridge, 2009
- [5] J. Devillet, G. Kiss, J.-L. Marichal, B. Teheux, Associativity and bisymmetry on reflexive, monotone functions, *in preparation*.
- [6] E. Lehtonen, J.-L. Marichal, B. Teheux, Associative string functions, *Asian-European Journal of Mathematics* **7**(4) (2014), 1450059.
- [7] J. Martin, G. Mayor, J. Torrens, On locally internal monotonic operations, *Fuzzy Sets and Systems* **137** (2003) 27–42.
- [8] D. Ruiz-Aguilera, J. Torrens, B. De Baets, J. Fodor, Some remarks on the characterization of idempotent uninorms, in: E. Hüllermeier, R. Kruse, F. Hoffmann (eds.), *Computational Intelligence for Knowledge-Based Systems Design*, Lecture Notes in Artificial Intelligence, vol. 6178, Springer, 2010, 425–434.