

Nonsuppurative (Aseptic) Meningoencephalomyelitis Associated with Neurovirulent Astrovirus Infections in Humans and Animals

Gábor Reuter,^{a,b} Péter Pankovics,^{a,b} Ákos Boros^{a,b}

^aRegional Laboratory of Virology, National Reference Laboratory of Gastroenteric Viruses, ÁNTSZ Regional Institute of State Public Health Service, Pécs, Hungary

^bDepartment of Medical Microbiology and Immunology, Medical School, University of Pécs, Pécs, Hungary

SUMMARY Astroviruses are thought to be enteric pathogens. Since 2010, a certain group of astroviruses has increasingly been recognized, using up-to-date random amplification and high-throughput next-generation sequencing (NGS) methods, as potential neurovirulent (Ni) pathogens of severe central nervous system (CNS) infections, causing encephalitis, meningoencephalitis, and meningoencephalomyelitis. To date, neurovirulent astrovirus cases or epidemics have been reported for humans and domesticated mammals, including mink, bovines, ovines, and swine. This comprehensive review summarizes the virology, epidemiology, pathology, diagnosis, therapy, and future perspective related to neurovirulent astroviruses in humans and mammals, based on a total of 30 relevant articles available in PubMed (searched by use of the terms "astrovirus/encephalitis" and "astrovirus/meningitis" on 2 March 2018). A paradigm shift should be considered based on the increasing knowledge of the causality-effect association between neurotropic astroviruses and CNS infection, and attention should be drawn to the role of astroviruses in unknown CNS diseases.

KEYWORDS animal, astrovirus, encephalitis, human, meningitis, meningoencephalomyelitis, neurotropic, neurovirulent

INTRODUCTION

Encephalitis is an inflammation of the brain tissue associated with clinical evidence of nervous system dysfunction. Encephalitis and central nervous system (CNS) infections are caused by a wide spectrum of known and unknown infectious agents, including bacteria, viruses, fungi, and parasites, in humans and animals (1, 2). Approximately 25% of the notifiable infectious diseases listed by the World Organization for Animal Health (OIE) are caused by neurotropic microbes (3). The microbial diagnosis of acute encephalitis and CNS diseases is challenging, especially for immunocompromised individuals, because of the atypical clinical signs and the potential presence of novel or uncommon infectious agents (4). Despite the application of various classical and modern diagnostic methods (e.g., culture, immunohistochemical, serologic, or molecular methods), a causative agent is not identified for a substantial proportion of encephalitis cases (5, 6). Failure to identify the etiology of neuroinfections impairs or delays appropriate therapeutics and influences the morbidity and mortality of these diseases (7).

Nonsuppurative (aseptic) encephalitis is an inflammation of the brain observed in postmortem histopathological examinations and is indicative of an infectious viral etiology. Perivascular cuffs with mononuclear cells, gliosis, and neuronal necrosis are common characteristic features of the disease (8). New unbiased molecular technologies, such as random amplification and high-throughput next-generation sequencing (NGS) methods, are valuable tools for differential diagnosis of unexplained diseases. An excellent example is that this technique raised the possibility that astroviruses may be associated with encephalitis in humans (9-11). Growing evidence of neurotropic/ neurovirulent astrovirus (Ni-AstV) strains has arisen for cases of encephalitis, meningitis, and meningoencephalomyelitis in humans and animals, including mink (12), bovines (13), ovines (14), and swine (15). If astrovirus infection reaches the CNS, the virus may infect multiple types of neurons (e.g., Purkinje cells, interneurons, and CA pyramidal neurons) and glial cells (e.g., astrocytes) in different parts of the CNS, including the brainstem, cerebellar/cerebral cortex, hippocampus, and spinal cord. Astrovirus infection may also cause neuronal degeneration, necrosis, neuronophagia, and gliosis (15-18). This means that the term "neuroinvasive astrovirus" may also be appropriate.

These unexpected initial results of astrovirus neuroinfection are starting to change our view about the astrovirus disease spectrum and pathogenesis, although there are several unanswered questions. At the same time, the evidence of astrovirus CNS infections opens an important and interesting new research field in the pathology of astroviruses.

NEUROVIRULENT ASTROVIRUS INFECTIONS IN HUMANS

Astroviruses are nonenveloped, single-stranded, positive-sense RNA viruses that are typically transmitted by the fecal-oral route and may cause gastrointestinal disease (19). One of the two astrovirus genera, Mamastrovirus, comprises viruses infecting mammals, including 4 species with human astroviruses. The first species, Mamastrovirus 1 (MAstV1), corresponds to "classical" (fecal origin) human astrovirus types 1 to 8 (HAstVs 1 to 8). Since 2008, a further three species (MAstV 6,8, and 9) of human astrovirus have been identified. MAstV6 includes human astrovirus clades MLB1 ("MLB" stands for Melbourne, the place of the first detection), MLB2, and MLB3 (20-22). The MAstV8 species includes astrovirus clades VA2 ("VA" stands for Virginia; also known as HMO-A [human-mink-ovine-like astroviruses]) (21) and VA4 (22). Finally, the MAstV9 species includes astrovirus clades VA1 (or HMO-C) (23) and VA3 (or HMO-B) (19, 21, 24). One further human astrovirus, VA5 (25), is presently unassigned (19). Although these novel astroviruses were initially detected in children with gastroenteritis, no definitive association has yet been established (19, 26, 27).

Both classical and, especially, novel astroviruses have been identified in unexpected CNS infections, indicating that some of these viruses are able to enter the host through the mucosal surfaces and infect extraintestinal cells and organs. In 2010, astrovirus-associated encephalitis was reported for the first time, in a 15-year-old immunocompromised boy with X-linked agammaglobulinemia in the United States (9). The patient had suicidal and homicidal ideation, headache, memory loss, and ataxia. He also had progressive cognitive decline and movement and communication problems, became comatose, and died 71 days after admission (9). The possible connection to a mink farm and the phylogenetic relationship of HAstV-PS to mink astroviruses led the authors to assume that the mink was a potential source of exposure (9). However, we cannot exclude the possibility of human-to-human transmission of the infection in this case. To date, a further eight human cases of astrovirus encephalitis have been reported, related to genotype VA1/HMO-C (n = 5) (9, 28-31), the MLB group (MLB1 and MLB2) (32, 33), and the "classical" HAstV4 species (34), although in the latter case, due to the undetermined complete capsid coding region (ORF2), the exact classification of the virus is not possible. The nine cases of neurotropic human astrovirus infections are summarized in Table 1. Except for an adult immuno-competent woman with MLB2 self-limited CNS infection, all other infections occurred in immunocompromised patients with severe underlying diseases. Six of the infections were fatal (19).

These results led to the conclusion that astroviruses may be a cause of previously unrecognized and severe neuroinfections, at least in immunocompromised humans, with a potential tropism for the CNS as a novel human disease (19, 35).

NEUROVIRULENT ASTROVIRUS INFECTIONS IN ANIMALS

Astrovirus-associated encephalitis is not restricted to humans. Based on pioneer studies, neurovirulent astroviruses have been reported

from four domesticated mammals: mink (*Mustela vison*)(12), bovines (*Bos taurus*)(13), ovines (*Ovis aries*)(14), and swine (*Sus scrofa domestica*)(15).

Mink

The first report of neurovirulent astrovirus infections ("shaking mink syndrome") from animals was reported retrospectively in 2010 for farmed minks in Denmark by the NGS method (12). Shaking, staggering gait, ataxia, and paraplegia were the characteristic clinical signs (Table 2). The symptoms were reproduced by the inoculation of brain homogenate from mink displaying "shaking mink syndrome" into experimentally infected healthy mink kits (12). A similar neurological disease was observed in farmed minks in Denmark in 2000 and 2002 and in Sweden and Finland in 2001 (12, 16), which suggests the epidemic spread of this disease.

Bovines

Many more data and cases are available about neurotropic astroviruses in bovines. While there are some historical reports since more than 60 years ago about "European sporadic bovine encephalitis/encephalomyelitis" with unknown origin in bovines (3642), the first confirmed cases of bovine neurovirulent astrovirus infections were reported only in 2013, from the United States (13). The neurovirulent astrovirus (bovine astrovirus [BoAstV] strain NeuroS1; accession number [KF233944](#)) was identified by NGS as the solely detectable virus in the CNS, and its presence was subsequently confirmed retrospectively for 3 of 32 other bovines that had bovine encephalitides (meningoencephalomyelitis and ganglioneuritis) of unknown etiology by reverse transcription-PCR (RT-PCR) and in situ RNA hybridization (ISH) (13). Viral RNA was detected in the spinal cord, brainstem, and cerebellum (13)(Table 2). The initial animal had lateral recumbency with opisthotonus and limb rigidity (13). Independently, shortly thereafter, similar series of cases identified by the NGS method were reported from Switzerland (17). In 2014, 5 (22.7%) of 22 cattle with nonsuppurative encephalitis of unknown etiology were associated with neurotropic astrovirus infections (17). The presence of astroviral RNA (BoAstV-CH13; accession number [KM035759](#)) at the site of the affected neurons was also confirmed by ISH in all 5 cases (17)(Table 2). This astrovirus has been present in the Swiss bovine population since at least 1995, particularly within the age group of

TABLE 1 Neurovirulent astrovirus infection cases associated with CNS disease in humans, in order of publication year (2010 to 2017)*

Patient (country)	Underlying disease and conditions	Clinical diagnosis	Symptoms	Outcome	Astrovirus type (method used for pathogen identification), strain name, available		
					GenBank accession no.	Potential source	Reference
15-yr-old boy (USA)	X-linked agammaglobulinemia (immunocompromised)	Encephalitis	Suicidal and homicidal ideation, headache, memory loss, ataxia, progressive cognitive decline	Fatal (within 71 days)	VA1/HMO-C (NGS), HuAstV-PS (Puget Sound) or HuAstV-SG, CQ891990	Exposure to mink?	9
3-mo-old boy (Switzerland)	Severe combined immunodeficiency, hematopoietic stem cell transplantation, chemotherapy (immunocompromised)	Meningoencephalitis	Multiorgan dysfunction (hepatic and respiratory)	Fatal (within 17 days posttransplantation)	HuAstV4 (NGS), HQ396880-HQ396890	Nosocomial	34
18-mo-old boy (UK)	Cartilage hair hypoplasia with hereditary immunodeficiency, hematopoietic stem cell transplantation, chemotherapy (immunocompromised)	Encephalopathy, encephalitis	Irritability, dystonia, reduced consciousness	Fatal (within 9 mo)	VA1/HMO-C (NGS), VA1/HMO-C UK1 (a) London, KJ920196	Community acquired?	28
42-yr-old man (UK)	Chronic lymphocytic leukemia (CLL); hematopoietic stem cell transplantation, chemotherapy (immunocompromised)	Progressive encephalitis, bilateral hearing loss	Tinnitus, sensorineural deafness, central dyspnea, hypotension, nausea, irritability, agitation	Fatal (within 7.5 mo)	VA1/HMO-C (NGS), VA1/HMO-C-UK1, KM 358468	Community acquired?	29
14-yr-old boy (France)	X-linked agammaglobulinemia (immunocompromised)	Progressive encephalitis	Progressive cognitive decline, recurring seizures, ataxia, erratic myoclonus, dysarthria	Alive at time of publication	VA1/HMO-C (NGS), VA1/HMO-C-PA, KM401565	Data not available	30
4-yr-old boy (Japan)	Congenital aplastic anemia, hematopoietic stem cell transplantation, chemotherapy (immunocompromised)	Encephalopathy	Cluster of convulsion, fever, diarrhea	Recovered	MLB1 (NGS), MLB1-NAGANO-1545, LC064152	Unknown	32
21-yr-old women (Switzerland)	Healthy	Acute meningitis	Severe headache, fever, neck stiffness	Recovered	MLB2 (NGS), MLB2/human/Geneva/2014, KT224358	Unknown (contact with children?)	33
37-year-old (Switzerland)	Acute myeloid leukemia, relapse hematopoietic stem cell transplantation, chemotherapy (immunocompromised)	Meningitis	Headache, meningeal involvement, vertigo, limb weakness, lightheadedness	Fatal (within 9 mo)	MLB2 (RT-PCR)	Unknown (contact with children?)	33
8-mo-old girl (UK)	Acute myeloid leukemia, hematopoietic stem cell transplantation, chemotherapy (immunocompromised)	Encephalitis	Encephalopathy, uncontrolled dystonic movement, poor respiratory effort	Fatal	VA1/HMO-C (NGS)	Unknown, no history of exposure to animals	31

*Abbreviations: VA/HMO, Virginia/human-mink-ovine-like astrovirus; NGS, next-generation sequencing.

1.5 to 2.5 years (17).

TABLE 2 Neurovirulent astrovirus infections associated with CNS disease in animals, in order of publication year (2004 to 2017)°

Host	Age (country)	Clinical signs	Natural outcome	Detection source	Method(s) of detection	Ni-AstV type	Abundance	Reference
Mink { <i>Mustela vison</i>)	3 to 4 mo (Denmark, Sweden, Finland)	Shaking syndrome (SMS), tremor/seizure, paraplegia	mink Mortality (27-28%)	Brain	NGS	MiAstV/MAstV10	0.2-0.8% morbidity	12, 16
Bovine (<i>Bos taurus</i>)	3 yr (USA)	Seizure, paraplegia, circling, and blindness	Not known	Brainstem, cerebellum, and/or spinal cord	NGS, ISH	BoAstV NeuroSI	3/32 animals (9.4%)	13
	1.5 to 7 yr	Abnormal gait	(or Not known)	Brain	NGS, ISH, BoAstV-CH1		6/22 affected	17, 43,

	(Switzerland)	recumbency), behavioral changes, and hyperreactivity or depression			tissue/medulla oblongata	RT-PCR	3 and/or BoAstV-CH1 5	animals, 0/33 healthy animals	45
m	4 (Switzerland)	yr Unspecified symptoms	CNS Not known		Medulla oblongata, cerebellar cortex, midbrain, cerebral cortex	NGS, RT-PCR	BoAstV-CH1 5	Single case	43
n									
>									
o									
o	1.5 to 10 yr (Switzerland)	Encephalitis	Deceased		Brain, medulla oblongata	NGS, RT-PCR	BoAstV-CH1 3/BoAstV NeuroSI	2/16 affected animals, 0/50 healthy animals	44
O									
C									
O	1 to >10 yr (Switzerland)	Encephalitis, meningoencephalitis	Not known		Brain, cerebellum, cerebrum, hippocampus	brainstem, ISH, RT-PCR	BoAstV-CH1 3/BoAstV NeuroSI	12/14 affected animals	18
	15 (Germany)	mo Central circling, inappétence, somnolence	blindness, Deceased		Brain, spleen, liver, pancreas	spinal cord, NGS, RT-PCR, ISH	BoAstV-BH8 9/14	Single case	47
	7 mo to 2 yr (Canada)	Encephalitis, seizures	Mortality (100%)		Brain	RT-PCR, ISH	BoAstV-CH1 3/BoAstV NeuroSI	2 single cases; 4/9 affected animals	48, 49
	1 to 11 yr (Switzerland)	Encephalitis, abnormalities, behavioral changes	gait Deceased		Gray matter of the CNS (brainstem), cerebellum, cerebrum, hippocampus	ISH, IHC	BoAstV-CH1 3/BoAstV NeuroSI	33/97 affected animals, 0/52 healthy animals	8
Ovine [Ovis aries)	4 yr or 10 days (Scotland)	Trembling, whole-body tremor, circling, recumbency	Not known		Cerebrum, cerebellum, obex, spinal cord, tonsil	NGS, RT-PCR	MAstV-13	Single cases	14
	7 (Switzerland)	yr Clinical nonsuppurative encephalitis	signs of Deceased		Medulla oblongata, cerebellum, thalamus, hippocampus, cortex, caudate nucleus	RT-PCR, IHC	MAstV13-CH 16	1/48 affected animals	51
Swine [Sus scrofa domestica)	1-6 (Sweden)	days Congenital type All	tremor Not known		Cerebrum, brainstem, cerebellum	RT-PCR	PoAstV-2 and PoAstV-5	3/3 affected animals, 3/3 healthy animals	52

3-5 wk (Hungarian pigs with presumed postweaning stress)	Paraplegia, pitching, flaccid paralysis, loss of consciousness	Mortality (100%)	Brainstem, cerebellum, spinal cord, tonsils, lung, nasal mucosa	RT-PCR, ISH	PoAstV-3	1.5-4% morbidity	15
5 wk (pigs and sows, USA)	Hind-limb weakness, quadriplegia, convulsions	Mortality (75-100%)	Cerebrum, cerebellum, spinal cord	NGS, RT-PCR	PoAstV-3	3/4 affected sows, 1/2 affected pigs	53

"Abbreviations: NGS, next-generation sequencing; ISH, in situ hybridization; IHC, immunohistochemistry; RT-PCR, reverse transcription-PCR.

Two years later, the same research group reported additional astrovirus-positive encephalitis cases from cattle, as determined by NGS and RT-PCR (43). This included a 4-year-old Swiss cow that had hyperreactivity and incoordination and was diagnosed with meningoencephalitis. The neurovirulent astrovirus (BoAstV-CH15; accession number [KT956903](#)) was genetically different from cattle BoAstV-CH13 strains within the cluster (VA1/HMO). By retrospective analysis, coinfection with BoAstV-CH13 and BoAstV-CH15 was detected in a case from the original 22 brain specimens from cattle with nonsuppurative encephalitis. Another specific study from Switzerland showed that 2 (12.5%) of 16 brain samples from cattle with unknown encephalitides, collected between 1995 and 2015, were astrovirus positive by NGS (44)(Table 2). The neurovirulent astroviruses from the 2- and 6-year-old diseased animals belonged to the BoAstV-CH13/BoAstV NeuroS1 astroviruses within the VA1/HMO cluster. None of 50 healthy control cattle were astrovirus positive by molecular methods (44). Taken together, the data show that around one quarter of the Swiss bovine population with nonsuppurative encephalitis of unresolved etiology had BoAstV-CH13 infection (45). Based on these results, formalin-fixed, paraffin-embedded (FFPE) brain tissues from histologically confirmed cases of European sporadic bovine encephalitis/encephalomyelitis between 1958 and 1976 were retested by ISH specific for BoAstV-CH13/BoAstV NeuroS1 in Switzerland (18). Astrovirus positivity was detected in 12 (86%) of the 14 cases (18). Using a combination of newly developed specific nucleic acid and viral protein detection methods (ISH and immunohistochemistry [IHC]), 33 (34%) of the 97 sporadic bovine encephalitis cases in Switzerland between 1985 and 2015 were BoAstV-CH13/BoAstV NeuroS1 positive, but none of the control group samples were positive (8). The mean age of the affected animals was 3.8 years in this study, and interestingly, all animals but one were female (8). Seasonality from the beginning of winter until the end of spring was observed, with peaks in November-December and March-April (8). Cattle with neurovirulent astrovirus had behavioral changes (e.g., aggression or anxiety), gastro-enteric symptoms (e.g., diarrhea or excessive salivation), and nystagmus (8). The most important conclusion from the Swiss studies is that using systematic investigation and specific methods to focus on the selected cases of bovine encephalitis increased the sensitivity; this underlines the importance and correlation of the causal relationship between neurological disease and astrovirus in the "post-Koch-postulate era" (8, 46).

Neurovirulent astrovirus infections in bovines were reported recently from Germany and Canada as well. In Germany, a 15-month-old cow died of encephalitis within 6 days after the onset of clinical signs (47). A neurovirulent astrovirus (BoAstV-BH89/14; accession number [LN879482](#)) was identified by NGS (47)(Table 2). Recent ISH and RT-PCR studies from Canada report that neurovirulent BoAstV-CH13/BoAstV NeuroS1 astroviruses are a common cause of encephalitis of unknown etiology (4 of 9 animals tested) in Canadian feedlot cattle (48, 49).

The most frequently observed clinical signs in Swiss bovines (n = 8) with astrovirus-associated encephalitis were decreased awareness of surroundings (87%), cranial nerve dysfunction (62%), such as dysphagia, decreased lingual and jaw tone, and reduced menace response and palpebral reflex, recumbency (62%), weakness, and tremor (50)(Table 2). Cell counts and protein concentration were increased in 44% and 60% of cerebrospinal fluid (CSF) samples, respectively. Astrovirus RNA was identified in only 1 of the 4 CSF samples. Other laboratory abnormalities (increased total and CSF protein concentrations, nonsuppurative pleocytosis, and increased creatinine kinase level) were non-specific (50). None of the affected bovines had a known immunocompromised status.

Ovines

A recent study reported cases of encephalitis and ganglionitis associated with astrovirus in domestic sheep from Scotland (14). In that study, a 4-year-old Welsh mountain ewe (from 2013) and a 10-day-old lamb (from 2014) from the same farm had progressive neurological signs (e.g., depressed sensorium, tremor, and unusual behavior) (Table 2). Nonsuppurative encephalomyelitis, particularly involving the cerebellar cortex and spinal cord, and dorsal root ganglionitis were detected by neurohistopathological methods. Nearly identical ovine astrovirus sequences (accession numbers [LT706530](#) and [LT706531](#)) within cluster VA1/HMO were detected from different tissue samples—including from the

CNS—although the cases presented 9 months apart (14). In addition, an ovine astrovirus sequence (accession number [KY859988](#)) was also reported for a 7-year-old Swiss white alpine ewe with nonsuppurative encephalitis in Switzerland (51)(Table 2).

Swine

Astrovirus as a possible cause of congenital tremor type AII in piglets (with brain and spinal cord demyelination) was first suggested in 2014 in a report from Sweden (52). Two lineages of partially characterized (only 296-bp-long RNA-dependent RNA polymerase [RdRp] gene) porcine astroviruses (accession numbers [KC790414](#) to [KC790418](#)), related to porcine astroviruses 2 and 5, were detected by nested RT-PCR in the cerebrum, brainstem, and cerebellum of affected and also healthy control piglets. The affected animals (n = 3) were 4 and 6 days old, while the control animals (n = 3) were only 1 day old (52). Furthermore, coinfection with porcine circovirus type 2 was found in healthy and diseased piglets (52). These preliminary results did not provide strong evidence for astrovirus as a cause of the disease.

Very recently (between 2011 and 2017), outbreaks of meningoencephalomyelitis associated with neurovirulent porcine astrovirus type 3 (PoAstV-3; accession numbers [KY073229](#) to [KY073232](#)), within the VA/HMO clade, were detected among 25- to 35-day-old domestic pigs in large, highly prolific swine farms in Hungary (15)(Table 2). Episodes of CNS disease of unknown etiology have persisted for 2 years at the index farm, and despite extensive decontamination efforts, sporadic cases are still present in 2018 (A. Boros and M. Albert, unpublished results). The disease has affected approximately 30 to 40 weaned pigs monthly (1.5% to 2% of the total), up to around 80 pigs (4%) in the autumn and winter seasons (15). Clinical symptoms typically appeared among weaned pigs 1 week after the weaning procedure (15). In stage 1, posterior leg weakness or paraplegia and pitching were visible. In stage 2, paralysis of both legs and skin pain were present. Later, in stage 3, paresis and serious spastic paralysis of muscles appeared, with loss of consciousness (15). Gastroenteric symptoms were not observed. Finally, in stage 3, the affected pigs were unable to eat or drink, and they died within a week due to exsiccosis (dehydration) (15). The neurovirulent astrovirus was detected in the cerebrum, brainstem, and spinal cord by RT-PCR, real-time PCR, and ISH analyses of samples from the affected group but not the healthy control animals (15).

Similar results were reported from the United States, where Ni-PoAstV-3 (accession number [KY940545](#)) was identified by NGS analysis of tissues from the CNS of piglets and sows with encephalomyelitis (53)(Table 2).

NEUROHISTOPATHOLOGY OF NEUROTROPIC ASTROVIRUS INFECTIONS

In general, the neurohistopathologic findings are consistent with nonsuppurative viral encephalitis. The most characteristic signs are severe neuronal degeneration, hypertrophic astrocytes, and infiltration of T lymphocytes and macrophages into the brain. Specifically, perivascular cuffs, composed of mononuclear cells, gliosis, and neuronal necrosis, are a common feature of all neurovirulent astrovirus infections (8,13, 15). The infection is present in different parts of the CNS (e.g., cerebrum, brainstem, and cerebellum), with variable severity, in all investigated host species (8, 13, 15). No inclusion bodies are observed.

Human Cases

The available neurohistopathological data are limited for human cases and are specific only to the brain. The human spinal cord was never investigated by histo-pathological methods in cases of neurovirulent astrovirus infection. Histopathology results are available for five VA1/HMO-C cases and the "classical" HAsV4 cases (9, 28-31). Biopsy of the frontal cortex found microgliosis and diffuse astrogliosis of white and gray matter, parenchymal and perivascular CD3+ T-cell infiltrates, axonal swelling, and neuronal loss in a boy with X-linked agammaglobulinemia (9). In another case, neuronal apoptosis, microglial reaction, and astrocytosis were observed in the brain biopsy specimen, without significant lymphocytic inflammation (28). Frontal lobe biopsy found gliosis, infiltration of CD3+/CD8+ lymphocytes, and astrovirus-infected neurons in the third case (29). In the fourth case, right frontal lobe biopsy found acute pan-encephalitis, microglial nodules, and perivascular infiltrates of CD3+/CD8+ lymphocytes (30). Scattered glial cells, perivascular and meningeal chronic inflammatory cell infiltrates, and large reactive astrocytes were identified in the fifth case caused by a VA1/HMO-C virus (31). In the case of HAsV4 infection, ventricular/leptomeningeal inflammatory infiltrates with macrophages and granulocytes and necrosis were detected in the brainstem, cerebellum, basal ganglia, and hippocampus (34).

Animal Cases

Mink. The lesions were most severe in the cerebellum and brainstem (Table 2) and consisted of neuronal necrosis (including neuronophagia), gliosis, perivascular cuffs with lymphocytes, plasma cells, and macrophages, and segmental loss of Purkinje cells (16).

Bovines. In bovines, encephalomyelitis cases were also characterized by neuronal degeneration and necrosis, gliosis, moderate to marked lymphohistiocytic perivascular cuffs, and meningitis with lymphocytes (13, 54). Nuclei of affected neurons were variably pyknotic, karyorrhectic, or karyomegalic, with absent or dispersed chromatin (13). Such lesions were most severe in the brainstem but were also present in the cerebellar cortex, the cerebrum, and the hippocampus (17, 18)(Table 2). The meningoencephalomyelitis was largely confined to gray matter in the CNS, with the most severe lesions in the cerebellar folia, brainstem, and spinal cord (13). Other parts of the brain were minimally (midbrain, thalamus, and basal nuclei) or not (cerebral cortex and underlying corona radiata) involved in inflammatory cell infiltrates (13).

Differences were also reported in certain studies. In one study, the histopathology revealed trigeminal ganglionitis with massive neuronal

necroses in both the brain and the ganglia (47). In another study, all layers of the hippocampus were seriously affected against the brainstem (18). In addition, viral RNA was also present in the medulla oblongata, cerebellar cortex, midbrain, and cerebral cortex (43)(Table 2). In general, the cerebellar folia had significant lesions with Purkinje cell necrosis and Bergmann glial and microglial proliferation (13). However, another study did not reveal astrovirus-positive Purkinje cells by ISH (8). Variations of the neuroanatomical distribution of the affected regions among individual animals might be associated with the possibility of nonrestricted neurovirulent astrovirus infection of gray matter of the CNS, with no preferred target region (8). It was also observed that the intensity of the lesions varied between individuals at different stages of the disease (8). This means that the stage of CNS infection may also have an effect on the results (8). This study also supports the probability of causality, as a close correlation of astrovirus and CNS lesions was found in most of the brain regions in cattle (8).

Swine. Histologically, moderate to marked perivascular lymphohistiocytic cell cuffing with marked vasculitis and neuronal degeneration, necrosis, and neurophagia with multifocal microgliosis and satellitosis were found in CNS lesions in pigs (15, 53). Neuronal necrosis was prominent in the gray matter of the cervical spinal cord but was also detected in neurons of the Purkinje layer (cerebellum), the medulla oblongata, cerebellar peduncles, and the midbrain (15). Necrotic neurons were variously swollen and hyper eosinophilic or shrunken with tinctorial changes, including faded, amphi-philic, or eosinophilic cytoplasm (15). Nuclei of affected neurons were pyknotic, kary-orrhectic, or losing border definition within the cytoplasm (15, 53). The intensity of the CNS lesions—as in cattle—was variable between individuals at different stages of the disease (Fig. 1).

For type AII congenital tremor, mild to moderate vacuolar changes of the white matter in the cerebrum, brain stem, and especially cerebellum were reported for pigs (52)(Table 2).




FIG 1 Representative sections from the anterior horn of the lumbar spinal cord for Ni-PoAstV-3-infected freshly weaned pigs with encephalomyelitis in stage 1 (posterior weakness) (A) and stage 3 (complete paralysis) (B) of the disease. The Ni-PoAstV-3-positive cells, which are presumed (based on the anatomical positions of the cells) to be motoneurons (arrows) and interneurons (arrowheads), were visualized using a digoxigenin-11-UTP (DIG)-labeled RNA probe and an *in situ* RNA hybridization technique. The 189-nt-long DIG-labeled antisense PoAstV-3 RNA probe targets the RdRp region of Ni-PoAstV-3 (accession numbers [KY073229](#) to [KY073232](#)). The sections were visualized using an anti-DIG Fab fragment conjugated to alkaline phosphatase (Roche) and NBT/BCIP solution (Roche) supplemented with 1 mM levamisole. The section in panel A was counterstained with FastRed. Note that Ni-PoAstV-3 ISH-positive apoptotic (shrinking) neurons (double arrows in inserts) are observable in all stages of the disease.

QUANTITATIVE AND MOLECULAR GENETIC ANALYSIS OF NEUROVIRULENT ASTROVIRUSES

Quantitative PCR Analysis

Viral loads in human brain biopsy specimens from a boy with X-linked agammaglobulinemia were as high as 1.53×10^7 RNA copies and were higher than those in postmortem samples (brain stem, 1.92×10^4 RNA molecules per reaction; cerebellum, 5.39×10^2 RNA molecules per reaction; and frontal lobe, 1.14×10^2 RNA molecules per reaction). No other postmortem samples were astrovirus positive (9). The brain tissue and the cerebrospinal fluid (CSF) usually contain viral RNA, but significantly lower viral loads were found in stool and serum, the latter of which confirmed viremia (28). The viral loads in brain biopsy specimens can be 103- and 106-fold higher than those in CSF or fecal samples, respectively (28). These and other results (30, 31) highlight that analysis of CSF or stools cannot be sufficient alone and that analysis of profound specimens (antemortem or postmortem brain biopsy specimens) may be required in order to make an appropriate laboratory diagnosis of neurovirulent astrovirus infection.

In ovines, the highest viral loads were detected in regions of the CNS (cerebrum, 2.16×10^7 /xl template; and cerebellum, 1.23×10^5 /xl template), including the obex (1.34×10^5 to 1.06×10^6 /xl template) and spinal cord (1.59×10^5 /xl template) (14). Other organs, such as the spleen, ileum, and pooled intestine (3.72×10^2 /xl template), showed remarkably low viral loads in comparison to those in organs of the CNS (14). Furthermore, low to moderate RNA loads were detected in lymphoid tissue of tonsils of sheep (14).

Similar results were reported for domestic pigs. The highest CNS viral loads were detected in the brain stem (1.85×10^6 to 7.43×10^7 /xg total RNA) and the cervical, thoracic, and lumbar spinal cord (1.47×10^6 to 3.39×10^7 /xg total RNA). However, moderate to high viral RNA loads were also found in the respiratory tract, such as the nasal mucosa (2.33×10^5 to 3.22×10^5 /xg total RNA), tonsil (9.08×10^2 to 4.65×10^4 /xg total RNA), and lung (1.34×10^2 to 5.01×10^2 /xg total RNA) (15). On the other hand, no or low copy numbers were detected in fecal

samples by nested RT-PCR (15).

Genomic Analysis of Neurovirulent Astroviruses

Due to their relatively recent discovery (in 2010) and the absence of model animals and cell cultures, little is known about the pathogenicity, replication cycle, or genomic features of Ni-AstVs, although the recently described Caco-2 cell-adapted Ni-HAstV strain VA1 may facilitate the cytopathogenic research on Ni-HAstVs (55). The majority of the current knowledge about the genome organization, viral peptide composition, and virion architecture of AstVs originated from experimental analysis of the cultivable turkey (Tu) and "classical" human astroviruses, mainly TuAstV-2, HAstV-1, HAstV-2, and HAstV-8, which have more than 40 years of scientific research background (56, 57). However, in silico genome analyses of Ni-AstVs and genome comparisons with the well-studied "classical" AstVs may reveal conserved sites and possible functional genome regions which may be important for the genome replication and even neurovirulence of Ni-AstVs. In this section, the results of in silico comparative genomic analyses of all Ni-AstV genomes are summarized. There are total of 26 Ni-AstV strains detected in CNS samples with known complete genome sequences or complete coding sequences (CDS) available in the GenBank database (as of 5 March 2018). The sequences originated from human (HAstV; n = 6), mink (MiAstV; n = 1), ovine (OvAstV; n = 3), bovine (BoAstV; n = 12), and porcine (PoAstV; n = 4) hosts located in Europe, the United States, and Japan (Table 3).

Phylogenetically, the majority (24/26 strains) of the Ni-AstVs belong to the VA/HMO phylogenetic clade (MAstV genogroup II), with the exception of Ni-HAstV-MLB strains (Fig. 2). Although other, non-HMO AstVs are also detectable from CNS samples, the dominance of Ni-AstVs in the VA/HMO clade suggests an increased neurotropic potential of this group of astroviruses compared to that of other AstVs. In the VA/HMO clade, the Ni-AstVs of the same host form distinct phylogenetic groups, with the exception of Ni-AstVs of bovine and ovine origins, which are present in a mixed group (Fig. 2) and suggest the capability of cross-species transmission of at least the Ni-BoAstVs (51, 58). Members of the same phylogenetic group may belong to the same AstV serotype/genotype, although the official classifications of the majority of Ni-AstVs are still absent.

Besides the phylogenetic distance, the genome organizations of all Ni-AstVs are essentially the same and follow the general astrovirus genome architecture, with three open reading frames (ORFs) flanked by 5' and 3' untranslated regions (UTRs) (Fig. 3)(57, 59). The 5'UTRs of AstVs, including the Ni-AstVs, predominantly start with the conserved pentamer CCAA (60), although there are some Ni-AstVs [such as MiAstV strain SMS-AstV, HAstV strain HMO-C-UK1(a)-London1, and HAstV-MLB2 strain MLB2/human/ Geneva/2014] in which this motif is missing, probably due to the incompletely determined 5' UTR (Fig. 3; Table 3). The full-length 5' UTRs of Ni-AstVs are relatively short, ranging from 14 nucleotides (nt) (HAstV-MLB1 strain NAGANO-1545; accession number [LC064152](#)) to 51 nt (BoAstV strain NeuroS1; accession number [KF233994](#)), and may contain a variable number of stem-loops (data not shown) (45).

An identity graph of the nt alignment (Fig. 3) of all (n = 26) Ni-AstVs revealed the presence of conserved regions, including the ribosomal frameshift signal (RFS; adenine hexamer, cysteine, and stem-loop), located in the junction of the ORFs, and the presumed subgenomic RNA (sgRNA) promoter sequence, located in the 3' end of ORF1b (57, 59, 61)(Fig. 3). Due to the presence of RFS, ORF1a and the adjacent ORF1b could be transcribed continuously (ORF1ab), resulting in production of nonstructural polyprotein 1ab (NSP-1ab), while ORF2 and the sgRNA copies could encode the capsid proteins (62). Besides the primary sequence conservation, the promoter sequence of sgRNA could be found in a conserved stem-loop structure (Fig. 3). Most of the conserved nt are located in ORF1b and the 5' end of ORF2 (Fig. 3).

The 3' UTRs of Ni-AstVs are relatively short, ranging from 32 nt (HAstV-MLB2 strain MLB2/human/Geneva/2014; accession number [KT224358.1](#)) to 101 nt (HAstV-PA strain Paris; accession number [KM401565.1](#)) without the poly(A) tail, and they may contain a variable number of stem-loops (45; data not shown). The highly conserved stem-loop II-like motif (s2m), identified in four different families of positive-sense single-stranded RNA (ssRNA) viruses—Astroviridae, Caliciviridae, Picornaviridae, and Coronaviridae (63, 64)—is predominantly present in the 3' end of the genome for all Ni-AstVs, with the exception of Ni-HAstV-MLB strains NAGANO1545 and MLB2/human/Geneva/2014, similar to other MLB AstVs (22). s2m is highly conserved at both the structural and

TABLE 3 Summary of currently known neurovirulent (identified from CNS samples) astroviruses with complete genome or coding sequences obtained from humans and animals

Host	Accession no.	Species/genotype	Strain/isolate	Genome completeness	Genome length (nt)	Detection source	Country	Reference
Bovine	KF233994	BoAstV NeuroS1	NeuroS1	3' end is missing	6,468	Brain tissue	USA	13
	KM035759	BoAstV-CH13	CH13	Complete	6,526	Medulla oblongata	Switzerland	17
	KX266901	BoAstV-CH13/NeuroS1	23871	Complete	6,526	Brainstem	Switzerland	45
	KX266902	BoAstV-CH13/NeuroS1	26730	Complete	6,525	Brainstem	Switzerland	45
	KX266903	BoAstV-CH13/NeuroS1	26875	Complete	6,526	Brainstem	Switzerland	45
	KX266904	BoAstV-CH13/NeuroS1	36716	Complete	6,526	Brainstem	Switzerland	45
	KX266905	BoAstV-CH13/NeuroS1	23985	Complete	6,526	Brainstem	Switzerland	45
	KX266906	BoAstV-CH13/NeuroS1	42799	Complete	6,525	Brainstem	Switzerland	45
	KX266907	BoAstV-CH13/NeuroS1	43661	Complete	6,526	Brainstem	Switzerland	45
	KX266908	BoAstV-CH13/NeuroS1	43660	Complete	6,526	Brainstem	Switzerland	45
KT956903.2	BoAstV	CH15	Complete	6,456	Medulla oblongata	Switzerland	43	
LN879482	BoAstV	BH89/14	Complete	6,450	Brain tissue	Germany	47	
Ovine	KY985988	MAstV-13	CH16	Complete	6,455	Brain tissue	Switzerland	51
	LT706530	MAstV-13	UK/2014/lamb/l1b01455	Complete	6,454	Organ pool containing brain, spinal cord, and spleen	UK	14
	LT706531	MAstV-13	UK/2013/ewe/l1b01454	Complete	6,454	Organ pool containing brain, spinal cord, and spleen	UK	14
Mink	GU985458	MiAstV/MAstV-10	SMS-AstV	5' end is missing	6,602	Brain tissue	Sweden	12
Swine	KY940545	PoAstV-3	USA/I A/7023/2017	Complete	6,430	Organ pool containing cerebrum, cerebellum, brainstem, and spinal cord	USA	53




FIG 2 Phylogenetic analysis of full-length capsid proteins (ORF2) of representative members of the family Astroviridae. The amino acid sequences were aligned with the MUSCLE algorithm (86), and evolutionary analysis was conducted in MEGA 6 (87), using an ML method based on the LG model with frequencies, a discrete gamma distribution (+G), and evolutionarily invariable sites (+I). Bootstrap values were determined for 1,000 replicates. The Ni-AstV strains with known full-length capsid proteins are indicated with red squares. The astrovirus clades representing the known Ni-AstVs are highlighted in orange. The Avastroviruses (accession numbers [AB033998](#), [FR727149](#), [HQ889774](#), [FR727146](#), [KJ020899](#), [JF414802](#), [Y15936](#), [AF206663](#), [EU143851](#), [FJ434664](#), and [FJ919228](#)) were chosen as a tree outgroup. Abbreviations:

BaAstV, bat astrovirus; BdAstV, bottlenose dolphin astrovirus; BoAstV, bovine astrovirus; CAstV, California sea lion astrovirus; CaAstV, canine astrovirus; ErAstV, European roller astrovirus; FeAstV, feline astrovirus; FoAstV, fox astrovirus; HAstV, human astrovirus; MiAstV, mink astrovirus; MuAstV, murine astrovirus; OvAstV, ovine astrovirus; PoAstV, porcine astrovirus; RaAstV, rabbit astrovirus; RAstV, rat astrovirus; WBAstV, wild boar astrovirus.





FIG 3 Representative genome map and identity graph (above the map) for Ni-AstVs. Vertical lines in the identity graph represent identical nucleotides (nt) in the alignment. Thicker lines represent multiple, consecutive identical nt. The identity graph was generated from the MUSCLE alignment of the complete genome/complete coding sequences of all known Ni-AstVs by use of UGENE ver 1.26 and CorelDraw ver. 12. The ruler above the identity graph represents nt positions in the alignment. The promoter sequence (gray background) of sgRNA is part of a conserved stem-loop (*). The secondary RNA structure of the sgRNA promoter was generated from the alignment of the corresponding genome region of all Ni-AstVs by use of the RNAalifold Web server (88) and VARNA visualization software, version 3.9. Note that s2m of the 3' end is missing in the Ni-HAstV strains of MLB-1 (accession number [LC064152](#)) and MLB-2 (accession number [KT224358](#)). ORF, open reading frame; RFS, ribosomal frameshift signal; NSP-1a, nonstructural polyprotein translated from ORF1a; NSP-1ab, nonstructural polyprotein continuously translated from ORF1a and ORF1b; sgRNA, subgenomic RNA.

sequence levels, is mostly present at the junction of ORF2 and the 3'UTR, and may contain the stop codon of ORF2 (CGMGGCCACGCCGAGUAGGAHCGAGGGUACAGS [the stop codon is underlined]), with the exception of Ni-MiAstV strain SMS-AstV (accession number [GU985458](#)) and Ni-BoAstV strain NeuroS1 (accession number [KF233994](#)). In Ni-MiAstV strain SMS-AstV, the stop codon is 11 nt upstream from s2m, and in Ni-BoAstV strain NeuroS1, the first nucleotide of s2m is uniquely U instead of C, which causes termination of ORF2 before s2m (data not shown). The first 15 nt of s2m (CGMGGCCACGCCGAG) encode the RGHAE amino acid (aa) motif, which is the highly conserved C-terminal 5 aa of the capsid protein in the majority of Ni-AstVs. Interestingly, identical or slightly modified (RTHAE) motifs are also present 15 aa upstream of the RGHAE motif of s2m in all Ni-BoAstVs (data not shown), which suggests the acquisition of two s2m by the ancestors of Ni-BoAstVs, similar to those found in certain corona- and picornaviruses (63). The exact role of s2m in the replication process of Ni-AstVs and other s2m-containing viruses is still unknown, although the conserved nature and wide distribution of the motif may suggest an important role in the replication process. The 3'UTRs of Ni-AstVs may contain deletions and sequence repeats, such as the G6382/6393AUUUCUUUNA sequence of Ni-PoAstV-3 of USA/IA/7023/ 2017, which is missing in the Hungarian Ni-PoAstV-3 strains (15).

Amino acid sequence comparisons of NSP-1ab and the capsid proteins transcribed from ORF1ab and ORF2 of all Ni-AstVs revealed the presence of highly conserved genomic regions located mainly in the presumed helicase (Hel), protease (Pro), and RNA-dependent RNA polymerase (RdRp) regions of NSP-1ab and the conserved, presumably assembly domain of the capsid (Fig. 4).

NSP-1ab (ORF1ab)



Capsid (ORF2)




FIG 4 Schematic maps and identity graphs (above the maps) for nonstructural protein 1ab (NSP-1ab) and capsid for all known Ni-AstVs. Vertical lines in the identity graphs represent identical amino acids in the alignment. Thicker lines indicate multiple, consecutive identical aa. The identity graphs were generated from the MUSCLE alignments of NSP-1ab and capsid proteins of all known Ni-AstVs with complete genome sequences or complete CDS by use of UGENE ver 1.26 and CorelDraw ver. 12. The rulers above the identity graphs represent aa positions in the alignments. Hel, helicase; cc 1 to 3, coiled coils; TM 1 to 7, transmembrane domains; VPg, genome-associated viral protein; NLS, nuclear localization sequence; HVR, hypervariable region; RdRp, RNA-dependent RNA polymerase; RBD, RNA-binding domain. The red arrow shows a possible trypsin cleavage site, and yellow arrows show possible caspase cleavage sites.

In NSP-1ab, the conserved aa motifs of Pro (65-68) as well as the two coiled coils (CC) and the multiple transmembrane (TM) helices upstream of Pro described for the "classical" HAstVs are present in all Ni-AstVs (57)(Fig. 4). Interestingly, NSP-1ab of Ni-AstVs may contain seven TM domains (Fig. 4 and 5) upstream of Pro, as predicted by the MEMSAT software of the PSIPRED Protein Sequence Analysis Workbench (69, 70), compared to the five or six TM helices of "classical" HAstVs (57), although no insertions were found in the corresponding genome regions of Ni-AstVs compared to those of "classical" AstVs.




FIG 5 Schematic representation of seven predicted transmembrane domains (yellow columns; TM1 to -7) identifiable in the N-terminal region of NSP-1ab of all known Ni-AstVs. Numbers in parentheses and italics indicate the first (numbers in the upper part) and last (numbers in the lower part) aa positions of the given TM domain. The predictions are based on the sequences listed on the left side of the figure and were determined using the MEMSAT software of the PSIPRED Protein Sequence Analysis Workbench (69, 70). All Ni-AstVs contain essentially the same number and topology of TM domains (data not shown).

One of the well-conserved genome regions of Ni-AstVs is located downstream of the second CC and contains the conserved RExx(I/V)LRxE motif (x = variable aa) of unknown function (Fig. 4). Furthermore, besides the well-known two CCs located in the presumed Hel region and between the Pro and VPg regions of all AstVs, a third CC is predictable in the hypervariable region (HVR) of Ni-AstVs (Fig. 4 and 6) by using the coiled-coil prediction software of ExpASy (71, 72). The HVR of "classical" HAstVs may contain several insertions/deletions and is known to be associated with the adaptation of certain cell lines, different RNA replication and growth rates, and different viral RNA levels in feces (73, 74). The exact role of the HVR in the replication cycle or neuropathogenic potential of Ni-AstVs is currently unknown, although the presence of a CC structure—CC proteins are usually involved in important biological functions, such as transcriptional control (75)—and some highly conserved aa [Gxx(P/L)FxQR] (Fig. 4) may indicate a pivotal role of this region.

The identity graph for all capsids of the 26 Ni-AstVs shows the presence of two major parts, an N-terminal conserved part and a C-terminal variable part, which may correspond to the "conserved" and "hypervariable" capsid regions of "classical" HAstVs (57, 76, 77)(Fig. 4). The N-terminal parts of the capsids of Ni-AstVs have a variable primary sequence (Fig. 7), although this region of all Ni-AstVs is equally rich in the basic aa lysine (K), arginine (R), and histidine (H), and therefore this region may act as an RNA-binding domain (RBD) required in the encapsidation process (57, 76, 77)(Fig. 4 and 7A). Interestingly, short glutamine-rich (Q-rich) motifs with unknown functions were identifiable in this region in the majority of Ni-AstVs of the VA/HMO clade (Fig. 7A). The conserved N-terminal part of the capsid may act as an assembly or core domain mainly responsible for the formation of the structural core of the virion, while the highly immunogenic hypervariable region is located on the surface of the viral particle and may be responsible for receptor interaction and the strain-specific tropism of the virus (76-78). The maturation process of AstV virions includes the trypsin-mediated cleavage of the capsid polyprotein between the conserved and hypervariable regions of "classical" HAstVs (79, 80). Trypsin is expressed predominantly in the gastroenteric tract. Currently, there is no experimental evidence about the role of trypsin in the maturation of Ni-AstV virions, although the aa alignment of the capsids of "classical" AstVs and Ni-AstVs revealed the potential presence of conserved arginine (R) and lysine (K) residues, in line with experimentally identified trypsin cleavage sites of "classical" HAstVs, in all Ni-AstVs (Fig. 7B). However, the first recent experimental investigation showed no significant effect of exogenous trypsin treatment on in vitro cultures of Ni-HAstV strain VA1 originated from human stool (55). The in vivo proteolytic processing


of Ni-AstVs is unclear. If this group of astroviruses does not absolutely require trypsin-dependent proteolytic activation, then Ni-AstVs may not depend on and be restricted to the gastrointestinal tract.

The potentially surface-exposed hypervariable region of the capsid of Ni-AstVs contains only nine, scattered identical aa identifiable in the MUSCLE alignment of capsid sequences (Fig. 4), which may be responsible for neuronal infectivity, although further sequences of Ni-AstVs and experimental evidence are needed to confirm this hypothesis.


The C-terminal parts of the capsid sequences of Ni-AstVs are rich in acidic aa, such as aspartic acid (D) and glutamic acid (E), similar to those in "classical" HAstVs. The acidic regions of the C-terminal end contain several presumed caspase cleavage sites (e.g., DXXD and XEXD), similar to those found in the "classical" HAstVs (81, 82). Interestingly, the conserved motif Q(I/L)QxR(F/Y), adjacent to the D/E-rich acidic section of unknown function, was identifiable in the majority of Ni-AstVs of the HMO clade (Fig. 7C).

In summary, the topology of the Ni-AstVs in the phylogenetic tree (Fig. 2) suggests a common lineage of neurovirulent astroviruses; however, the determination of the specific nt and aa motifs that are involved in neuroinvasion warrants further experimental investigation.


Ni-BoAstV strain NeuroS1 - NSP-1ab [AGO50637]




Ni-MiAstV strain SMS-AstV - NSP-1ab [ADR65075]




Ni-OvAstV strain lib01455 - NSP-1ab [SIP85305]



Ni-HAstV-PA strain Paris - NSP-1ab [AKI81871]



Ni-PoAstV3 strain NI-Brain9-2016a - NSP-1ab [ARR73566]



Ni-HAstV-MLB1 strain NAGANO1545 - NSP-1ab [BAU68079-80]




FIG 6 Representative probability (y axis) diagrams for predicted coiled-coil structures and their localizations in the aa sequences of NSP-1ab (x axis). The diagrams were generated using Ni-AstV reference sequences (indicated above the diagrams) and screening windows of 14, 21, and 28 aa (blue, green, and red lines) by use of the coiled-coil prediction software of ExPASy (71, 72). All other Ni-AstV sequences showed essentially the same localization of coiled coils (data not shown).




FIG 7 (A) Amino acid alignment of the N-terminal ends (RNA-binding domain [RBD]) of representative capsid sequences of neurovirulent (Ni) AstVs. Conservative and basic amino acids (R, K, and H) are shown with black and green backgrounds. Glutamine (Q) residues in Q-rich regions are marked with a deep yellow background and red boxes. Amino acid positions are shown according to the scale above the sequences. (B) Amino acid alignment and presumed trypsin cleavage sites (arrows) of the junctions of the conserved and hypervariable regions of representative neurovirulent (Ni) and "classical" AstV capsids. Red squares indicate N-terminal aa sequences of VP26 (HAstV2) and VP27/VP28 (HAstV8), experimentally identified by N-terminal sequencing (80, 89). Identical and highly conserved aa are marked with a black background in the alignment. Colored arrows indicate the conserved arginine (R) and lysine (K) residues which may act as potential trypsin cleavage sites (90), among which the conserved R/K residues marked with a green background and a red arrow are the most likely to be active. Numbers in parentheses indicate the first and last aa positions in the capsid polyprotein. (C) Representative amino acid alignment of the surroundings of the conserved acidic section (blue boxes), located in the C-terminal hypervariable regions, for all known neurovirulent (Ni; VA/HMO clade) and "classical" AstV capsids. Conservative and acidic amino acids (inside blue boxes) are shown with black and yellow backgrounds, respectively. Putative caspase cleavage sites of HAstV8 are marked in bold and underlined (81, 82). The conserved genome region that contains the Q(I/L)QxR(F/Y) motif, which is predominantly present among Ni-AstVs, is marked with a red box. Numbers in parentheses indicate the first and last aa positions in the capsid polyprotein.

POSSIBLE TRANSMISSION AND PATHOGENESIS OF NEUROVIRULENT ASTROVIRUSES

It is a general "dogma" that astroviruses are transmitted by the fecal-oral route, as they are associated with enteric infections and shedding in the feces. For humans, it has therefore also been postulated that invasion of the CNS occurs via the enteric tract as a consequence of immunodeficiency. This concept is supported by the finding that the VA1/HMO-C astroviruses have also been identified in the feces of a diseased patient as well as in healthy individuals (43). For humans, close contact with animals (mink) and young children and medical treatments (intravenous immunoglobulin and stem cell grafts) have all been suggested as possible—but not confirmed—sources of infection (9). While systematic investigations related to all potential sources and transmission modes of neurovirulent astrovirus are not available, there are some interesting novel findings which suggest the potential for extraintestinal virus shedding in different hosts. (i) The neurotropic astrovirus strains have not yet been identified in cattle feces (43). (ii) The human astrovirus VA1 was identified by NGS in a nasopharyngeal specimen from a

child with respiratory disease of unknown origin (83). (iii) Furthermore, Ni-HAstV was also detectable in throat swab and urine samples from an immunocompromised patient with HAstV-MLBI-associated acute encephalopathy (32). (iv) Neurovirulent PoAstV-3 (VA/HMO clade) was detected in high viral loads in nasal mucosa and tonsil samples from domestic pigs with encephalomyelitis, but no or consistently lower copy numbers were found in fecal samples from the affected animals by nested RT-PCR (15). (v) Overall, 33 (32%) nasal swabs and none of the 24 analyzed anal swabs, collected as sample pairs from 18- to 21-day-old healthy pigs on four different swine farms with no history of encephalitis in Hungary, were positive for PoAstV-3 by nested RT-PCR (Boros et al., unpublished data). These results indicate that certain astroviruses may be capable of being shed extraintestinally, and therefore investigations should be extended to the respiratory tract for determination of the possible respiratory source and/or route of infections.

The pathogenesis of and influential risk factors (host, virus, and environmental) for the neurovirulent astroviruses are unknown. Immunosuppression and an immunocompromised condition are common underlying diseases in neurovirulent astrovirus cases in humans and weaned pigs (15,19). Animals in highly prolific and intensive production management environments are often exposed to stressors which may also cause immunosuppression in cattle and pigs (15, 43).

LABORATORY DIAGNOSIS OF NEUROVIRULENT ASTROVIRUS INFECTIONS

There are several classical and modern laboratory methods for detection of classical (human) astroviruses, especially from feces (84). However, these methods are limited for the detection of the newly discovered group of neurovirulent astroviruses. Neither the genetic diversity of the circulating astroviruses nor the specificity of the available tests is known for neurovirulent astroviruses. In addition, it seems that the optimal specimen(s) for detection of neurovirulent astrovirus infection is probably something (e.g., CNS biopsy specimen or respiratory tract samples) other than feces (see Possible Transmission and Pathogenesis of Neurovirulent Astroviruses).

There is limited knowledge related to the *in vitro* tissue culture propagation of neurovirulent astroviruses. A recent study reported, for the first time, that the human astrovirus VA1 of stool origin can grow in cell cultures (without cytopathic effect), including cultures of human enteric Caco-2, human embryonic kidney HEK293T, and human respiratory A549 cells but not those of BHK-21 (baby hamster kidney) cells (55). Interestingly, VA1 could be cultured equally well with and without the addition of exogenous trypsin (55). There is no report of the propagation of VA1/HMO-C strains directly from the affected CNS tissues.

At present, electron microscopy (EM), ISH, RT-PCR, real-time PCR (with strain-specific primer pairs), and viral metagenomics/NGS research-based methods are available for detection of neurovirulent astroviruses; however, there are no commercially available assays. A hyperimmune antibody against the conserved N terminus of the viral capsid protein was recently developed for detection of BoAstV-CH13/BoAstV NeuroSI infections in cattle brain tissue samples (51). This immunohistochemistry (IHC) assay is sensitive and specific (51) for BoAstV-CH13/BoAstV NeuroSI; however, it is not available for commercial use.

Development and use of a sensitive routine laboratory test (e.g., using an enzyme-linked immunosorbent assay [ELISA], immunochromatography, or molecular method) for detection of neurovirulent astroviruses is necessary, especially for veterinary medicine.

THERAPY

Combinations of intravenous immunoglobulin (IVIG), methylprednisolone, ribavirin, and pegylated interferon (PEG-IFN) alpha-2b were applied as nonspecific antiviral treatments in a human case of neurovirulent astrovirus infection (30). In that study, progression of motor function and mild memory improvement were found at month 12 of follow-up (30). On the other hand, there was no response to ribavirin and IVIG therapy (after the patient previously received steroids and IVIG) in another fatal human case of neurovirulent astrovirus encephalitis (29). At present, there is no specific antiviral therapy for neurovirulent astrovirus infection of either humans or animals.

FUTURE PERSPECTIVE

Astrovirus was discovered in a fecal sample in 1975 (85). Until now, astrovirus has been thought to be an infectious agent only of the gastroenteric tract in humans and animals. The most recent and advanced high-throughput molecular techniques have changed our knowledge of astrovirus infection (84). The possibility of an extraintestinal infection or complication of astrovirus infection was first suggested in 2010 (9). Between 2010 and 2017, significant numbers of studies reported an association between astrovirus and CNS infection. To date, this association has been reported for humans and four mammalian species. These results demonstrate a previously unrecognized neurotropism of neurovirulent astrovirus, a causative agent of neuroinfections, such as encephalitis, meningitis, myelitis, and ganglionitis. The first studies not only support the association but also suggest that certain astroviruses may be relatively common infectious agents of the CNS in cases of undifferentiated viral encephalitis. Retrospective and prospective systematic investigations of astrovirus should be performed, focusing on cases and outbreaks of unknown encephalitis—whose economic cost is profound for cases in both humans and domestic animals. It would also be interesting to determine if all astroviruses have an affinity for the CNS or if this affinity is restricted to certain neurovirulent astrovirus lineages.

Astrovirus should be considered in the differential diagnosis of CNS diseases, including encephalitis, especially for

nonimmunocompetent patients. In addition, we cannot exclude the possibility that a certain proportion of unknown encephalitis cases in domesticated mammals are also related to astrovirus. But timely diagnosis is hindered by the lack of available laboratory assays to rapidly survey the potential agents responsible for encephalitis. Development of a specific routine laboratory test is necessary for detection of astroviruses in encephalitis and other CNS infections. In particular, an astrovirus detection assay should be included in the diagnostic armory and routinely used in laboratories of veterinary medicine.

The optimal specimen type for laboratory diagnosis of neurovirulent astrovirus infection is also problematic. Based on the present data, no or very low levels of astrovirus RNA are present in the feces, serum, and CSF, which leads to misdiagnosis of neurological infections. Invasive brain biopsy for correct diagnosis of astrovirus encephalitis clearly has related medical and ethical issues. A new report suggests that nasal and respiratory specimens are better and noninvasive samples for laboratory detection of neurovirulent astrovirus (15). According to the present research, a broad range of specimens should be collected for detection of astrovirus in CNS diseases.

The first reports indicated that the neurotropic astroviruses are geographically widespread in human, mink, bovine, and porcine species and that epidemic spread is also possible (12, 15). In addition, while neurovirulent astrovirus infections have been reported for humans and four mammalian species, it is possible that this type of infection occurs in other domestic and wild animal species as well. Extended investigations of known hosts and other related mammals (e.g., wild boar, roe deer, goat, etc.) with unknown meningoencephalitis causes should also be done. The first reports have drawn our attention to neurovirulent astrovirus infections. More comprehensive epidemiological studies on the clinical presentation, prevalence, distribution, extraintestinal transmission, immunology, virology, and diversity of neurovirulent astrovirus infections, with determination of complete genome sequences of further neurovirulent astroviruses, should be performed. The detailed pathogenesis of neurovirulent astrovirus infections is also lacking, and exogenous and endogenous factors determining the tropism and neuroinvasion of these viruses are still unknown. In vitro cell culture and in vivo animal models (e.g., pig and mink models) would facilitate studies of the neurotropism, replication sites, neuropathogenesis, and immune control of astroviruses.

In the future, more attention should be paid to the specific prevention of CNS infections, especially among domesticated animals, and to the pathogenesis and epidemiological investigation of neurovirulent astroviruses. Development and use of a specific vaccine against neurovirulent astroviruses would be an important, logical, and probably cost-effective step, especially for domesticated animals (e.g., pigs). Astroviruses show typical features suitable for a "one health" approach, in which human and animal health can be discussed together. Future research into astrovirus infections should include collaborative, multidisciplinary efforts in both veterinary and human medicine. Human and veterinary physicians should be aware of astrovirus infection as a potential differential diagnosis with corresponding signs of neurologic disease.

ACKNOWLEDGMENTS

This work was supported by a grant from the Hungarian Scientific Research Fund (grant OTKA/NKFIH K111615). P.P. and A.B. were supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences.

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

REFERENCES

1. Glaser CA, Honarmand S, Anderson LJ, Schnurr DP, Forghani B, Cossen CK, Schuster FL, Christie LJ, Tureen JH. 2006. Beyond viruses: clinical profiles and etiologies associated with encephalitis. *Clin Infect Dis* 43: 1565-1577. <https://doi.org/10.1086/509330>.
2. Long MT. Overview of meningitis, encephalitis, and encephalomyelitis. MSD veterinary manual. <http://www.msdsvetmanual.com/nervous-system/meningitis,-encephalitis,-and-encephalomyelitis/overview-of-meningitis,-encephalitis,-and-encephalomyelitis>. Accessed 2 November 2017.
3. World Organization for Animal Health. 2004. Manual of diagnostic tests and vaccines for terrestrial animals, 5th ed. World Organization for Animal Health, Paris, France.
4. Saylor D, Thakur K, Venkatesan A. 2015. Acute encephalitis in the immunocompromised individual. *Curr Opin* 28:330-336. <https://doi.org/10.1097/YCO.000000000000169>.
5. Glaser CA, Gilliam S, Schnurr D, Forghani B, Honarmand S, Khetsuriani N, Fischer M, Cossen CK, Anderson LJ. 2003. In search of encephalitis etiologies: diagnostic challenges in the California Encephalitis Project, 1998-2000. *Clin Infect Dis* 36:731-742. <https://doi.org/10.1086/367841>.
6. Kennedy PGE, Quan PL, Lipkin WI. 2017. Viral encephalitis of unknown cause: current perspective and recent advances. *Viruses* 9:138. <https://doi.org/10.3390/v9060138>.
7. Vora NM, Holman RC, Mehal JM, Steiner CA, Blanton J, Sejvar J. 2014. Burden of encephalitis-associated hospitalizations in the United States, 1998-2010. *Neurology* 82:443-451. <https://doi.org/10.1212/WNL.000000000000086>.
8. Selimovic-Hamza S, Boujon CL, Hilbe M, Oevermann A, Seuberlich T. 2017. Frequency and pathological phenotype of bovine astrovirus CH13/ NeuroSI infection in neurologically-diseased cattle: towards assessment of causality. *Viruses* 9:12.

<https://doi.org/10.3390/v9010012>.

9. Quan PL, Wagner TA, Briese T, Torgerson TR, Hornig M, Tashmukhame-dova A, Firth C, Palacios G, Baisre-De-Leon A, Paddock CD, Hutchison SK, Egholm M, Zaki SR, Goldman JE, Ochs HD, Lipkin WI. 2010. Astrovirus encephalitis in boy with X-linked agammaglobulinemia. *Emerg Infect Dis* 16:918-925. <https://doi.org/10.3201/eid1606.091536>.
10. Calistri A, Palu G. 2015. Editorial commentary: unbiased next-generation sequencing and new pathogen discovery: undeniable advantages and still-existing drawbacks. *Clin Infect Dis* 60:889-891. <https://doi.org/10.1093/cid/ciu913>.
11. Schubert RD, Wilson MR. 2015. A tale of two approaches: how metagenomics and proteomics are shaping the future of encephalitis diagnostics. *Curr Opin Neurol* 28:283-287. <https://doi.org/10.1097/WCO.0000000000000198>.
12. Blomstrom AL, Widen F, Hammer AS, Belak S, Berg M. 2010. Detection of a novel astrovirus in brain tissue of mink suffering from shaking mink syndrome by use of viral metagenomics. *J Clin Microbiol* 48:4392-4396. <https://doi.org/10.1128/JCM.01040-10>.
13. Li L, Diab S, McGraw S, Barr B, Traslavina R, Higgins R, Talbot T, Blanchard P, Rimoldi G, Fahsbender E, Page B, Phan TG, Wang C, Deng X, Pesavento P, Delwart E. 2013. Divergent astrovirus associated with neurologic disease in cattle. *Emerg Infect Dis* 19:1385-1392. <https://doi.org/10.3201/eid1909.130682>.
14. Pfaff F, Schlottau K, Scholes S, Courtenay A, Hoffmann B, Höper D, Beer M. 2017. A novel astrovirus associated with encephalitis and ganglionitis in domestic sheep. *Transbound Emerg Dis* 64:677-682. <https://doi.org/10.1111/tbed.12623>.
15. Boros A, Albert M, Pankovics P, Birö H, Pesavento PA, Phan TG, Delwart E, Reuter G. 2017. Outbreaks of neuroinvasive astrovirus associated with encephalomyelitis, weakness, and paralysis among weaned pigs, Hungary. *Emerg Infect Dis* 23:1999-2010. <https://doi.org/10.3201/eid2312.170804>.
16. Gavier-Widen D, Brojer C, Dietz HH, Englund L, Hammer AS, Hedlund KO, Hard af Segerstad C, Nilsson K, Nowotny N, Puurula V, Thoren P, Uhlhorn H, Weissenböck H, Agren E, Klingeborn B. 2004. Investigations into shaking mink syndrome: an encephalomyelitis of unknown cause in farmed mink (*Mustela vison*) kits in Scandinavia. *J Vet Diagn Invest* 16:305-312. <https://doi.org/10.1177/104063870401600408>.
17. Bouzalas IG, Wüthrich D, Walland J, Drögemüller C, Zubriggen A, Vande-velde M, Overmann A, Bruggmann R, Seuberlich T. 2014. Neurotropic astrovirus in cattle with nonsuppurative encephalitis in Europe. *J Clin Microbiol* 52:3318-3324. <https://doi.org/10.1128/JCM.01195-14>.
18. Selimovic-Hamza S, Bouzalas IG, Vandevelde M, Oevermann A, Seuberlich T. 2016. Detection of astrovirus in historical cases of European sporadic bovine encephalitis, Switzerland 1958-1976. *Front Vet Sci* 3:91. <https://doi.org/10.3389/fvets.2016.00091>.
19. Vu DL, Bosch A, Pinto RM, Guix S. 2017. Epidemiology of classic and novel human astrovirus: gastroenteritis and beyond. *Viruses* 9:33. <https://doi.org/10.3390/v9020033>.
20. Finkbeiner SR, Allred AF, Tarr PI, Klein EJ, Kirkwood CD, Wang D. 2008. Metagenomic analysis of human diarrhea: viral detection and discovery. *PLoS Pathog* 4:e1000011. <https://doi.org/10.1371/journal.ppat.1000011>.
21. Finkbeiner SR, Holtz LR, Jiang Y, Rajendran P, Franz CJ, Zhao G, Kang G, Wang D. 2009. Human stool contains a previously unrecognized diversity of novel astroviruses. *Virol J* 6:161. <https://doi.org/10.1186/1743-422X-6-161>.
22. Jiang H, Holtz LR, Bauer I, Franz CJ, Zhao G, Bodhidatta L, Shrestha SK, Kang G, Wang D. 2013. Comparison of novel MLB-clade, VA-clade and classic human astroviruses highlights constrained evolution of the classic human astrovirus nonstructural genes. *Virology* 436:8-14. <https://doi.org/10.1016/j.virol.2012.09.040>.
23. Finkbeiner SR, Li Y, Ruone S, Conrardy C, Gregoricus N, Toney D, Virgin HW, Anderson LJ, Vinje J, Wang D, Tong S. 2009. Identification of a novel astrovirus (astrovirus VA1) associated with an outbreak of acute gastroenteritis. *J Virol* 83:10836-10839. <https://doi.org/10.1128/JVI.00998-09>.
24. Kapoor A, Li L, Victoria J, Oderinde B, Mason C, Pandey P, Zaidi SZ, Delwart E. 2009. Multiple novel astrovirus species in human stool. *J Gen Virol* 90:2965-2972. <https://doi.org/10.1099/vir.0.014449-0>.
25. Phan TG, Nordgren J, Ouermi D, Simpore J, Nitiema LW, Deng X, Delwart E. 2014. New astrovirus in human feces from Burkina Faso. *J Clin Virol* 60:161-164. <https://doi.org/10.1016/j.jcv.2014.03.024>.
26. Holtz LR, Bauer IK, Rajendran P, Kang G, Wang D. 2011. Astrovirus MLB1 is not associated with diarrhea in a cohort of Indian children. *PLoS One* 6:e28647. <https://doi.org/10.1371/journal.pone.0028647>.
27. Meyer CT, Bauer IK, Antonio M, Adeyemi M, Saha D, Oundo JO, Ochieng JB, Omere R, Stine OC, Wang D, Holtz LR. 2015. Prevalence of classic, MLB-clade and VA-clade astroviruses in Kenya and The Gambia. *Virol J* 12:78. <https://doi.org/10.1186/s12985-015-0299-z>.
28. Brown JR, Morfopoulou S, Hubb J, Emmett WA, Ip W, Shah D, Brooks T, Paine SM, Anderson G, Virasami A, Tong CY, Clark DA, Plagnol V, Jaques TS, Qasim W, Hubank M, Breuer J. 2015. Astrovirus VA1/HMO-C: an increasingly recognized neurotropic pathogen

- in immunocompromised patients. *Clin Infect Dis* 60:881-888. <https://doi.org/10.1093/cid/ciu940>.
29. Naccache SN, Peggs KS, Mattes FM, Phadke R, Garson JA, Grant P, Samayoa E, Federman S, Miller S, Lunn MP, Gant V, Chiu CY. 2015. Diagnosis of neuroinvasive astrovirus infection in an immunocompromised adult with encephalitis by unbiased next-generation sequencing. *Clin Infect Dis* 60:919-923. <https://doi.org/10.1093/cid/ciu912>.
 30. Frémond ML, Perot P, Muth E, Cros G, Dumarest M, Mahlaoui N, Seilhean D, Desguerre I, Hebert C, Corre-Catelin N, Neven B, Lecuit M, Blanche S, Picard C, Eliot M. 2015. Next-generation sequencing for diagnosis and tailored therapy: a case report of astrovirus-associated progressive encephalitis. *J Pediatr Infect Dis Soc* 4:e53-e57. <https://doi.org/10.1093/jpids/piv040>.
 31. Lum SH, Turner A, Guiver M, Bonney D, Martland T, Davies E, Newbould M, Brown J, Morfopoulou S, Breuer J, Wynn R. 2016. An emerging opportunistic infection: fatal astrovirus (VA1/HMO-C) encephalitis in a pediatric stem cell transplant recipient. *Transpl Infect Dis* 18:960-964. <https://doi.org/10.1111/tid.12607>.
 32. Sato M, Kuroda M, Kasai M, Matsui H, Fukuyama T, Katano H, Tanaka-Taya K. 2016. Acute encephalopathy in an immunocompromised boy with astrovirus-MLB1 infection detected by next generation sequencing. *J Clin Virol* 78:66-70. <https://doi.org/10.1016/j.jcv.2016.03.010>.
 33. Cordey S, Vu D-L, Schibler M, L'Huillier AG, Brito F, Docquier M, Posfay-Barbe KM, Petty TJ, Turin L, Zdobnov EM, Kaiser L. 2016. Astrovirus MLB2, a new gastroenteric virus associated with meningitis and disseminated infection. *Emerg Infect Dis* 22:846-853. <https://doi.org/10.3201/eid2205.151807>.
 34. Wunderli W, Meerbach A, Güngör T, Berger C, Greiner O, Caduff R, Trkola A, Bossart W, Gerlach D, Schibler M, Cordey S, McKee TA, Van Belle S, Kaiser L, Tapparel C. 2011. Astrovirus infection in hospitalized infants with severe combined immunodeficiency after allogeneic hematopoietic stem cell transplantation. *PLoS One* 6:e27483. <https://doi.org/10.1371/journal.pone.0027483>.
 35. Vu DL, Cordey S, Brito F, Kaiser L. 2016. Novel human astroviruses: novel human diseases? *J Clin Virol* 82:56-63. <https://doi.org/10.1016/j.jcv.2016.07.004>.
 36. Fankhauser R. 1961. Sporadic meningo-encephalomyelitis in cattle. *Schweiz Arch Tierheilkd* 103:225-235. (In German.)
 37. Munday BL, Mason RW, Cumming R. 1973. Observations on diseases of the central nervous system of cattle in Tasmania. *Aust Vet J* 49:451-455. <https://doi.org/10.1111/j.1751-0813.1973.tb09290.x>.
 38. Billing J. 1974. Epizootic investigations on sporadically non-suppurative encephalomyelitis in cattle. PhD thesis. University of Munich, Munich, Germany. (In German.)
 39. Jeffrey M, Wilesmith JW. 1992. Idiopathic brainstem neuronal chromatinolysis and hippocampal sclerosis: a novel encephalopathy in clinically suspect cases of bovine spongiform encephalopathy. *Vet Rec* 131: 359-362. <https://doi.org/10.1136/vr.131.16.359>.
 40. Theil D, Fatzer I, Schiller P, Caplazi A, Zurbriggen M, Vandeveld M. 1998. Neuropathological and aetiological studies of sporadic non-suppurative meningoencephalitis of cattle. *Vet Rec* 143:244-249. <https://doi.org/10.1136/vr.143.9.244>.
 41. Bozzetta E, Caramelli M, Casalone C, Acutis PL, Ru G. 2003. BSE surveillance in Italy: neuropathological findings in cattle in the frame of the passive surveillance programme. *J Vet Med A Physiol Pathol Clin Med* 50:48-49. <https://doi.org/10.1046/j.1439-0442.2003.00488.x>.
 42. Sanchez S, Clark EG, Wobeser GA, Janzen ED, Philibert H. 2013. A retrospective study of non-suppurative encephalitis in beef cattle from western Canada. *Can Vet J* 54:1127-1132.
 43. Seuberlich T, Wüthrich D, Selimovic-Hamza S, Drögemüller C, Oevermann A, Bruggmann R, Bouzalas I. 2016. Identification of a second encephalitis-associated astrovirus in cattle. *Emerg Microbes Infect* 5:e5. <https://doi.org/10.1038/emi.2016.5>.
 44. Wüthrich D, Boujon CL, Truchet L, Selimovic-Hamza S, Oevermann A, Bouzalas IG, Bruggmann R, Seuberlich T. 2016. Exploring the virome of cattle with non-suppurative encephalitis of unknown etiology by metagenomics. *Virology* 493:22-30. <https://doi.org/10.1016/j.virol.2016.03.009>.
 45. Bouzalas IG, Wüthrich D, Selimovic-Hamza S, Drögemüller C, Bruggmann R, Seuberlich T. 2016. Full-genome based molecular characterization of encephalitis-associated bovine astroviruses. *Infect Genet Evol* 44: 162-168. <https://doi.org/10.1016/j.meegid.2016.06.052>.
 46. Lipkin WI, Anthony SJ. 2015. Virus hunting. *Virology* 479-480:194-199. <https://doi.org/10.1016/j.virol.2015.02.006>.
 47. Schlottau K, Schulze C, Bilk S, Hanke D, Höper D, Beer M, Hoffmann B. 2016. Detection of a novel bovine astrovirus in a cow with encephalitis. *Transbound Emerg Infect* 63:253-259. <https://doi.org/10.1111/tbed.12493>.
 48. Selimovic-Hamza S, Sanchez S, Philibert H, Clark EG, Seuberlich T. 2017. Bovine astrovirus infection in feedlot cattle with neurological disease in western Canada. *Can Vet J* 58:601-603.
 49. Spinato MT, Vince A, Cai H, Ojkic D. 2017. Identification of bovine astrovirus in cases of bovine non-suppurative encephalitis in eastern Canada. *Can Vet J* 58:607-609.
 50. Deiss R, Selimovic-Hamza S, Seuberlich T, Meylan M. 2017. Neurologic clinical signs in cattle with astrovirus-associated encephalitis. *J*

- Vet Intern Med 31:1209-1214. <https://doi.org/10.1111/jvim.14728>.
51. Boujon CL, Koch MC, Wüthrich D, Werder S, Jakupovic D, Bruggmann R, Seuberlich T. 2017. Indication of cross-species transmission of astrovirus associated with encephalitis in sheep and cattle. *Emerg Infect Dis* 23: 1604-1606. <https://doi.org/10.3201/eid2309.170168>.
 52. Blomström AL, Ley C, Jacobson M. 2014. Astrovirus as a possible cause of congenital tremor type AII in piglets? *Acta Vet Scand* 56:82. <https://doi.org/10.1186/s13028-014-0082-y>.
 53. Arruda B, Arruda P, Hensch M, Chen Q, Zheng Y, Yang C, Gatto IRH, Ferreyra FM, Gauger P, Schwartz K, Bradner L, Harmon K, Hause B, Li G. 2017. Porcine astrovirus type 3 in central nervous system of swine with polioencephalomyelitis. *Emerg Infect Dis* 23:2097-2100. <https://doi.org/10.3201/eid2312.170703>.
 54. Vandeveld M, Higgins RJ, Oevermann A. 2012. *Veterinary neuropathology*, p 57. John Wiley & Sons, Chichester, United Kingdom.
 55. Janowski AB, Bauer IK, Holtz LR, Wang D. 2017. Propagation of astrovirus VA1, a neurotropic human astrovirus, in cell culture. *J Virol* 91:e00740-17. <https://doi.org/10.1128/JVI.00740-17>.
 56. Madeley CR, Cosgrove BP. 1975. Letter: 28 nm particles in faeces in infantile gastroenteritis. *Lancet* 2:451-452. [https://doi.org/10.1016/S0140-6736\(75\)90858-2](https://doi.org/10.1016/S0140-6736(75)90858-2).
 57. Méndez E, Murillo A, Velázquez R, Burnham A, Arias CF. 2013. Replication cycle of astroviruses, p 19-47. In Schultz-Cherry S (ed), *Astrovirus research: essential ideas, everyday impacts, future directions*. Springer Science + Business Media, New York, NY.
 58. Boujon CL, Selimovic-Hamza S, Bouzalas I, Seuberlich T. 2017. Development and validation of an immunohistochemistry procedure for the detection of a neurotropic bovine astrovirus. *J Virol Methods* 239:26-33. <https://doi.org/10.1016/j.jviromet.2016.10.013>.
 59. Bosch A, Méndez E, Krishna NK, Pantin-Jackwood M, Schultz-Cherry S, Monroe SS, Guix S. 2011. Astroviridae, p 953-960. In King AMQ, Adams MJ, Carstens EB, Lefkowitz EJ (ed), *Virus taxonomy*, 9th ed. Elsevier, Oxford, United Kingdom.
 60. Xiao CT, Gimenez-Lirola LG, Gerber PF, Jiang YH, Halbur PG, Opriessnig T. 2013. Identification and characterization of novel porcine astroviruses (PAstVs) with high prevalence and frequent co-infection of individual pigs with multiple PAstV types. *J Gen Virol* 94:570-582. <https://doi.org/10.1099/vir.0.048744-0>.
 61. Bosch A, Pintó RM, Guix S. 2014. Human astroviruses. *Clin Microbiol Rev* 27:1048-1074. <https://doi.org/10.1128/CMR.00013-14>.
 62. Monroe SS, Jiang B, Stine SE, Koopmans M, Glass RI. 1993. Subgenomic RNA sequence of human astrovirus supports classification of Astroviridae as a new family of RNA viruses. *J Virol* 67:3611-3614.
 63. Tengs T, Kristoffersen AB, Bachvaroff TR, Jonassen CM. 2013. A mobile genetic element with unknown function found in distantly related viruses. *Virol J* 10:132. <https://doi.org/10.1186/1743-422X-10-132>.
 64. Tengs T, Jonassen CM. 2016. Distribution and evolutionary history of the mobile genetic element s2m in coronaviruses. *Diseases* 4:27. <https://doi.org/10.3390/diseases4030027>.
 65. Bazan JF, Fletterick RJ. 1988. Viral cysteine proteases are homologous to the trypsin-like family of serine proteases: structural and functional implications. *Proc Natl Acad Sci USA* 85:7872-7876.
 66. Jiang B, Monroe SS, Koonin EV, Stine SE, Glass RI. 1993. RNA sequence of astrovirus: distinctive genomic organization and a putative retrovirus-like ribosomal frameshifting signal that directs the viral replicase synthesis. *Proc Natl Acad Sci USA* 90:10539-10543.
 67. Al-Mutairy B, Walter JE, Pothen A, Mitchell DK. 2005. Genome prediction of putative genome-linked viral protein (VPg) of astroviruses. *Virus Genes* 31:21-30. <https://doi.org/10.1007/s11262-004-2196-1>.
 68. Fuentes C, Bosch A, Pintó RM, Guix S. 2012. Identification of human astrovirus genome-linked protein (VPg) essential for virus infectivity. *J Virol* 86:10070-10078. <https://doi.org/10.1128/JVI.00797-12>.
 69. Nugent T, Jones DT. 2009. Transmembrane protein topology prediction using support vector machines. *BMC Bioinformatics* 10:159. <https://doi.org/10.1186/1471-2105-10-159>.
 70. Buchan DWA, Minneci F, Nugent TCO, Bryson K, Jones DT. 2013. Scalable web services for the PSIPRED Protein Analysis Workbench. *Nucleic Acids Res* 41:W349-W357. <https://doi.org/10.1093/nar/gkt381>.
 71. Lupas A, Van Dyke M, Stock J. 1991. Predicting coiled coils from protein sequences. *Science* 252:1162-1164. <https://doi.org/10.1126/science.252.5009.1162>.
 72. Combet C, Blanchet C, Geourjon C, Deleage G. 2000. NPS@: network protein sequence analysis. *Trends Biochem Sci* 25:147-150. [https://doi.org/10.1016/S0968-0004\(99\)01540-6](https://doi.org/10.1016/S0968-0004(99)01540-6).
 73. Guix S, Caballero S, Bosch A, Pintó RM. 2005. Human astrovirus C-terminal nsP1a protein is involved in RNA replication. *Virology* 333: 124-131. <https://doi.org/10.1016/j.virol.2004.12.023>.
 74. Guix S, Bosch A, Pintó RM. 2008. Astrovirus replication: an overview, p 571-595. In Cheng RH, Miyamura T (ed), *Structure-based study of viral replication*. World Scientific, Hackensack, NJ.
 75. Mason JM, Arndt KM. 2004. Coiled coil domains: stability, specificity, and biological implications. *ChemBiochem* 5:170-176. <https://doi.org/10.1002/cbic.200300781>.

76. Krishna NK. 2005. Identification of structural domains involved in astro-virus capsid biology. *Viral Immunol* 18:17-26. <https://doi.org/10.1089/vim.2005.18.17>.
77. Toh Y, Harper J, Dryden KA, Yeager M, Arias CF, Méndez E, Tao YJ. 2016. Crystal structure of the human astrovirus capsid protein. *J Virol* 90: 9008 -9017. <https://doi.org/10.1128/JVI.00694-16>.
78. York RL, Yousefi PA, Bogdanoff W, Haile S, Tripathi S, DuBois RM. 2015. Structural, mechanistic, and antigenic characterization of the human astrovirus capsid. *J Virol* 90:2254-2263. <https://doi.org/10.1128/JVI.02666-15>.
79. Bass DM, Qiu S. 2000. Proteolytic processing of the astrovirus capsid. *J Virol* 74:1810-1814. <https://doi.org/10.1128/JVI.74.4.1810-1814.2000>.
80. Méndez E, Fernández-Luna T, López S, Méndez-Toss M, Arias CF. 2002. Proteolytic processing of a serotype 8 human astrovirus ORF2 polyprotein. *J Virol* 76:7996-8002. <https://doi.org/10.1128/JVI.76.16.7996-8002.2002>.
81. Méndez E, Salas-Ocampo E, Arias CF. 2004. Caspases mediate processing of the capsid precursor and cell release of human astroviruses. *J Virol* 78:8601-8608. <https://doi.org/10.1128/JVI.78.16.8601-8608.2004>.
82. del Rocío Banos-Lara M, Méndez E. 2010. Role of individual caspases induced by astrovirus on the processing of its structural protein and its release from the cell through a non-lytic mechanism. *Virology* 401: 322-332. <https://doi.org/10.1016/j.virol.2010.02.028>.
83. Cordey S, Brito F, Vu DL, Turin L, Kilowoko M, Kyungu E, Genton B, Zdobnov EM, D'Acremont V, Kaiser L. 2016. Astrovirus VA1 identified by next-generation sequencing in a nasopharyngeal specimen of a febrile Tanzanian child with acute respiratory disease of unknown etiology. *Emerg Microbes Infect* 5:e67. <https://doi.org/10.1038/emi.2016.67>.
84. Pérot P, Lecuit M, Eliot M. 2017. Astrovirus diagnostics. *Viruses* 9:10. <https://doi.org/10.3390/v9010010>.
85. Appleton H, Higgins PG. 1975. Letter: viruses and gastroenteritis in infants. *Lancet* i:1297.
86. Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32:1792-1797. <https://doi.org/10.1093/nar/gkh340>.
87. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol Biol Evol* 30:2725-2729. <https://doi.org/10.1093/molbev/mst197>.
88. Bernhart SH, Hofacker IL, Will S, Gruber AR, Stadler PF. 2008. RNAalifold: improved consensus structure prediction for RNA alignments. *BMC Bioinformatics* 9:474. <https://doi.org/10.1186/1471-2105-9-474>.
89. Sánchez-Fauquier A, Carrascosa AL, Carrascosa JL, Otero A, Glass RI, Lopez JA, San Martin C, Melero JA. 1994. Characterization of a human astrovirus serotype 2 structural protein (VP26) that contains an epitope involved in virus neutralization. *Virology* 201:312-320. <https://doi.org/10.1006/viro.1994.1296>.
90. Olsen JV, Ong SE, Mann M. 2004. Trypsin cleaves exclusively C-terminal to arginine and lysine residues. *Mol Cell Proteomics* 3:608-614. <https://doi.org/10.1074/mcp.T400003-MCP200>



Gábor Reuter, M.D., Ph.D., D.Sc., is a virologist and specialist in medical microbiology. He studied medicine at the University of Pécs and holds a Ph.D. in virology from the University of Pécs Medical School in Pécs, Hungary. He worked for more than 15 years as a virologist of the State Public Health Service in Pécs, Hungary. He currently works as a full professor at the University of Pécs, in the Department of Medical Microbiology and Immunology. His research interests focus on clinical virology, molecular epidemiology, and pathogenesis of viral infections. His lab is interested in identification and characterization of novel viruses, investigation of the genetic diversity and evolution of RNA viruses, and description of viral coinfections and viromes of human and animal hosts. His scientific activity includes more than 130 publications in peer-reviewed international journals. He is a member of the International Committee on Taxonomy of Viruses, in the Picorna-viridae Study Group.



Ákos Boros, M.Sc., Ph.D., graduated from the Faculty of Natural Sciences, University of Pécs, with an M.Sc. in biology in 2007 and with a Ph.D. in neuroanatomy in 2011. He acquired a clinical microbiology qualification in 2016 from the Medical School, University of Pécs. He worked from 2009 as a molecular virologist and from 2016 as a clinical microbiologist in the Regional Laboratory of Virology, ÁNTSZ Regional Institute, State Public Health Service, and in the Department of Medical Microbiology and Immunology, Medical School, University of Pécs, in Pécs, Hungary. He is supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences and the National Excellence Program of the Ministry of Human Capacities. His fields of interest include virus discovery, virus evolution, and the pathogenesis of enteric single-stranded RNA viruses of humans and vertebrates. His scientific history includes 60 publications in peer-reviewed scientific journals.



Péter Pankovics, M.Sc., Ph.D., received his M.Sc. in biology-chemistry (2007) and his Ph.D. in medical sciences (2014) from the University of Pécs. His Ph.D. thesis examined the identification and genetic characterization of novel picornaviruses (Picornaviridae) by molecular biological methods. Since 2008, he has served as a molecular microbiologist in the Regional Laboratory of Virology, ÁNTSZ Regional Institute, State Public Health Service. His fields of interest include discovery of novel viruses and virus-like agents in animals and humans by molecular methods and viral metagenomics. His ongoing research focuses on novel picornaviruses (Picornaviridae), hepatitis viruses (Hepeviridae), picobirnaviruses (Picobirnaviridae), and astroviruses (Astroviridae). His research programs have been supported by grants from the Hungarian Scientific Research Fund (OTKA/NKFIH) and by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences. He has been actively involved in research since 2007, having published 57 publications in peer-reviewed international scientific journals.