

Chiral symmetry and bulk–boundary correspondence in periodically driven one-dimensional systems

J. K. Asbóth

*Institute for Solid State Physics and Optics, Wigner Research Centre,
Hungarian Academy of Sciences, H-1525 Budapest P.O. Box 49, Hungary*

B. Tarasinski

Instituut-Lorentz, Universiteit Leiden, P.O. Box 9506, 2300 RA Leiden, The Netherlands

P. Delplace

Laboratoire de Physique, Ecole Normale Supérieure de Lyon, 47 allée d'Italie, 69007 Lyon, France

(Dated: May 2014)

In periodically driven lattice systems, the effective (Floquet) Hamiltonian can be engineered to be topological: then, the principle of bulk–boundary correspondence guarantees the existence of robust edge states. However, such setups can also host edge states not predicted by the Floquet Hamiltonian. The exploration of such edge states, and the corresponding unique bulk topological invariants, has only recently begun. In this work we calculate these invariants for chiral symmetric periodically driven one-dimensional systems. We find simple closed expressions for these invariants, as winding numbers of blocks of the unitary operator corresponding to a part of the time evolution. This gives a robust way to tune these invariants using sublattice shifts. We illustrate our ideas on the periodically driven Su-Schrieffer-Heeger model, which, as we show, can realize a discrete time quantum walk: this opens a useful connection between periodically driven lattice systems and discrete time quantum walks. Our work helps interpret the results of recent simulations where a large number of Floquet Majorana fermions in periodically driven superconductors have been found.

Controlling the topological phases of matter is an important challenge in solid state physics. In the recent years, periodic driving has emerged as an important tool to meet this challenge. Topologically protected edge states, the hallmarks of topological phases, have been predicted and observed in periodically driven systems, such as materials irradiated by light^{1–4}, in shaken optical lattices^{5,6}, and in photonic crystals⁷. In the above cases, the principle of bulk–boundary correspondence⁸ was applied to the effective (Floquet) Hamiltonian of the periodically driven system.

The variety of topological phases that periodically driven systems can display, however, is much wider than those of their Floquet Hamiltonians, and the systematic exploration of these phases has only just begun⁹. An important example is the case of periodically driven one-dimensional topological superconductors, where, the bulk \mathbb{Z}_2 invariant is replaced by a pair of \mathbb{Z}_2 invariants, whose calculation necessitates information beyond that represented by the Floquet Hamiltonian¹⁰. The edge states then are the Floquet Majorana fermions, with potential applications in quantum information processing¹¹. Such states, not predicted by the bulk Floquet Hamiltonian, have also been observed in optical realization of a one-dimensional quantum walk¹².

Simulations of one-dimensional periodically driven superconductors have shown that they can host a large number of Floquet Majorana fermions at their ends^{13,14}. This can be explained by an extra chiral symmetry (CS) of the Floquet Hamiltonian, which prevents Majorana fermions on the same sublattice from recombining into complex fermions. Although this explanation is sufficient

in some cases^{13,14}, it cannot be general as it only relies on the Floquet Hamiltonian. Thus, the question is still open: what are the bulk topological invariants for periodically driven systems with CS?

In this paper, we find the bulk–boundary correspondence for periodically driven one-dimensional quantum systems with chiral symmetry, building on the theory of CS in discrete-time quantum walks^{15–18}. We show how CS can be ensured in a periodically driven system, whose time evolution in a period starts with a unitary operator F , by choosing an appropriate second part for the period. We show that the topological invariants predicting the number of 0 and π quasienergy end states are the winding numbers of the blocks of F in a canonical basis. Our formulas give a direct recipe to tune the topological invariants using a sublattice shift operation. We give an example of how to realize this operation in the simplest periodically driven one-dimensional Floquet insulator with CS, the periodically driven Su-Schrieffer-Heeger (PDSSH) model. We show how this model realizes a discrete-time quantum walk, and how this can be used to calculate the topological invariants of particle-hole symmetric quantum walks.

Floquet formalism. We consider periodically driven single-particle lattice Hamiltonians, $H(t + 1) = H(t)$. The long-time dynamics of $H(t)$, i.e., over many periods, is governed by the time-evolution operator of one period, the Floquet operator $U(\tau)$,

$$U(\tau) = \mathbb{T}e^{-i \int_{\tau}^{\tau+1} H(t) dt}, \quad (1)$$

where \mathbb{T} stands for time ordering. If at time τ the system is in an eigenstate $|\Psi\rangle$ of the Floquet operator, $U(\tau)|\Psi\rangle =$

$e^{-i\varepsilon}|\Psi\rangle$, then at all times $\tau + n$, for $n \in \mathbb{Z}$, it will be in state $e^{-in\varepsilon}|\Psi\rangle$. In this sense, the periodically driven system acts as a stroboscopic simulator of the effective (Floquet) Hamiltonian H_{eff} ,

$$H_{\text{eff}}(\tau) = i\ln U(\tau). \quad (2)$$

We fix the branch of the logarithm by restricting the eigenvalues ε of H_{eff} , the quasienergies, to $-\pi < \varepsilon \leq \pi$.

The Floquet operator $U(\tau)$, and thus also the effective Hamiltonian $H_{\text{eff}}(\tau)$, depend on the choice of the starting time of the period, τ . Changing τ amounts to a unitary transformation of the Floquet operator and the effective Hamiltonian (quasienergies are independent of τ).

Chiral symmetry of periodically driven systems. Ensuring CS of the periodically driven system amounts to ensuring that there is an initial time τ such that the corresponding effective Hamiltonian has CS, i.e., there is a unitary, Hermitian, and local (within a unit cell) operator Γ , that satisfies

$$\Gamma H_{\text{eff}}(\tau)\Gamma = -H_{\text{eff}}(\tau) \quad \Leftrightarrow \quad \Gamma U(\tau)\Gamma = U^{-1}(\tau). \quad (3)$$

The effective Hamiltonian does not inherit CS from the instantaneous Hamiltonian, as is the case with particle-hole symmetry¹⁹. However, CS of the periodically driven system is ensured if there is an intermediate time $0 < t_1 < 1$ that splits the period into a first and second part in a special way. Let F denote the time evolution of the first part of the cycle,

$$F = \mathbb{T}e^{-i \int_{\tau'}^{\tau'+t_1} H(t)dt}. \quad (4)$$

The second part of the cycle has to fulfil

$$\Gamma F^\dagger \Gamma = \mathbb{T}e^{-i \int_{\tau'+t_1}^{\tau'+1} H(t)dt}. \quad (5)$$

It is easy to check that in that case, not only $U' \equiv U(\tau')$, but also $U'' \equiv U(\tau'')$ have CS, where $\tau'' = \tau' + t_1$. These Floquet operators read

$$U' = \Gamma F^\dagger \Gamma F; \quad U'' = F \Gamma F^\dagger \Gamma. \quad (6)$$

Topological invariants of the effective Hamiltonians due to chiral symmetry. Consider a one-dimensional Floquet insulator: a long chain, with a translation invariant insulating bulk part, whose quasienergy spectrum has gaps around $\varepsilon = 0$ and π . If the system has CS, a local basis transformation can be performed that diagonalizes Γ , so that each lattice site has a sublattice index A or B, defined via the projectors $\Pi_{A/B} = (1 \pm \Gamma)/2$. We call such a basis a *canonical basis*. For the system to be a Floquet insulator, the number of A and B sites in each bulk unit cell has to be equal (or else the system would have flat bands at 0 or π quasienergy). We denote this number by N . In a canonical basis, the CS operator acts in each unit cell independently, as $\Gamma = \sigma_z \otimes 1_N$.

The spectrum of an effective Hamiltonian with CS is symmetric: stationary states $|\Psi'\rangle$ of H'_{eff} with

quasienergy $\varepsilon \neq 0, \pi$ have chiral symmetric partners $\Gamma|\Psi'\rangle$, that are also eigenstates with quasienergy $-\varepsilon$. Such states can be chosen to have equal support on both sublattices. The system can also host states $|\Psi'\rangle_{L/R}$ with quasienergy $\varepsilon = 0$ or π , whose wavefunctions are expelled from the bulk to the left/right by the gaps in the bulk spectrum. These *end states* can be chosen to have support only on one sublattice.

The effective Hamiltonians H'_{eff} and H''_{eff} have CS, as per Eqs. (6), and thus can be assigned topological invariants ν' and ν'' . These are obtained by standard procedure⁸, whereby we first isolate the bulk part of H'_{eff} and H''_{eff} , by imposing periodic boundary conditions on the translation invariant central part of these Hamiltonians, and taking the thermodynamic limit. The bulk Hamiltonians are periodic functions of the quasimomentum $k \in [-\pi, \pi)$, and, in the canonical basis, are block off-diagonal,

$$H_{\text{eff}}(k) = \begin{pmatrix} 0 & h(k) \\ h^\dagger(k) & 0 \end{pmatrix}. \quad (7)$$

Here, and later on, H_{eff} refers to either of H'_{eff} or H''_{eff} , and similarly for U and h . The topological invariants are

$$\nu' = \nu[h']; \quad \nu'' = \nu[h''], \quad (8)$$

where the function $\nu[h]$ is a winding number,

$$\nu[h] = \frac{1}{2\pi i} \int_{-\pi}^{\pi} dk \frac{d}{dk} \ln \det h(k). \quad (9)$$

These integers cannot change under adiabatic deformation of the bulk Hamiltonians, and so are equal to the winding numbers of the flat band limits of these Hamiltonians, which are the topological invariants of Ryu et al⁸. They can be interpreted as the dimensionless bulk sublattice polarization²⁰ of the effective Hamiltonians, at times τ' and τ'' .

Topological invariants of the driven system. To derive the topological invariants of the periodically driven system, we start by adopting the results obtained for discrete-time quantum walks (DTQW) with CS¹⁸ to periodically driven systems. The derivations follow very closely those of Ref. 18, and so we omit them here, but for completeness, we give details in Appendix A. As with DTQWs, also in periodically driven systems, the wavefunctions of quasienergy π end states switch sublattices as they evolve from time τ' to τ'' , and so, neither ν' , nor ν'' , on their own, give useful information about the number of end states (observations to the contrary in specific models^{13,14} do not generalize). The winding numbers ν' and ν'' must be combined to obtain the bulk topological invariants controlling the number of end states,

$$\nu_0 = \frac{\nu' + \nu''}{2}; \quad \nu_\pi = \frac{\nu' - \nu''}{2}. \quad (10)$$

We now proceed to simplify Eqs. (10), and express them using the blocks of F in the canonical basis:

$$F(k) = \begin{pmatrix} a(k) & b(k) \\ c(k) & d(k) \end{pmatrix}. \quad (11)$$

Along the way, we will use simple properties of the function $\nu[A(k)]$ of Eq. (9): $\nu[AB] = \nu[A] + \nu[B]$ and $\nu[A^\dagger] = -\nu[A]$, for arbitrary $A(k)$ and $B(k)$.

There are two constraints on the winding numbers of the blocks of the Floquet operator F representing the first part of the drive cycle, both following from the unitarity of F . First, substituting Eqs. (11) directly into $F(k)F(k)^\dagger = 1$ gives $ac^\dagger = -bd^\dagger$. Taking the winding numbers of the two sides gives

$$\nu[c] - \nu[a] = \nu[d] - \nu[b]. \quad (12)$$

Second, F represents an operation on an open chain, terminated at its ends. Thus, the average displacement of a state in the bulk, with this average going over all possible states, has to be zero: Otherwise, unitarity of F would be violated in the end regions. This average displacement is given by the winding number of F itself¹⁹, which, since F is unitary, can be written as

$$\nu[F] = \frac{1}{2\pi i} \int dk \text{Tr} F^\dagger(k) \frac{d}{dk} F(k). \quad (13)$$

Inserting the decomposition of F in the canonical basis, Eq. (11), into $\nu[F] = 0$, gives

$$\nu[F] = \nu[a] + \nu[c] + \nu[b] + \nu[d] = 0. \quad (14)$$

To use the relations derived above, we note, that

$$U = e^{-iH_{\text{eff}}} = \cos H_{\text{eff}} - i \sin H_{\text{eff}}. \quad (15)$$

Because of the block off-diagonal structure of H_{eff} , the first term in the sum above corresponds to the block diagonal and the second to the block off-diagonal parts of U . Now since $\text{sign}(\varepsilon) = \text{sign}(\sin \varepsilon)$ for $\varepsilon \in [-\pi, \pi]$, the winding number of H_{eff} is the same as that of $\sin H_{\text{eff}}$. Therefore, in Eq. (9) above, we can substitute the off-diagonal block of U in a canonical basis: $h \rightarrow iU_{12}$. For the topological invariants of the effective Hamiltonians H'_{eff} and H''_{eff} , using Eqs. (6), substituting the blocks of F , we obtain $\nu' = \nu[a^\dagger b - c^\dagger d]$ and $\nu'' = \nu[-ac^\dagger + bd^\dagger]$. We can simplify these using the unitarity of F , whereby $a^\dagger b + c^\dagger d = 0$ and $ac^\dagger + bd^\dagger = 0$, and the fact that $\nu[\alpha c] = \nu[c]$ for any $\alpha \in \mathbb{C}$. We obtain

$$\nu' = \nu[b] - \nu[a] = \nu[d] - \nu[c]; \quad (16a)$$

$$\nu'' = \nu[a] - \nu[c] = \nu[b] - \nu[d]. \quad (16b)$$

Inserting these equations into Eqs. (10), together with Eqs. (14) and (12), gives us

$$\nu_0 = \nu[b]; \quad \nu_\pi = \nu[d]. \quad (17)$$

These equations are the central result of our paper: In one-dimensional periodically driven systems with CS, the windings of the determinant of the off-diagonal and the diagonal blocks of the Floquet operator in a canonical basis fix the number of end states at quasienergy 0 and π , respectively.

Eqs. (17) determine the topological invariant ν_0 (ν_π) even if the gap of H_{eff} at quasienergy $\varepsilon = \pi$ ($\varepsilon = 0$) is closed, a problem raised by Tong et al.¹³. Consider

$$\cos H'_{\text{eff}} = 1 - 2 \begin{pmatrix} c^\dagger c & 0 \\ 0 & b^\dagger b \end{pmatrix} = 2 \begin{pmatrix} a^\dagger a & 0 \\ 0 & d^\dagger d \end{pmatrix} - 1. \quad (18)$$

If there is a quasimomentum k where the gap of H'_{eff} closes around $\varepsilon = 0$, then $\cos H'_{\text{eff}}(k)$ has a doubly degenerate eigenvalue +1. At that k , using the first relation of Eq. (18), either $c(k)$ or $b(k)$ (or both) have an eigenvalue zero. This means ν_0 is not well defined, and neither are ν' or ν'' . However, ν_π of Eq. (17) is still well defined. Similarly, if at some k the gap of H'_{eff} around $\varepsilon = \pi$ closes, then, using the second relation of Eq. (18), $a(k)$ or $d(k)$ must have an eigenvalue zero, and ν_π is not well defined, but ν_0 is.

Geometrical picture. In case of a two-band 1D Floquet insulator with CS, we can give a geometrical interpretation for the topological invariants ν_0 and ν_π . We relegate details to Appendix B, and just summarize the results here.

Disregarding an irrelevant global phase, the evolution operator for the first half of the period reads $F(k) = e^{-i\vec{f}(k)\vec{\sigma}}$, with $\vec{f}(k)$ a three-dimensional real vector inside a unit sphere of radius π , all points on whose surface are identified with each other, and $\vec{\sigma}$ the vector of Pauli matrices. As k traverses the Brillouin zone $[-\pi, \pi[$, $\vec{f}(k)$ describes a directed, smooth, closed loop. If the gap around $\varepsilon = 0$ is open, the loop of $\vec{f}(k)$ cannot touch the z -axis or the surface of the sphere, and we find that the invariant ν_0 is given by the winding of the loop around the z axis. If the gap of H_{eff} around $\varepsilon = \pi$ is open, the path of $\vec{f}(k)$ cannot touch the circle in the xy plane of radius $\pi/2$. In that case, ν_π is given by the winding of the loop around that circle.

Tuning the invariants. Formulas (17) allow for a simple way to tune the topological invariants of a periodically driven system, using a unitary sublattice shift operation $S(n)$, whose bulk part reads

$$S(n, k) = \exp(-in\Gamma k). \quad (19)$$

In the bulk, $S(n)$ displaces sites on sublattice A (B) to the right (left) by n sites. Therefore, at the left/right end, under the effect of $S(n)$, n states must switch sublattices, transitioning $B \rightarrow A$ / $A \rightarrow B$ (if n is negative, vice versa). How this transition happens depends on the details of $S(n)$ that have no influence on the topological invariants (nor on the number of end states).

To tune the invariants of a periodically driven system, obeying Eq. (6), with some $F = F^{(0)}$, insert extra sublattice shifts before and after $F^{(0)}$,

$$F^{(1)} = S(m)F^{(0)}S(n). \quad (20)$$

Substituting into Eqs. (17), we obtain directly the topological invariants of the modified driven system,

$$\nu_0^{(1)} = \nu_0^{(0)} + m - n; \quad \nu_\pi^{(1)} = \nu_\pi^{(0)} - m - n. \quad (21)$$




FIG. 1: Floquet eigenstates of a periodically driven SSH chain of 40 sites. (a) Time dependence of the intracell (continuous) and intercell (dotted) hopping amplitudes. (b) The curve $\vec{f}(k)$, which winds -1 times around the circle of radius $\pi/2$ on the xy plane, showing that $\nu_0 = -1$ and $\nu_\pi = -2$. (c) Local Density of States of the effective Hamiltonian $H_{\text{eff}}(0)$. (d) Time evolution of the position distribution $|\langle \Psi(t) | x \rangle|^2$ of the single end state with $\epsilon = 0$, and (e,f) of two orthogonal end states with $\epsilon = \pi$.

Example: the periodically driven SSH model. We now illustrate the concepts introduced above on the PDSSH model, given by

$$H_{\text{SSH}}(t) = \sum_{j=1}^M \left(v(t) c_{2j} c_{2j-1}^\dagger + w(t) c_{2j+1} c_{2j}^\dagger \right) + \text{h.c.}, \quad (22)$$

where c_x annihilates the fermion on site x . For simplicity, we keep the intracell hopping amplitudes $v(t)$ and the intercell hopping amplitudes $w(t)$ real, homogeneous in space, and modulated periodically, with period 1. We fix open boundary conditions by identifying $c_{2M+1} = 0$ (as opposed to periodic boundary conditions, which would require $c_{2M+1} = c_1$).

The sublattice shift operator $S(n)$ can be realized⁹ by the following drive sequence: a pulse of v of area $\pi/2$, followed by a pulse of w of area $-\pi/2$. This allows us to realize a discrete time quantum walk as a periodically driven lattice Hamiltonian.

As a concrete example, we consider the PDSSH model on an open chain of 40 sites ($M = 20$ unit cells). The drive sequence, shown in Fig. 1(a), consists of a train of nine pulses, chosen to be Gaussian for numerical convenience, applied to v and w homogeneously. We ensure CS by way of Eq. (6), with $t_1 = 0.5$, by choosing both $v(t)$ and $w(t)$ to be even functions of time.

We follow the recipe of Eq. (20), to realize $\nu_0 = -1, \nu_\pi = -2$. The role of $F^{(0)}$ is played by the first half of the central Gaussian pulse, where $w = 5v$: thus, it is a short pulse $e^{-i\pi/2H_1}$, where H_1 is an SSH Hamiltonian in the topologically nontrivial phase. So, we have $\nu_0^{(0)} = 1, \nu_\pi^{(0)} = 0$. To test the robustness of the recipe, we realize the sublattice displacement $S(n=2)$ only approximately by allowing considerable overlaps between the $\pi/2$ area pulses of v and the $-\pi/2$ area pulses of w .

We find that the bulk topological invariants and the end states agree perfectly with the theory above. The invariants are the winding numbers of the curve of Fig. 1 (b), which are $\nu_0 = -1, \nu_\pi = -2$. Correspondingly, in the local density of states, Fig. 1 (c), at each end, we find 2 end states at $\epsilon = \pi$, and 1 end state at $\epsilon = 0$, exclusively localized on B/A sublattice at the left/right end. The time dependence of these end states, Fig. 1 (d-f), shows that they indeed spread over both sublattices at intermediate times, but return to a single sublattice at $t = 0.5$. For the $0/\pi$ energy end states, this is the same/opposite sublattice as that occupied at $t = 0$.

Since we restricted the hopping amplitudes v and w to be real, the instantaneous SSH Hamiltonian, Eq. (22), has particle-hole symmetry (PHS), represented by ΓK , where K denotes complex conjugation. The PDSSH model inherits this symmetry, and therefore, its end states are analogous to 0 and π quasienergy Floquet Majorana fermions. If CS is violated, but PHS is maintained, only the parity of the number of the Floquet Majorana fermions at each edge and at each quasienergy $0, \pi$ is protected. There is a corresponding pair of bulk \mathbb{Z}_2 topological invariants¹⁰. In the case of the PDSSH model, we can follow the construction of Jiang et al.¹⁰, and find that the \mathbb{Z}_2 invariants can simply be obtained from the complete areas of the pulses of v and w . For details, see Appendix C.

Outlook. The topologically protected states our theory predicts should have experimental signatures in different kinds of setups. Optical experiments, where edge states are routinely imaged directly^{12,21}, are in the best position to test our predictions. Alternatively, in transport measurements, the end states should give rise to transmission resonances, similar to the ones predicted for Floquet Majorana fermions²².

Our work leaves a couple of theoretical questions open. First, is the decomposition of the drive cycle U into F and $\Gamma F^\dagger \Gamma$, as per Eqs. (4-6), a necessary requirement for a periodically driven Hamiltonian to have CS? For previously studied cases^{13,14} we can find such a decomposition, but if a counterexample were to be found, the theory we presented here would need to be expanded. Second, the bulk effective Hamiltonian $H_{\text{eff}}(\tau, k)$ of a one-dimensional Floquet insulator (with or without CS) is periodic in both τ and k , and thus has a Chern number. In all the examples we examined numerically, we found this Chern number to be zero, but can it take on a nonzero value? If so, what is the physical interpretation of this number?

Last, how can the topological invariants we found here be formulated in the frequency domain⁹? This is especially an interesting question, as previous work on the PDSSH model using this approach⁷ has not detected the pair of topological invariants we found.

We thank J. Dahlhaus, J. Li, A. Gábris and J. Edge, for useful discussions. PD acknowledges useful comments from A. Bernevig. This research was realized in the frames of TAMOP 4.2.4. A/1-11-1-2012-0001 "National Excellence Program – Elaborating and operating an in-

land student and researcher personal support system", subsidized by the European Union and co-financed by the European Social Fund. This work was also supported by the Hungarian National Office for Research and Technology under the contract ERC_HU_09 OPTOMECH and the Hungarian Academy of Sciences (Lendület Program, LP2011-016). This research was supported by the Foundation for Fundamental Research on Matter (FOM) and the Netherlands Organization for Scientific Research (NWO/OCW).

-
- ¹ L. N. H., G. Refael, and V. Galitski, *Nature Physics* **7**, 490495 (2011).
- ² B. Dóra, J. Cayssol, F. Simon, and R. Moessner, *Phys. Rev. Lett.* **108**, 056602 (2012), URL <http://link.aps.org/doi/10.1103/PhysRevLett.108.056602>.
- ³ T. Kitagawa, T. Oka, A. Brataas, L. Fu, and E. Demler, *Phys. Rev. B* **84**, 235108 (2011), URL <http://link.aps.org/doi/10.1103/PhysRevB.84.235108>.
- ⁴ Y. H. Wang, H. Steinberg, P. Jarillo-Herrero, and N. Gedik, *Science* **342**, 453 (2013), <http://www.sciencemag.org/content/342/6157/453.full.pdf>, URL <http://www.sciencemag.org/content/342/6157/453.abstract>.
- ⁵ P. Hauke, O. Tieleman, A. Celi, C. Ölschläger, J. Simonet, J. Struck, M. Weinberg, P. Windpassinger, K. Senegstock, M. Lewenstein, et al., *Phys. Rev. Lett.* **109**, 145301 (2012), URL <http://link.aps.org/doi/10.1103/PhysRevLett.109.145301>.
- ⁶ M. Reichl and E. Mueller, arXiv preprint arXiv:1404.3217 (2014).
- ⁷ M. C. Rechtsman, J. M. Zeuner, Y. Plotnik, Y. Lumer, D. Podolsky, F. Dreisow, S. Nolte, M. Segev, and A. Szameit, *Nature* **496**, 196 (2013), ISSN 0028-0836, URL <http://dx.doi.org/10.1038/nature12066>.
- ⁸ S. Ryu, A. P. Schnyder, A. Furusaki, and A. W. W. Ludwig, *New Journal of Physics* **12**, 065010 (2010).
- ⁹ M. S. Rudner, N. H. Lindner, E. Berg, and M. Levin, *Phys. Rev. X* **3**, 031005 (2013), URL <http://link.aps.org/doi/10.1103/PhysRevX.3.031005>.
- ¹⁰ L. Jiang, T. Kitagawa, J. Alicea, A. R. Akhmerov, D. Pekker, G. Refael, J. I. Cirac, E. Demler, M. D. Lukin, and P. Zoller, *Phys. Rev. Lett.* **106**, 220402 (2011), URL <http://link.aps.org/doi/10.1103/PhysRevLett.106.220402>.
- ¹¹ D. E. Liu, A. Levchenko, and H. U. Baranger, *Phys. Rev. Lett.* **111**, 047002 (2013), URL <http://link.aps.org/doi/10.1103/PhysRevLett.111.047002>.
- ¹² T. Kitagawa, M. A. Broome, A. Fedrizzi, M. S. Rudner, E. Berg, I. Kassal, A. Aspuru-Guzik, E. Demler, and A. G. White, *Nature Communications* **3**, 882 (2012).
- ¹³ Q.-J. Tong, J.-H. An, J. Gong, H.-G. Luo, and C. H. Oh, *Phys. Rev. B* **87**, 201109 (2013), URL <http://link.aps.org/doi/10.1103/PhysRevB.87.201109>.
- ¹⁴ M. Thakurathi, A. A. Patel, D. Sen, and A. Dutta, *Phys. Rev. B* **88**, 155133 (2013), URL <http://link.aps.org/doi/10.1103/PhysRevB.88.155133>.
- ¹⁵ T. Kitagawa, M. S. Rudner, E. Berg, and E. Demler, *Phys. Rev. A* **82**, 033429 (2010), URL <http://link.aps.org/doi/10.1103/PhysRevA.82.033429>.
- ¹⁶ T. Kitagawa, *Quantum Information Processing* **11**, 1107 (2012).
- ¹⁷ J. K. Asbóth, *Phys. Rev. B* **86**, 195414 (2012), URL <http://link.aps.org/doi/10.1103/PhysRevB.86.195414>.
- ¹⁸ J. K. Asbóth and H. Obuse, *Phys. Rev. B* **88**, 121406 (2013), URL <http://link.aps.org/doi/10.1103/PhysRevB.88.121406>.
- ¹⁹ T. Kitagawa, E. Berg, M. Rudner, and E. Demler, *Phys. Rev. B* **82**, 235114 (2010), URL <http://link.aps.org/doi/10.1103/PhysRevB.82.235114>.
- ²⁰ I. Mondragon-Shem, J. Song, T. L. Hughes, and E. Prodan, arXiv preprint arXiv:1311.5233 (2013).
- ²¹ Y. E. Kraus, Y. Lahini, Z. Ringel, M. Verbin, and O. Zeitlinger, *Phys. Rev. Lett.* **109**, 106402 (2012), URL <http://link.aps.org/doi/10.1103/PhysRevLett.109.106402>.
- ²² A. Kundu and B. Seradjeh, *Phys. Rev. Lett.* **111**, 136402 (2013).
- ²³ B. Tarasinski, J. K. Asbóth, and J. P. Dahlhaus, *Phys. Rev. A* **89**, 042327 (2014), URL <http://link.aps.org/doi/10.1103/PhysRevA.89.042327>.

Appendix A: Derivation of Eqs. (10)

To derive Eqs. (10), we follow closely the line of thought of Ref. 18. We consider an open, periodically driven chain with CS, which has one bulk and two ends. Let $n'_{A/B,0/\pi}$ denote the number of end states at the left end on the A/B sublattice at quasienergy $0/\pi$ of the Hamiltonian H'_{eff} , and $n''_{A/B,0/\pi}$ the corresponding quantities for H''_{eff} . The bulk–boundary correspondence for the effective Hamiltonians H'_{eff} and H''_{eff} reads

$$\nu' = n'_{A,0} - n'_{B,0} + n'_{A,\pi} - n'_{B,\pi}; \quad (\text{A1a})$$

$$\nu'' = n''_{A,0} - n''_{B,0} + n''_{A,\pi} - n''_{B,\pi}. \quad (\text{A1b})$$

Topologically protected end states of periodically driven one-dimensional lattices with CS can be divided to two classes: a), they have quasienergy 0 and are on the same sublattice at τ' and τ'' , or b) have quasienergy π and are on opposite sublattices. Indeed, consider a topologically protected end state $|\Psi'\rangle$, which is an eigenstate of U' with eigenvalue $e^{-i\varepsilon}$, with $\varepsilon \in \{0, \pi\}$. It is only on a single sublattice: $\Gamma|\Psi'\rangle = e^{-i\gamma}|\Psi'\rangle$, with $\gamma = 0/\pi$ corresponding to sublattice A/B . Now consider the same end

state at the other special time τ'' , $|\Psi''\rangle = F|\Psi'\rangle$. This is an eigenstate of U'' with the same quasienergy ε . This state is also on one sublattice only, because $\Gamma F|\Psi'\rangle = \Gamma F \Gamma e^{i\gamma} |\Psi'\rangle = \Gamma F \Gamma e^{i(\gamma-\varepsilon)} \Gamma F^{-1} \Gamma F \Psi' = e^{i(\gamma-\varepsilon)} F|\Psi'\rangle$. So $|\Psi''\rangle$ is on the same (opposite) sublattice as $|\Psi'\rangle$ if $\varepsilon = 0$ ($\varepsilon = \pi$). This can be written succinctly as

$$n''_{A,\pi} - n'_{B,\pi} = n''_{B,\pi} - n'_{A,\pi} = 0; \quad (\text{A2a})$$

$$n''_{A,0} - n'_{A,0} = n''_{B,0} - n'_{B,0} = 0. \quad (\text{A2b})$$

Using Eqs. (A2) to simplify $\nu' + \nu''$ and $\nu' - \nu''$ from Eqs. (A1), we obtain

$$\nu_0 = \frac{\nu' + \nu''}{2}; \quad \nu_\pi = \frac{\nu' - \nu''}{2}, \quad (\text{A3})$$

which are Eqs. (10) we set out to demonstrate.

Appendix B: Geometrical picture

For a two-band 1D Floquet insulator with CS, we can give a direct geometrical picture for the topological invariants ν_0 and ν_π . Since the global phase cannot wind (F cannot have quasienergy winding), it can safely be disregarded, and the evolution operator for the first half of the period then reads $F(k) = e^{-i\vec{f}(k)\vec{\sigma}}$. Here \vec{f} is a 3-dimensional vector, of magnitude $f \in [0, \pi]$ and $\vec{\sigma}$ the vector of Pauli matrices. The k -dependent vector $\vec{f}(k)$ is restricted inside a spherical ball of radius π , with all points on the surface identified with each other. The a, b, c, d in Eq. (11) are just complex number valued functions of k ,

$$F = \begin{pmatrix} \cos f - i \sin f \cos \theta & -i \sin f \sin \theta e^{-i\phi} \\ -i \sin f \sin \theta e^{i\phi} & \cos f + i \sin f \cos \theta \end{pmatrix}, \quad (\text{B1})$$

using spherical coordinates. As k traverses the Brillouin zone, $\vec{f}(k)$ describes a directed, smooth, closed loop, that can at some k exit the ball at a point on the surface and reenter at the same k at the antipodal point.

If the gap around $\varepsilon = 0$ is open, the loop of $\vec{f}(k)$ cannot touch the z -axis, nor the surface of the sphere. Thus, the loop has a well defined winding number around the z axis,

$$\nu_0 = \frac{1}{2\pi} \int dk \frac{d}{dk} \phi(k). \quad (\text{B2})$$

Since both $f(k), \theta(k) \in]0, \pi[$ for all k , this is the same as the winding number ν_0 obtained by substituting (B1) into Eq. (17).

The gap of H_{eff} around $\varepsilon = \pi$ closes when $\vec{f}(k)$ is on the circle on the $n_z = 0$ plane of radius $\pi/2$ ($n_z = 0$ and $f = \pi/2$). Thus, if the gap around $\varepsilon = \pi$ is open, the loop of $\vec{f}(k)$ has a well defined winding number around that circle. To calculate this winding number, first discard the ϕ information, by setting $\phi = 0$. This transforms the

3D closed path of $\vec{f}(k)$ into a 2D path in a semicircle, with the points on the circular boundary with the same x coordinate identified. We need the winding of this path around the single point, $f = \pi/2, n_z = 0$. This is found by deforming the semicircle yet again, by the transformation $(f \sin \theta, f \cos \theta) \rightarrow (\cos f, \sin f \cos \theta)$, into circle, into whose origin the point $f = \pi/2, n_z = 0$ is mapped. The winding number is then

$$\nu_\pi = \frac{1}{2\pi} \int dk \frac{d}{dk} \arctan \frac{\cos f(k)}{\sin f(k) \cos \theta(k)} \quad (\text{B3})$$

which is the same as ν_π obtained by substituting Eq. (B1) into Eq. (17b).

Appendix C: The $\mathbb{Z}_2 \times \mathbb{Z}_2$ invariant

The PDSSH model, Eq. (22), has particle-hole symmetry (PHS), represented by ΓK , where K stands for complex conjugation. This antiunitary symmetry is inherited by the effective Hamiltonian from the instantaneous Hamiltonian¹⁹.

If we break CS in the PDSSH model, an end state can remain protected if it can have no PHS partner. This happens whenever the number of end states at a given energy and at a given end is odd: then, after breaking CS, a single end state is still protected by PHS. We illustrate this on the PDSSH model. If we break CS by delaying the intracell hopping amplitude v by δt with respect to the intercell hopping w pulses, as shown in Fig. 2(a), the lone end state at $\varepsilon = 0$ is still topologically protected, while the pair of end states at $\varepsilon = \pi$ hybridize and move away from the edge of the energy Brillouin zone (except for a time shift of 0.5, where the conditions for CS are again fulfilled). To break PHS, we can add a sublattice potential to the SSH model, obtaining the periodically driven Rice-Mele (PDRM) model,

$$H_{\text{RM}}(t) = H_{\text{SSH}}(t) + u(t) \sum_{x=1}^M \left(c_{2x-1}^\dagger c_{2x-1} - c_{2x}^\dagger c_{2x} \right). \quad (\text{C1})$$

Now, CS still holds if in addition to $v(t)$ and $w(t)$ being even functions of time, $u(t)$ is odd: $u(t) = -u(-t)$. We choose $u(t) = \sin(2\pi t)$. This time, if we break CS by shifting the $v(t)$ pulse in time with respect to the $w(t)$ and $u(t)$ pulses, as shown in Fig. 2(b), all end states move away from their original energies (again except for the time shift of 0.5).

The extra PHS of the PDSSH model brings with it an extra pair of bulk topological invariants, $(Q_0, Q_\pi) \in \mathbb{Z}_2 \times \mathbb{Z}_2$, which predict the number of end states protected by PHS at 0 and π energy. If we have CS, the invariants are just $Q_\varepsilon = \nu_\varepsilon \bmod 2$; if CS is broken, however, they can only be obtained by a procedure involving analytic continuation based on the full cycle $H(t)$, as found by Jiang et al.¹⁰.




FIG. 2: Effect of breaking CS by time-shifting the pulse of the intracell hopping $v(t)$ with respect to the other pulses. (a) In the PDSSH model, the extra PHS protects the end states at $\varepsilon = 0$. (b) In the PDRM model, there is no PHS, and all end state energies are affected by the time shift.

We find that for the PDSSH model, the invariant of Jiang et al.¹⁰ can be given by simple closed formulas. At the momenta $k = 0$ and $k = \pi$, the Hamiltonians at different times all commute with each other, and therefore, all that matters is the total area under the v and w pulses,

$$V = \int_0^1 v(t) dt; \quad W = \int_0^1 w(t) dt. \quad (\text{C2})$$

A short calculation gives

$$Q_0 = \text{sgn} \left(\sin \frac{V+W}{2} \sin \frac{V-W}{2} \right); \quad (\text{C3})$$

$$Q_0 Q_\pi = \text{sgn}(\sin(V+W) \sin(V-W)). \quad (\text{C4})$$

Appendix D: Mapping to the discrete time quantum walk

The PDSSH model, besides being the simplest periodically driven topological insulator, also gives a lattice realization of the discrete time split-step quantum walk. For the quantum walk, we need to define the basis states $|R/L, x\rangle$, for coin state predicting the next step right/left, and the walker at position x . These basis states are identified with states on the SSH chain as

$$c_{2x+1}^\dagger |0\rangle = |R, x\rangle; \quad (\text{D1})$$

$$c_{2x}^\dagger |0\rangle = -i |L, x\rangle. \quad (\text{D2})$$

The basic operations of the split-step walk are rotations of the internal state of the walker, $R(\theta) = e^{-i\theta\sigma_y}$, and shifts of the R/L internal state to the right/left, given by $S_\pm = e^{-ik(\sigma_z \pm 1)}$. One timestep of the split-step walk is defined as

$$U = S_- e^{-i\theta_2 \sigma_y} S_+ e^{-i\theta_1 \sigma_y}. \quad (\text{D3})$$

A pulse of v of area V followed by a pulse of w of area W , in the basis of Eq. D2, can be written as

$$U = e^{-iW(\cos k\sigma_y - \sin k\sigma_x)} e^{-iV\sigma_y}, \quad (\text{D4})$$

which reproduces the timestep of the split-step walk with the angles

$$\theta_2 = W + \pi/2; \quad \theta_1 = V - \pi/2. \quad (\text{D5})$$

The above mapping is important as it allows us to apply results about the topological phases of periodically driven systems to quantum walks.

As an example, consider the invariants due to CS, via Eqs. (17), for the simple quantum walk, given by $U = S_- S_+ e^{-i\theta\sigma_y}$. According to the mapping above, the winding numbers are $\nu_0 = \nu[-i(s + ce^{ik})]$, $\nu_\pi = \nu[c - se^{-ik}]$, with $c = \cos(\pi/4 + \theta/2)$, $s = \sin(\pi/4 + \theta/2)$. We get $(\nu_0, \nu_\pi) = (+1, 0)$ if $|c| > |s|$, i.e., if $\theta \in [-\pi, 0]$, and $(0, -1)$ if $\theta \in [0, \pi]$. This is shifted by $(1/2, -1/2)$ from the invariants obtained by the scattering matrix method²³, but such a shift is not physical: both methods predict a pair of end states at 0 and π quasienergy at an interface between bulks with $\theta < 0$ and $\theta > 0$, as seen in simulations¹⁷.

Another example is the calculation of the invariants due to PHS in the split-step quantum walk. Compared to the invariants $Q_0^{(\text{gap})}, Q_\pi^{(\text{gap})}$, defined via gap closings in the parameter space¹⁷, the above mapping to the PDSSH model, together with Eqs. (C4) gives $Q_0 = Q_0^{(\text{gap})}$, and $Q_\pi = 1 - Q_\pi^{(\text{gap})}$, which agrees in all the predictions concerning end states at interfaces. Compared to the scattering matrix topological invariants²³, we of course find the same constant shift by $(1/2, -1/2)$ as for the invariants due to CS, which has no influence on the physical predictions.