

On the weighted trigonometric Bojanov-Chebyshev extremal problem

Béla Nagy and Szilárd Gy. Révész

September 13, 2023

Dedicated to Vitalii Vladimirovich Arestov, a leading mathematician and teacher of generations, on the occasion of his 80th anniversary

Abstract

We investigate the weighted Bojanov-Chebyshev extremal problem for trigonometric polynomials, that is, the minimax problem of minimizing $\|T\|_{w, C(\mathbb{T})}$, where w is a sufficiently nonvanishing, upper bounded, nonnegative weight function, the norm is the corresponding weighted maximum norm on the torus \mathbb{T} , and T is a trigonometric polynomial with prescribed multiplicities ν_1, \dots, ν_n of root factors $|\sin(\pi(t - z_j))|^{\nu_j}$. If the ν_j are natural numbers and their sum is even, then T is indeed a trigonometric polynomial and the case when all the ν_j are 1 covers the Chebyshev extremal problem.

Our result will be more general, allowing, in particular, so-called generalized trigonometric polynomials. To reach our goal, we invoke Fenton's sum of translates method. However, altering from the earlier described cases without weight or on the interval, here we find different situations, and can state less about the solutions.

Keywords: minimax and maximin problems, kernel function, sum of translates function, vector of local maxima, equioscillation, majorization

2020 Mathematics subject classification: 26A51, 26D07, 49K35

1 Introduction

In this paper our aim is to solve the Bojanov-Chebyshev extremal problem in the setting of weighted trigonometric polynomials. As in our earlier papers on related subjects, our approach is the so-called “sum of translates method” of Fenton, what he introduced in [10]. However, here we do not develop the whole theory for two reasons: first, in the periodic, i.e., torus setup, in [5] we have already developed much of what is possible, and second, much of what we could find useful here, is simply not holding true. In this regard, our Example 20 is an important part of the study, showing the limits of any proof in this generality.

The analogous problems for the unweighted periodic case and the weighted and unweighted algebraic polynomial cases on the interval were already solved in [5] and in [7] and [9]. The results available to date do not imply, not in a direct and easy way, the corresponding result to the weighted trigonometric polynomial Bojanov-Chebyshev problem. In fact, some of them simply does not remain valid. So, the weighted trigonometric polynomial case poses new challenges and requires a careful adaptation of our methods, with an avoidance of certain obstacles – for example a bagatelle-looking, but in fact serious dimensionality obstacle in the way of proving a homeomorphism theorem, analogous to the earlier cases – and recombining our existing knowledge about the torus setup with all what can be saved and reused from the interval case. So, we heavily rely on all our earlier papers [5] [8] [7] [6] [9] on the subject, while these in themselves will not suffice to reach our goals. We will still need to devise new proofs or at least new versions for various existing arguments.

We note that proving minimax- and equioscillation type results in certain contexts may be attempted without rebuilding the whole theory, just by transferring some existing results of an already better explored case to the new settings. This has already been done in [5] for the (unweighted) algebraic polynomial case of the interval, deriving it from the (unweighted) trigonometric polynomial case, explored in the major part of [5]. However, the transference was not easy and broke down for general weights (even if for even weights it seemed working). Similarly, in [14] Tatiana Nikiforova succeeded in transferring certain results to the real line and semiaxis cases from the interval case – while leaving unresolved some of the related and still interesting questions. However, in both cases we can expect a more detailed and complete picture when we take the time to build up the method and explore the full strength of it right in the given context. Therefore, we did not settle with the results which could be transferred from [5], but worked out the interval case fully in [7], [8], [6]; we also think that it would be worthwhile to do so in the cases of the real line and the semiaxis. In particular, the relevant variant of the homeomorphism theorem is missed very much for the real line and the semiaxis. However, as already said and as will be explained in due course later, in the current setup that buildup does not seem to be possible, and we must be satisfied by a combination of transferred results and ad hoc arguments.

From our point of view, however, the weighted trigonometric polynomial Bojanov-Chebyshev problem is not a main goal, but more of an application, which testifies the strength of the method. We try to work in a rather general framework, and prove more general results than that. In particular, the results will be valid also for generalized trigonometric polynomials (GTPs), which are introduced, e.g., in [3] Chapter A4 as follows.

$$\mathcal{T} := \bigcup_{n=1}^{\infty} \mathcal{T}_n, \quad \mathcal{T}_n := \left\{ T(\mathbf{z}, t) := c_0 \prod_{j=1}^n |\sin \pi(t - z_j)|^{\nu_j} : c_0 > 0, \right. \quad (1)$$

$$\left. \nu_j > 0 (j = 1, \dots, n), \mathbf{z} = (z_1, \dots, z_n) \in \mathbb{C}^n \right\}.$$

By periodicity one can assume $0 \leq \Re z_j < 1$, and in the below extremal problems it is obvious that replacing $\Re z_j$ for z_j can only decrease the quantity to be minimized, so that we will assume that all the z_j 's are real. However, fixing the ordering of z_j (or $\Re z_j$) has a role, with different fixed orderings posing separate extremal problems, and the ordering-specific solution being much stronger, than just a “global” minimization. As this issue has already been discussed in [5] and [7], e.g., we leave the details to the reader simply addressing the order-specific, stronger question here. One particular result ahead of us will be the following.

Theorem 1. *Let $n \in \mathbb{N}$ and $\nu_1, \dots, \nu_n > 0$ be given. Put $\nu := (\nu_1, \dots, \nu_n)$.*

Further, let $w : \mathbb{R} \rightarrow [0, \infty)$ be an upper bounded, nonnegative, 1-periodic weight function, attaining positive values at more than n points of $\mathbb{T} := \mathbb{R}/\mathbb{Z}$.

Denote the weighted sup norm by $\|\cdot\|_w$ and consider the minimax problem

$$M := M(w, \nu) := \inf \left\{ \|T(\mathbf{z}, \cdot)\|_w : T(\mathbf{z}; t) = \prod_{j=1}^n |\sin \pi(t - z_j)|^{\nu_j} \in \mathcal{T}_n, \right. \\ \left. \exists c \in [0, 1) \quad \text{such that} \quad c \leq \Re z_1 \leq \dots \leq \Re z_n \leq c + 1 \right\}.$$

Then there exists a minimax point $\mathbf{z}^ = (z_1^*, \dots, z_n^*) \in \mathbb{T}^n$ and with the prescribed cyclic ordering of the nodes, satisfying $\|T(\mathbf{z}^*, \cdot)\|_w = M(w, \nu)$.*

Moreover, all z_j^ s are distinct and real, and their cyclic ordering is strict in the sense that there exists $c \in \mathbb{R}$ such that $c < z_1^* < \dots < z_n^* < c + 1$ (as in the prescribed order, but with strict inequalities).*

Furthermore, this extremal point has the equioscillation property, that is, $\max\{T(\mathbf{z}^, t) : z_1^* \leq t \leq z_2^*\} = \dots = \max\{T(\mathbf{z}^*, t) : z_j^* \leq t \leq z_{j+1}^*\} = \dots = \max\{T(\mathbf{z}^*, t) : z_n^* \leq t \leq z_1^* + 1\} = M(w, \nu)$.*

The occurrence of c in the description is another simple-looking, yet important difference between the interval and torus setup. Basically, we fix here the ordering of nodes z_j^* only cyclically, that is, as they follow each other when one covers the circle once, moving continuously from some appropriate $c \in \mathbb{T}$ in the positive (counter-clockwise) orientation until return.

2 Basics for the Bojanov-Chebyshev problem

2.1 Trigonometric polynomials and generalized trigonometric polynomials

It is well known that (real) trigonometric polynomials can be factorized as follows. Let

$$T(t) := a_0 + \sum_{j=1}^n a_j \sin(2\pi j t) + b_j \cos(2\pi j t) \quad (2)$$

be a (real) trigonometric polynomial ($a_0, a_1, b_1, a_2, b_2, \dots, a_n, b_n \in \mathbb{R}$, $a_n^2 + b_n^2 \neq 0$) of degree n (with period 1). Then (see, e.g., [3] p. 10) there exist uniquely

$c_0 \in \mathbb{R}$, $c_0 \neq 0$, $z_1, \dots, z_{2n} \in \mathbf{C}$ such that nonreal z_j 's occur in conjugate pairs and

$$T(t) = c_0 \prod_{j=1}^{2n} \sin(\pi(t - z_j)). \quad (3)$$

This explains that GTPs are indeed generalizations of trigonometric polynomials¹.

By taking logarithm of a generalized trigonometric polynomial (1), we have

$$\log T(t) = \log c_0 + \sum_{j=1}^n \nu_j \log |\sin(\pi(t - z_j))|.$$

In this work we assume for normalization that our trigonometric polynomial or GTP is *monic*, i.e., the “leading coefficient” is $c_0 = 1$. However, we consider weights, which are fixed, but can as well be constants, so that the weighted norm can incorporate any other prescribed leading coefficient as well. Obviously, the weighted minimax problem is equivalent to minimizing $\log \|T(\mathbf{z}, \cdot)\|_w = \sup_{\mathbb{T}} \log w(t) + \sum_{j=1}^n \nu_j \log |\sin(\pi(t - z_j))|$. This reformulation leads to considering sums (and positive linear combinations) of translated copies of the basic “kernel function” $\log |\sin(\pi t)|$, instead of products of root factors. That reformulation, so standard in logarithmic potential theory, will be the starting point of our presentation of the Fenton method for the current setup.

2.2 Basics of Fenton’s sum of translates approach

In this paragraph we present the by now standard notations and terminology how we use Fenton’s method. There is particular need for this clarification because we will use it in *two* setups, needing it for the torus \mathbb{T} , but also time to time referring to and invoking into our arguments corresponding results for the interval case. Here we start with notations and terminologies which can be equally interpreted for the torus and real line case, so with a slight abuse of notation we do not distinguish between them.

However, in the next subsection we set a separated terminology with quantities for the periodic case denoted by a star, because in these notions there are some essential alterations. With this long, and sometimes doubled list of definitions, notions and terminology, these paragraphs will be boring and longish, but in later sections we will need it for precise references. Note that in most of our definitions we will *not* assume that the considered functions and setups were periodic, and handle the periodic cases only as special cases, especially pointing out the periodicity assumption.

¹The somewhat curious fact is that root factorization of trigonometric polynomials relies on *pairs of factors* of the form $\sin(\pi(t - z_j))$, where one such factor in itself is not a trigonometric polynomial (as it is only antiperiodic, but not periodic by 1). Considering general products of root factors thus leads to GTPs.

A function $K : (-1, 0) \cup (0, 1) \rightarrow \mathbb{R}$ is called a *kernel function* if it is concave on $(-1, 0)$ and on $(0, 1)$, and if it satisfies

$$\lim_{t \downarrow 0} K(t) = \lim_{t \uparrow 0} K(t). \quad (4)$$

By the concavity assumption these limits exist, and a kernel function has one-sided limits also at -1 and 1 . We set

$$K(0) := \lim_{t \rightarrow 0} K(t), \quad K(-1) := \lim_{t \downarrow -1} K(t) \quad \text{and} \quad K(1) := \lim_{t \uparrow 1} K(t).$$

We note explicitly that we thus obtain the extended continuous function $K : [-1, 1] \rightarrow \mathbb{R} \cup \{-\infty\} =: \underline{\mathbb{R}}$, and that we have $\sup K < \infty$. Also note that a kernel function is almost everywhere differentiable.

A kernel function K is called *singular* if

$$K(0) = -\infty. \quad (\infty)$$

We say that the kernel function K is *strictly concave* if it is strictly concave on both of the intervals $(-1, 0)$ and $(0, 1)$.

In this paper we consider only systems of kernels which are constant multiples of each other, i.e.,

$$K_j(t) = \nu_j K(t) \quad (5)$$

for some $\nu_1, \dots, \nu_n > 0$ and some kernel function $K(t)$.

The condition

$$K'(t) - K'(t-1) \geq c \quad \text{for a.e. } t \in [0, 1], \quad (PM_c)$$

was called “periodized c -monotonicity” in [8] and [7]. The particular case $c = 0$ deserves special attention. Then we have

$$K'(t) - K'(t-1) \geq 0 \quad \text{for a.e. } t \in [0, 1]. \quad (PM_0)$$

Our main objective is the study of kernels which extend to \mathbb{R} 1-periodically:

$$K(t-1) = K(t), \quad t \in \mathbb{R}, \quad (6)$$

but sometimes we will invoke more general, not necessarily periodic kernels, too. It is straightforward that (6) implies (PM_0) .

Note that the log-trigonometric kernel

$$K(t) := \log |\sin(\pi t)| \quad (t \in \mathbb{R}), \quad (7)$$

which is in the focus of our analysis, is periodic (6), strictly concave and singular (∞) (and in particular $K(1) = K(-1) = -\infty$, too).

We will call a function $J : \mathbb{R} \rightarrow \underline{\mathbb{R}}$ an *external n -field function* on \mathbb{R} if it is 1-periodic, bounded above and it assumes finite values at more than n different points from $[0, 1)$. Comparing this definition with that of [7], we see that if J

is an external n -field function on \mathbb{R} , then J is an external n -field function on $[0, 1]$. In the opposite direction, if J is an external n -field function on $[0, 1]$ and $J(0) = J(1)$, then it can be extended 1-periodically to \mathbb{R} , to an external n -field function on \mathbb{R} . We use external n -field functions on \mathbb{R} from now on and for simplicity, we call them field functions. With a slight abuse of notation we will also consider them as external field functions on $\mathbb{T} := \mathbb{R}/\mathbb{Z}$: again, on \mathbb{T} the defining properties are that $J : \mathbb{T} \rightarrow \underline{\mathbb{R}}$, $J > -\infty$ at more than n points of \mathbb{T} , and J is upper bounded.

For a field function J we define its *singularity set* and *finiteness domain* by

$$X := X_J := J^{-1}(\{-\infty\}) \cap [0, 1] \quad \text{and} \quad X^c := [0, 1] \setminus X = J^{-1}(\mathbb{R}) \cap [0, 1]. \quad (8)$$

Then X^c has cardinality exceeding n , in particular $X \neq [0, 1]$. Considering J as defined on \mathbb{T} , we can replace $[0, 1]$ with \mathbb{T} in all the above.

Given $n \in \mathbb{N}$ and n kernel functions K_1, \dots, K_n , and an n -field function J , pure sum of translates and sum of translates functions are defined as

$$f(\mathbf{x}, r) := \sum_{j=1}^n K_j(r - x_j), \quad (9)$$

$$F(\mathbf{x}, r) := J(r) + f(\mathbf{x}, r) \quad (10)$$

where $r \in \mathbb{R}$, and $\mathbf{x} = (x_1, x_2, \dots, x_n) \in [0, 1]^n$; or, analogously, we can define $f(\mathbf{y}, t)$ and $F(\mathbf{y}, t)$ also for $t \in \mathbb{T}$, $\mathbf{y} \in \mathbb{T}^n$.

2.3 Differences between the torus and the interval setting

A useful step in several of our arguments—already in [5]—is the “cutting up” of the torus at an arbitrary point $c \in \mathbb{T}$. To formalize it, we introduce the mapping $\pi_c : \mathbb{R} \rightarrow \mathbb{T}$, $\pi_c(r) := \{r + c\} = r + c \pmod{1}$. That constitutes a (multiple) covering mapping of \mathbb{T} (hence in particular it is continuous), and it is bijective on $[0, 1)$, so its inverse $\pi_c^{-1} := (\pi_c|_{[0, 1)})^{-1} : \mathbb{T} \rightarrow [0, 1)$ is bijective, too. However, in the inverse direction the mapping ceases to remain continuous: it is continuous at all $t \neq c$, $t \in \mathbb{T}$, but at c it has a jump.

Cutting up is particularly useful when we want to prove local results like e.g. continuity of some mappings at $\mathbf{x} \in \mathbb{T}^n$. Choosing c appropriately, node systems from \mathbb{T}^n , subject to some ordering restriction and close to \mathbf{x} may correspond to node systems in $[0, 1)^n$ admitting a specific ordering in the interval. However, ordered node systems do not have a global match on \mathbb{T} . We will detail this phenomenon below.

In [5] we introduced, for any permutation $\sigma : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$, the corresponding simplex on $[0, 1]$ as

$$S_{[0, 1]}^{(\sigma)} := \{(x_1, \dots, x_n) \in [0, 1]^n : 0 < x_{\sigma(1)} < x_{\sigma(2)} < \dots < x_{\sigma(n)} < 1\} \subset \mathbb{R}^n.$$

Its closure is

$$\overline{S}_{[0, 1]}^{(\sigma)} = \{(x_1, \dots, x_n) \in [0, 1]^n : 0 \leq x_{\sigma(1)} \leq x_{\sigma(2)} \leq \dots \leq x_{\sigma(n)} \leq 1\} \subset \mathbb{R}^n.$$

An essential difference between the torus and interval setup is that in $[0, 1]$ we cannot perturb in both directions the nodes lying at the endpoints: they can be moved only towards the interval center. That restriction could be considered responsible for the need of some monotonicity assumption about the kernels when proving minimax etc. results for the interval case. However, in the torus a monotonicity assumption is in fact impossible: a periodic and monotone kernel function would necessarily be constant. On the other hand we had already seen in [5] that perturbation of node systems on the torus, in particular when we are free to decide about the direction of change of the nodes, are very useful. As a consequence, we need to consider all node systems, which may arise by means of such a perturbation, together. So, we need to consider the case, when a node passes over 0 and reappears at 1, as the same ordering. In fact, that is very natural: on the torus there is no strict ordering, but only an orientation, and the “order of nodes” can only be fixed as up to rotation. This we may call cyclic ordering. In this sense we may write $x_1 \preccurlyeq \dots \preccurlyeq x_n$ if starting from $x_1 \in \mathbb{T}$ and moving in the counterclockwise direction (that is, according to the positive orientation of the circle), we pass the points in the order of their listing until after a full rotation we arrive at the initial point x_1 . Similarly for the strict precedence notation $x_1 \prec \dots \prec x_n$. Correspondingly, arcs are defined as the set of points between two endpoints: $[a, b] := \{x \in \mathbb{T} : a \preccurlyeq x \preccurlyeq b\}$ etc. Note that for $n = 2$ $x_1 \preccurlyeq x_2$ and also $x_2 \preccurlyeq x_1$ hold simultaneously for all points $x_1, x_2 \in \mathbb{T}$, and that cyclic ordering of n nodes is possible in $(n - 1)!$ different ways.

The “large simplex on the torus” and its closure is defined as

$$L := \{(y_1, y_2, \dots, y_n) \in \mathbb{T}^n : y_1 \prec y_2 \prec \dots \prec y_n (\prec y_1)\}, \quad (11)$$

$$\overline{L} := \{(y_1, y_2, \dots, y_n) \in \mathbb{T}^n : y_1 \preccurlyeq y_2 \preccurlyeq \dots \preccurlyeq y_n (\preccurlyeq y_1)\}. \quad (12)$$

One may consider various permutations for the cyclic ordering, but by relabeling we will always assume that the ordering is just this natural cyclic order.

Pulling back by any π_c^{-1} coordinatewise we see that

$$\pi_c^{-1}(\overline{L}) = \cup_{j=0}^{n-1} \overline{S}_{[0,1]}^{(\sigma_j)}, \quad \sigma_j(\ell) := \begin{cases} \ell + j, & \text{if } \ell = 1, \dots, n - j \\ \ell + j - n, & \text{if } \ell = n - j + 1, \dots, n. \end{cases} \quad (13)$$

This decomposition is not disjoint, given that \overline{L} is connected. Similarly, no representation of L by disjoint $S_{[0,1]}^{(\sigma)}$ exists, as L is connected, too.

For $\mathbf{x} \in \overline{S}_{[0,1]}^{(\sigma)}$, we have defined the intervals

$$I_0(\mathbf{x}) := [0, x_{\sigma(1)}], \quad I_j(\mathbf{x}) := [x_{\sigma(j)}, x_{\sigma(j+1)}] \quad (1 \leq j \leq n - 1), \quad I_n(\mathbf{x}) := [x_{\sigma(n)}, 1],$$

and the corresponding “interval maxima” (in fact, supremums, not maximums) as

$$m_j(\mathbf{x}) := \sup\{F(\mathbf{x}, t) : t \in I_j(\mathbf{x})\}, \quad j = 0, 1, \dots, n.$$

Analogously, for an arbitrary $\mathbf{y} \in \overline{L}$ we define

$$I_j^*(\mathbf{y}) := \{t \in \mathbb{T} : y_j \preccurlyeq t \preccurlyeq y_{j+1}\}, \quad j = 1, 2, \dots, n - 1, \quad (14)$$

$$I_n^*(\mathbf{y}) := \{t \in \mathbb{T} : y_n \preccurlyeq t \preccurlyeq y_1\}, \quad (15)$$

and the corresponding ‘‘arc maximums’’

$$m_j^*(\mathbf{y}) := \sup\{F(\mathbf{y}, t) : t \in I_j^*(\mathbf{y})\}, \quad j = 1, 2, \dots, n.$$

The correspondence between $I_k^*(\mathbf{y})$ and $I_j(\mathbf{x})$, and also between $m_k^*(\mathbf{y})$ and $m_j(\mathbf{x})$, can be described easily. Fix $c \in \mathbb{T}$ arbitrarily and let $\mathbf{y} \in \bar{L}$. Then $\pi_c^{-1}(y_1), \dots, \pi_c^{-1}(y_n) \in [0, 1)$, moreover, setting $x_j := \pi_c^{-1}(y_j)$, the coordinates of the vector $\mathbf{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$ follow according to the cyclic ordering of the coordinates of \mathbf{y} , that is, if $y_j \prec c \preccurlyeq y_{j+1}$, then $\mathbf{x} \in \bar{S}_{[0,1]}^{(\sigma_j)}$. Extending the definition of π_c and its inverse to vectors, we may write $\pi_c^{-1}(\mathbf{y}) = \mathbf{x} \in \bar{S}_{[0,1]}^{(\sigma_j)}$. Then we find $\pi_c(I_k(\mathbf{x})) = \pi_c([x_{\sigma_j(k)}, x_{\sigma_j(k+1)}]) = I_k^*(\mathbf{y})$ for $k = 1, \dots, n-1$, while $\pi_c(I_0(\mathbf{x})) \cup I_n(\mathbf{x}) = I_n^*(\mathbf{y})$. Accordingly, the interval and arc maxima correspond to each other as follows.

$$m_k^*(\mathbf{y}) = m_k(\mathbf{x}) \quad (1 \leq k \leq n-1), \quad m_n^*(\mathbf{y}) = \max(m_0(\mathbf{x}), m_n(\mathbf{x})). \quad (16)$$

Note that this representation does depend on the ordering of the x_k , because $I_0(\mathbf{x}) = [0, x_{\sigma_j(1)}] = [0, \pi_c^{-1}(y_{j+1})] = \pi_c^{-1}([c, y_{j+1}])$, and $I_n(\mathbf{x}) = [x_{\sigma_j(n)}, 1] = [\pi_c^{-1}(y_j), 1] = \pi_c^{-1}([y_j, c])$.

In [7, 9, 6] we have already investigated the following minimax and maximin problems on the interval $[0, 1]$.

$$\begin{aligned} \bar{m}(\mathbf{x}) &:= \max_{j=0,1,\dots,n} m_j(\mathbf{x}) = \sup\{F(\mathbf{x}, r) : r \in [0, 1]\}, & \underline{m}(\mathbf{x}) &:= \min_{j=0,1,\dots,n} m_j(\mathbf{x}), \\ M(\bar{S}_{[0,1]}) &:= \inf\{\bar{m}(\mathbf{x}) : \mathbf{x} \in \bar{S}_{[0,1]}\}, & m(\bar{S}_{[0,1]}) &:= \sup\{\underline{m}(\mathbf{x}) : \mathbf{x} \in \bar{S}_{[0,1]}\}. \end{aligned}$$

The analogous quantities on the torus are

$$\begin{aligned} \bar{m}^*(\mathbf{y}) &:= \max_{j=1,\dots,n} m_j^*(\mathbf{y}) = \sup\{F(\mathbf{y}, t) : t \in \mathbb{T}\}, & \underline{m}^*(\mathbf{y}) &:= \min_{j=1,\dots,n} m_j^*(\mathbf{y}), \\ M^*(\bar{L}) &:= \inf\{\bar{m}^*(\mathbf{y}) : \mathbf{y} \in \bar{L}\}, & m^*(\bar{L}) &:= \sup\{\underline{m}^*(\mathbf{y}) : \mathbf{y} \in \bar{L}\}. \end{aligned}$$

Note that $m_n^*(\pi_c(\mathbf{x})) = \max(m_n(\mathbf{x}), m_0(\mathbf{x}))$,

$$\bar{m}^*(\pi_c(\mathbf{x})) = \bar{m}(\mathbf{x}), \quad (17)$$

and in view of (13) we also have² $M^*(\bar{L}) = \min_{j=1,\dots,n} M(\bar{S}^{(\sigma_j)})$. However, there is no similar easy formula for $m^*(\bar{L})$.

3 Continuity results

In this section, we collect the continuity properties of maximum functions. First, we recall Lemma 3.3 from [9]:

²In fact, this formula provides an alternative way to prove Theorem 12, but not of Theorem 13, so we have decided to follow the forthcoming approach.

Lemma 2. *Let $n \in \mathbb{N}$, $\nu_j > 0$ ($j = 1, \dots, n$), let J be an n -field function on $[0, 1]$, and let K be a kernel function on $[-1, 1]$.*

Then $\overline{m} : [0, 1]^n \rightarrow \mathbb{R}$ is continuous.

From this we easily deduce the following.

Proposition 3. *Let $n \in \mathbb{N}$, $\nu_j > 0$ ($j = 1, \dots, n$), let J be a field function and let K be a kernel function.*

Then $\overline{m}^ : \mathbb{T}^n \rightarrow \mathbb{R}$ is continuous.*

Proof. Let $\mathbf{a} \in \mathbb{T}^n$, $\mathbf{a} = (a_1, \dots, a_n)$ be fixed and $c \in \mathbb{T} \setminus \{a_1, \dots, a_n\}$. We show that \overline{m}^* is continuous in a small neighborhood of \mathbf{a} . Let $\delta_0 < \min_{j=1, \dots, n} \text{dist}_{\mathbb{T}}(c, a_j)$. We pull back \mathbf{y} to $[0, 1]^n$ coordinatewise: $x_j := \pi_c^{-1}(y_j)$, $j = 1, \dots, n$. Then $x_j \in (0, 1)$ and $x_j = x_j(y_j)$ is continuous (since $\text{dist}_{\mathbb{T}}(y_j, a_j) < \delta_0$, and π_c^{-1} is continuous save at c), so $\mathbf{x} := \mathbf{x}(\mathbf{y}) := \pi_c^{-1}(\mathbf{y})$ is changing continuously in the given neighborhood of \mathbf{a} . We may write that $F(\mathbf{y}, t) = F(\pi_c(\mathbf{x}(\mathbf{y})), \pi_c(r))$ where $\pi_c(r) = t$ and after simplifying the notation, we simply write $F(\mathbf{y}, t) = F(\mathbf{x}(\mathbf{y}), r)$, with $t \in \mathbb{T}$ corresponding to $r := \pi_c^{-1}(t) \in [0, 1]$.

Therefore, with (17), we see that $\overline{m}^*(\mathbf{y}) = \overline{m}(\mathbf{x}(\mathbf{y}))$.

The continuity of \overline{m}^* at \mathbf{a} follows from the continuity of $\mathbf{x}(\mathbf{y})$ at \mathbf{a} and the continuity of \overline{m} at \mathbf{x} , the latter coming from Lemma 2. \square

We show continuity of the arc maxima functions m_k^* in some important cases.

Proposition 4. *Let $n \in \mathbb{N}$ and $k \in \{1, 2, \dots, n\}$ be fixed and let K_1, \dots, K_n be arbitrary kernel functions.*

- (a) *Suppose that J is an arbitrary n -field function and all K_j , $j = 1, 2, \dots, n$ satisfy (∞) . Then m_k^* is extended continuous on \overline{L} .*
- (b) *Suppose that J is an extended continuous field function. Then m_k^* is extended continuous on \overline{L} .*
- (c) *If J is an upper semicontinuous n -field function, then m_k^* is upper semicontinuous on \overline{L} .*

Proof. To see (a), let $\mathbf{a} \in \overline{L}$ be fixed. If $I_k^*(\mathbf{a}) \neq \mathbb{T}$, then let $c \in \mathbb{T} \setminus I_k^*(\mathbf{a})$, $c \notin \{a_1, \dots, a_n\}$. Then there is a $j \in \{0, 1, \dots, n-1\}$ (see (13)) such that $\pi_c^{-1}(\mathbf{a}) \in \overline{S}_{[0,1]}^{(\sigma_j)}$. Moreover, $0 < \pi_c^{-1}(a_1), \dots, \pi_c^{-1}(a_n) < 1$.

So, with (16) we can write $m_k^*(\mathbf{y}) = m_k(\mathbf{x})$ when $\mathbf{y} \in \overline{L}$ is close to \mathbf{a} and $\mathbf{x} = \pi_c^{-1}(\mathbf{y})$, $\mathbf{x} \in \overline{S}_{[0,1]}^{(\sigma_j)}$. Since $\pi_c^{-1} : \overline{L} \rightarrow \overline{S}_{[0,1]}^{(\sigma_j)}$ is continuous near \mathbf{a} , and by Lemma 3.1 from [9], $m_j : \overline{S}_{[0,1]}^{(\sigma_j)} \rightarrow \mathbb{R}$ is continuous, we obtain the assertion of this part.

If $I_k^*(\mathbf{a}) = \mathbb{T}$, then we follow the same steps with $c \in \mathbb{T} \setminus \{a_k\}$, but the arc $I_k^*(\mathbf{a})$ —and all arcs $I_k^*(\mathbf{y})$ with \mathbf{y} close to \mathbf{a} —necessarily split into two intervals via $\pi_c^{-1}(\cdot)$, so we use the second half of (16), and we get $m_k^*(\mathbf{y}) = \max(m_0(\mathbf{x}), m_n(\mathbf{x}))$ when $\mathbf{y} \in \overline{L}$ is close to \mathbf{a} . Continuing with the same steps, we obtain the assertion.

The proof of (b) is straightforward.

The proof of (c) follows the same steps as that of (a), using Proposition 3.6 (a) from [9] and that the maximum of two upper semicontinuous function is again upper semicontinuous. \square

Example 5. *If J is not continuous, and K_j does not satisfy (∞) then m_j^* is not continuous on \bar{L} .*

To see this, take $y^ \in \mathbb{T}$, where J is not continuous: we may assume $y^* = 1/2$. Let $n = 2$ and consider $\mathbf{x} = (x, x)$ where $x \approx 1/2$. Then $m_1^*(\mathbf{x}) = \sup_{I_1^*(\mathbf{x})} F(\mathbf{x}, \cdot) = F(\mathbf{x}, x) = J(x) + 2K(0)$ and $m_1^*((1/2, 1/2)) = F((1/2, 1/2), 1/2) = J(1/2) + 2K(0)$. Hence $m_1^*(\cdot)$ is not continuous at $(1/2, 1/2)$.*

Let us remark the Berge proved a maximum theorem about partial maxima of bivariate functions, see, e.g., [1], but that result is not applicable here since his approach requires bivariate continuity. In our case, J may be discontinuous, and continuity of m_j or \bar{m} is thus nontrivial.

4 Perturbation lemmas

The first perturbation lemma describes the behavior of sum of translates functions when two nodes are pulled apart. It appeared in several forms, e.g., in [5] (see Lemma 11.5), [12], Lemma 10 on p. 1069, or [7], Lemma 3.1. A similar form can be found in [10] (see around formula (15) too).

Lemma 6 (Perturbation lemma). *Let K be a kernel function which is periodic (6). Let $0 \leq \alpha < a < b < \beta \leq 1$ and $p, q > 0$. Set*

$$\mu := \frac{p(a - \alpha)}{q(\beta - b)}. \quad (18)$$

(a) *If $\mu = 1$, then*

$$pK(t - \alpha) + qK(t - \beta) \leq pK(t - a) + qK(t - b). \quad (19)$$

holds for every $t \in [0, \alpha] \cup [\beta, 1]$.

(b) *Additionally, if K is strictly concave, then (19) holds with strict inequality.*

(c) *If $\mu = 1$, then*

$$pK(t - \alpha) + qK(t - \beta) \geq pK(t - a) + qK(t - b). \quad (20)$$

holds for every $t \in [a, b]$.

(d) *Additionally, if K is strictly concave, then (20) holds with strict inequality.*

Lemma 7 (Trivial Lemma). *Let $f, g, h : D \rightarrow \underline{\mathbb{R}}$ be functions on some Hausdorff topological space D and assume that*

- (i) either f, g, h are all upper semicontinuous,
- (ii) or f, g are extended continuous and h is locally upper bounded, but otherwise arbitrary.

Let $\emptyset \neq A \subseteq B \subseteq D$ be arbitrary. Assume

$$f(t) < g(t) \quad \text{for all } t \in A. \quad (21)$$

If $A \subseteq B$ is a compact set, then

$$\sup_A (f + h) < \sup_B (g + h) \quad \text{unless} \quad h \equiv -\infty \quad \text{on} \quad A. \quad (22)$$

Proof. The straightforward proof of (i) was given in [7] as Lemma 3.2. The proof of (ii) is similar, so we leave it to the reader. \square

The following lemma is rather similar to Lemma 4.1. of [7]. However, there are several differences, too, in which the below version is stronger than the former version. First, here we do not assume upper semicontinuity of the field function, which was made possible by the observation that there is a version of the Trivial Lemma which relaxes on that condition on h (even if using a little more assumption regarding continuity of f and g , which, on the other hand, are clearly available). Second, we assume non-degeneracy $w_{i+1} > w_i$ only for indices $i \in \mathcal{I}$, again a delicate novelty in the current version. Third, we drop the condition that K be monotone, an essentially necessary assumption for the interval $[0, 1]$, but, as is already told in the Introduction, not required in the periodic case. In view of all these differences, as well as in regard of the slightly different setup of having only n arcs defined by the n node points (and not $n + 1$ intervals), we will present the full proof of the Lemma, even if its basic idea and a large part of the details are repeating the former argument.

Lemma 8 (General maximum perturbation lemma on the torus). *Let $n \in \mathbb{N}$ be a natural number, and let $\nu_1, \dots, \nu_n > 0$ be given positive coefficients. Let K be a kernel function on \mathbb{T} , and let J be an arbitrary n -field function.*

Let $\mathbf{w} \in \overline{L}$ and $\mathcal{I} \cup \mathcal{J} = \{1, \dots, n\}$ be a non-trivial partition, and assume that for all $i \in \mathcal{I}$, we have $w_i < w_{i+1}$ (which holds in particular, independently of \mathcal{I} , if $\mathbf{w} \in L$).

Then, arbitrarily close to \mathbf{w} , there exists $\mathbf{w}' \in \overline{L} \setminus \{\mathbf{w}\}$, essentially different from and less degenerate than \mathbf{w} in the sense that

$$w'_\ell \neq w_\ell \quad \text{unless} \quad \{\ell - 1, \ell\} \subset \mathcal{I} \quad \text{or} \quad \{\ell - 1, \ell\} \subset \mathcal{J} \quad (23)$$

and³

$$w'_\ell \neq w'_{\ell+1} \quad \text{unless} \quad \{\ell - 1, \ell, \ell + 1\} \subset \mathcal{J} \quad \text{and} \quad w_\ell = w_{\ell+1}, \quad (24)$$

³Note that here $w_\ell = w_{\ell+1}$ excludes $\ell \in \mathcal{I}$, so only \mathcal{J} can contain all three indices listed.

(in particular, if $\mathbf{w} \in L$ then necessarily $\mathbf{w}' \in L$), and such that it satisfies

$$F(\mathbf{w}', t) \leq F(\mathbf{w}, t) \text{ for all } t \in I_i^*(\mathbf{w}') \quad \text{and} \quad I_i^*(\mathbf{w}') \subseteq I_i^*(\mathbf{w}) \text{ for all } i \in \mathcal{I}; \quad (25)$$

$$F(\mathbf{w}', t) \geq F(\mathbf{w}, t) \text{ for all } t \in I_j^*(\mathbf{w}) \quad \text{and} \quad I_j^*(\mathbf{w}') \supseteq I_j^*(\mathbf{w}) \text{ for all } j \in \mathcal{J}. \quad (26)$$

As a result, we also have

$$m_i^*(\mathbf{w}') \leq m_i^*(\mathbf{w}) \text{ for } i \in \mathcal{I} \quad \text{and} \quad m_j^*(\mathbf{w}') \geq m_j^*(\mathbf{w}) \text{ for } j \in \mathcal{J} \quad (27)$$

for the corresponding torus maxima.

Moreover, if K is strictly concave, then the inequalities in (25) and (26) are strict for all points in the respective arcs where $J(t) \neq -\infty$.

Furthermore, for strictly concave K the inequalities in (27) are also strict for all indices k with non-singular $I_k^*(\mathbf{w})$.

Proof. Before the main argument, we observe that the assertion in (27) is indeed a trivial consequence of the previous inequalities (26) and (25), so we need not give a separate proof for that.

Second, the inequalities (25) and (26) follow from

$$f(\mathbf{w}', t) \leq f(\mathbf{w}, t) \quad (\forall t \in I_i^*(\mathbf{w}')) \quad \text{and} \quad I_i^*(\mathbf{w}') \subseteq I_i^*(\mathbf{w}) \quad \text{for all } i \in \mathcal{I}; \quad (28)$$

$$f(\mathbf{w}', t) \geq f(\mathbf{w}, t) \quad (\forall t \in I_j^*(\mathbf{w})) \quad \text{and} \quad I_j^*(\mathbf{w}') \supseteq I_j^*(\mathbf{w}) \quad \text{for all } j \in \mathcal{J}. \quad (29)$$

Moreover, strict inequalities for all points t with $J(t) \neq -\infty$ will follow from (25) and (26) if we can prove strict inequalities in (28) and (29) for all values of t in the said compact arcs.

Furthermore, in case we have strict inequalities in (28) and (29) for all points t , then for non-singular $I_k^*(\mathbf{w})$ this entails strict inequalities also in (27) (for the corresponding k). To see this, one may refer back to the Trivial Lemma 7 (ii) with $\{f, g\} = \{f(\mathbf{w}, \cdot), f(\mathbf{w}', \cdot)\}$, $h = J$, $\{A, B\} = \{I_k^*(\mathbf{w}), I_k^*(\mathbf{w}')\}$. Here in the case when $k = i \in \mathcal{I}$ we need to use that the arc $I_i^*(\mathbf{w})$ is not degenerate for $i \in \mathcal{I}$, hence if it is nonsingular, too, then either $J|_{I_i^*(\mathbf{w}')} \equiv -\infty$ and then we have the strict inequality $m_i^*(\mathbf{w}) > -\infty = m_i^*(\mathbf{w}')$, or $J(t) > -\infty$ at some points of $I_i^*(\mathbf{w}')$, and then using this Lemma furnishes the required strict inequality. (The other case with $k = j \in \mathcal{J}$ is easier because nonsingularity of $I_j^*(\mathbf{w})$ implies nonsingularity of the larger arc $I_j^*(\mathbf{w}')$, too, and then the application of the Lemma need not be coupled by considerations of an identically $-\infty$ field.)

So the proof hinges upon showing (23), (24), (28) and (29), for any n -field function and any kernel function, coupled with the strict inequality assertion in (28) and (29) for all t belonging to the said compact arcs, in case K is strictly concave.

For $n = 0$ or $n = 1$ there is no nontrivial partition of the index set $\{1, \dots, n\}$, hence the assertion is void and true.

For $n = 2$ there is essentially only one way to split the index set in a nontrivial way, so the statement will be part of the following, more general setup in Case

0, which we prove directly. Actually, the $n = 2$ case can be proved directly from Lemma 6, but we will need the more general Case 0 anyway.

Case 0. We prove directly the assertion when \mathcal{I}, \mathcal{J} contain no neighboring indices, so n must be even and \mathcal{I} and \mathcal{J} partition $\{1, \dots, n\}$ into the subsets of odd and even natural numbers from 1 to n .

We can assume that $\mathcal{I} = (2\mathbb{N} + 1) \cap \{1, \dots, n\}$ and $\mathcal{J} = 2\mathbb{N} \cap \{1, \dots, n\}$ (the other case being a simple change of the cut, i.e., starting of the listing of the cyclic ordering of nodes from one node later).

Note that whenever $w_j \in \mathcal{J}$ (i.e., when j is even), then we necessarily have $j - 1, j + 1 \in \mathcal{I}$, and $w_{j-1} < w_j \leq w_{j+1} < w_{j+2}$.

Denote $\delta := \min_{i \in \mathcal{I}} |I_i^*(\mathbf{w})|$, (where $|I_i^*(\mathbf{w})|$ is the length of the arc $[w_i, w_{i+1}]$) which is positive by condition.

Our new perturbed node system \mathbf{w}' will be, with an arbitrary $0 < h < \frac{1}{2}\delta / \max\{\nu_1, \dots, \nu_n\}$, the system

$$\mathbf{w}' := (w'_1, \dots, w'_n) \quad \text{with} \quad w'_\ell := w_\ell - (-1)^\ell \frac{1}{\nu_\ell} h, \quad \ell = 1, 2, \dots, n. \quad (30)$$

The definition guarantees, that $w'_\ell = w_\ell$ happens for no index ℓ , furnishing (23).

It is easy to see that by the choice of the perturbation lengths, no two consecutive nodes will change ordering or reach each other: for $j \in \mathcal{J}$ w_j, w_{j+1} are changed to become farther away, while for $i \in \mathcal{I}$ w_i, w_{i+1} are moved closer, but only by $\nu_i h + \nu_{i+1} h < \delta < w_{i+1} - w_i$. It follows that the ordering of nodes remains in \bar{L} . (In a minute we will see that, moreover, there remains no degenerate arc, so $\mathbf{w}' \in L \setminus \{\mathbf{w}\}$.)

Obviously, $I_j^*(\mathbf{w}') \supset I_j^*(\mathbf{w})$ holds for all $j \in \mathcal{J}$, and $I_i^*(\mathbf{w}') \subset I_i^*(\mathbf{w})$ holds for all $i \in \mathcal{I}$, with the inclusions strict, furnishing the second parts of (25) and (26) (matching the second parts of (28) and (29), too). In particular, even if $I_j^*(\mathbf{w})$ may be degenerate for some $j \in \mathcal{J}$, i.e., for some even j , the j th arc of the perturbed system will not be such: $w'_j < w_j \leq w_{j+1} < w'_{j+1}$ for any even $j \in \mathcal{J}$. We get $\mathbf{w}' \in L$, entailing (24).

Take now an even indexed arc $I_{2k}^*(\mathbf{w}) = [w_{2k}, w_{2k+1}]$, so $2k \in \mathcal{J}$. (When $2k = n$, then we must read $2k + 1 = n + 1 \equiv 1$, i.e. $w_{2k+1} = w_1$.) Our perturbation of nodes in (30) can now be grouped as *pairs of changing nodes* $w_{2\ell-1}, w_{2\ell}$ among w_1, \dots, w_{2k} , and then again among w_{2k+1}, \dots, w_n , recalling that n is even. Now, *the pairs* are always changed so that the arcs in between shrink, and shrink exactly as is described in Lemma 6. We apply this lemma for each pair of such nodes with the choices $a = w'_{2\ell-1}$, $b = w'_{2\ell}$, $\alpha = w_{2\ell-1}$, $\beta = w_{2\ell}$, $p = \nu_{2\ell-1}$, $q = \nu_{2\ell}$. This gives that for each such pair of changes, for t *outside of the enclosed arc* $(w_{2\ell-1}, w_{2\ell}) = I_{2\ell-1}^*(\mathbf{w})$ we have

$$\nu_{2\ell-1} K(t - w'_{2\ell-1}) + \nu_{2\ell} K(t - w'_{2\ell}) \geq \nu_{2\ell-1} K(t - w_{2\ell-1}) + \nu_{2\ell} K(t - w_{2\ell}). \quad (31)$$

Note that $I_{2k}^*(\mathbf{w})$, hence any $t \in I_{2k}^*(\mathbf{w})$, is *always outside* of the arcs $I_{2\ell-1}^*(\mathbf{w})$,

therefore (31) holds for the given, fixed $t \in I_{2k}^*(\mathbf{w})$ and for all ℓ . So we find

$$\begin{aligned} f(\mathbf{w}, t) &= \sum_{\ell=1}^{n/2} (\nu_{2\ell-1} K(t - w_{2\ell-1}) + \nu_{2\ell} K(t - w_{2\ell})) \\ &\leq \sum_{\ell=1}^{n/2} (\nu_{2\ell-1} K(t - w'_{2\ell-1}) + \nu_{2\ell} K(t - w'_{2\ell})) = f(\mathbf{w}', t). \end{aligned} \quad (32)$$

Furthermore, all the appearing inequalities are strict in case K is strictly concave. We have proved (29), even with strict inequality under appropriate assumptions.

The proof of (28) runs analogously by grouping the change of nodes as a change of pairs $w_{2\ell}, w_{2\ell+1}$ for $\ell = 1, \dots, n/2$, writing $w_{2(n/2)+1} = w_{n+1} = w_1$ according to periodicity. For these arcs $2\ell \in \mathcal{J}$ and the arcs are getting larger after the perturbation, so outside these enlarged arcs—that is, for all points which belong to any $I_{2k-1}^*(\mathbf{w}')$ for some fixed k —the changed value $f(\mathbf{w}', t)$ will not exceed (and in case of strict concavity, will be strictly smaller than) $f(\mathbf{w}, t)$. This means a nonincreasing (decreasing) change for all $I_i^*(\mathbf{w}')$, entailing (28) together with the respective strict inequality statement.

The proof of Case 0 is thus completed.

Therefore, we have also the case $n = 2$ proved. From here we continue our argumentation by induction. Let now $n > 2$ and assume, as inductive hypothesis, the validity of the assertions for all $n^* \leq \tilde{n} := n - 1$ and for any choice of kernel- and n^* -field functions.

In view of Case 0 above, there remains the case when there are neighboring indices $k - 1, k$ belonging to the same index set \mathcal{I} or \mathcal{J} . In view of the cyclic ordering and to avoid indexing complications, assume that we also have $1 < k < n$, which is a possibility for any n at least 3. We separate two cases.

Case 1. Assume first that $w_{k-1} < w_k < w_{k+1}$ holds.

Then we consider the kernel function $\tilde{K} := K$, and the \tilde{n} -field function $\tilde{\mathcal{J}} := \nu_k K(\cdot - w_k)$ (which is indeed an \tilde{n} -field function because it attains $-\infty$ only at most at one point, namely w_k , in case K is singular.)

Correspondingly, now the sum of translates function \tilde{F} is formed by using $\tilde{n} = n - 1$ translates with coefficients $\nu_1, \dots, \nu_{k-1}, \nu_{k+1}, \dots, \nu_n$ and with respect to the node system

$$\tilde{\mathbf{w}} := (w_1, w_2, \dots, w_{k-1}, w_{k+1}, \dots, w_n).$$

Formally, the indices change: $\tilde{w}_\ell = w_\ell$ for $\ell = 1, \dots, k - 1$, but $\tilde{w}_\ell = w_{\ell+1}$ for $\ell = k, \dots, n - 1$, the k th coordinate being left out.

We apply the same change of indices in the partition: k is dropped out (but the corresponding index set \mathcal{I} or \mathcal{J} will not become empty, for it contains $k - 1$); and then shift indices one left for $\ell > k$: so

$$\begin{aligned} \tilde{\mathcal{I}} &:= \{i \in \mathcal{I} : i < k\} \cup \{i - 1 : i \in \mathcal{I}, i > k\} \text{ and} \\ \tilde{\mathcal{J}} &:= \{j \in \mathcal{J} : j < k\} \cup \{j - 1 : j \in \mathcal{J}, j > k\}. \end{aligned}$$

Observe that $\tilde{F}(\tilde{\mathbf{w}}, t) = f(\mathbf{w}, t)$ for all $t \in \mathbb{T}$, while

$$I_\ell^*(\tilde{\mathbf{w}}) = \begin{cases} I_\ell^*(\mathbf{w}), & \text{if } \ell < k-1, \\ I_{k-1}^*(\mathbf{w}) \cup I_k^*(\mathbf{w}), & \text{if } \ell = k-1, \\ I_{\ell+1}^*(\mathbf{w}), & \text{if } \ell \geq k. \end{cases}$$

(Here we make a little use of the choice that $1 < k < n$, so we need not bother too much with the cyclic renumbering etc.)

Note that $I_i^*(\tilde{\mathbf{w}}) = [\tilde{w}_i, \tilde{w}_{i+1}]$ is still nondegenerate whenever $i \in \tilde{\mathcal{I}}$, for the arc is either a former arc belonging to some $i \in \mathcal{I}$, or the union of two such arcs. Also, ordering of nodes is kept intact, so $\tilde{\mathbf{w}} \in \overline{\mathcal{L}}^{(\tilde{n})}$ (where $\overline{\mathcal{L}}^{(n)}$ and $\overline{\mathcal{L}}^{(\tilde{n})}$ denote the cyclic simplices of the corresponding dimension).

Now we apply the inductive hypothesis for the new configuration. This yields a perturbed node system $\tilde{\mathbf{w}}' \in \overline{\mathcal{L}}^{(\tilde{n})} \setminus \{\tilde{\mathbf{w}}\}$, arbitrarily close to $\tilde{\mathbf{w}}$, with the asserted properties. It is important that here the ordering of the nodes remain the order fixed in $\overline{\mathcal{L}}^{(\tilde{n})}$, so if $\tilde{\mathbf{w}}'$ was closer to $\tilde{\mathbf{w}}$ than the distance δ of w_k from $\{w_{k-1}, w_{k+1}\}$, then the n -term node system \mathbf{w}' , obtained by keeping the nodes from $\tilde{\mathbf{w}}'$ and inserting back w_k to the k th place (and shifting the following indices by one) will again be ordered as \mathbf{w} was, i.e., $\mathbf{w}' \in \overline{\mathcal{L}}^{(n)}$ (and of course $\neq \mathbf{w}'$, as already $\tilde{\mathbf{w}}' \neq \tilde{\mathbf{w}}$). Moreover, $w'_k = w_k$ is still not equal to any of the nodes w'_{k-1}, w'_{k+1} , because $\text{dist}_{\mathbb{T}^n}(\mathbf{w}', \mathbf{w}) < \delta$.

We now set to prove (23). First, if for some $\ell < k$ we have $w'_\ell = w_\ell$, then $\tilde{w}'_\ell =: w'_\ell = w_\ell =: \tilde{w}_\ell$, and then by the inductive hypothesis $\{\ell-1, \ell\} \subset \tilde{\mathcal{I}}$ or $\tilde{\mathcal{J}}$. This gives (23) in case $\ell < k$, because below k the partition sets \mathcal{I} and $\tilde{\mathcal{I}}$ (and \mathcal{J} and $\tilde{\mathcal{J}}$, respectively), consist of the same indices.

Take now some $\ell > k$ with $w'_\ell = w_\ell$. Then $\tilde{w}'_{\ell-1} =: w'_\ell = w_\ell =: \tilde{w}_{\ell-1}$ and $\{\ell-2, \ell-1\} \subset \tilde{\mathcal{I}}$ or $\tilde{\mathcal{J}}$ by the inductive hypothesis. If it was $\ell > k+1$, too, then this means $\{\ell-1, \ell\} \subset \mathcal{I}$ or \mathcal{J} , that is, (23).

If, however, $\ell = k+1$, then $\ell-2 = k-1$ and $\ell-1 = k$, and by construction we get that $\{k-1, k+1\} \subset \mathcal{I}$ or \mathcal{J} . Given that we already have from the outset that k belongs to the same index set as $k-1$, this altogether gives $\{k-1, k, k+1\} \subset \mathcal{I}$ or \mathcal{J} , which is more than needed for (23).

Finally, if $\ell = k$, then $w'_k = w_k$ by construction, but then we had by assumption that $k-1$ and k belonged to the same set \mathcal{I} or \mathcal{J} , so that (23) is satisfied.

Consider now the assertions of (24). As above, there is no problem with respective index sets all remaining the same or all being shifted by one, i.e., if either $\ell < k-1$ or if $\ell > k+1$.

Recall that we assumed $w_{k-1} < w_k < w_{k+1}$ at the outset, and chose the perturbation small enough to keep this strict ordering. Therefore, $w'_{k-1} = w'_k$ and $w'_k = w'_{k+1}$ are excluded, and only the case of $\ell = k+1$ and $w'_{k+1} = w'_{k+2}$ remains to be dealt with. Now, $\tilde{w}'_k =: w'_{k+1} = w'_{k+2} := \tilde{w}'_{k+1}$, so $\{k-1, k, k+1\} \subset \tilde{\mathcal{J}}$ and $\tilde{w}_k := \tilde{w}_{k+1}$ according to the inductive hypothesis. The latter means $w_{k+1} = w_{k+2}$, while for the indices we obtain $\{k-1, k+1, k+2\} \subset \mathcal{J}$.

But $k - 1$ and k belong to the same index set, so that also k must belong to \mathcal{J} , and therefore $\{k - 1, k, k + 1, k + 2\} \subset \mathcal{J}$, entailing (24).

Using that $k - 1$ and k belong to the same index set \mathcal{I} or \mathcal{J} , it is easy to check that $I_i^*(\tilde{\mathbf{w}}') \subseteq I_i^*(\tilde{\mathbf{w}})$ for all $i \in \tilde{\mathcal{I}}$ is equivalent to $I_i^*(\mathbf{w}') \subseteq I_i^*(\mathbf{w})$ for all $i \in \mathcal{I}$, and $I_j^*(\tilde{\mathbf{w}}') \supseteq I_j^*(\tilde{\mathbf{w}})$ for all $j \in \tilde{\mathcal{J}}$ is equivalent to $I_j^*(\mathbf{w}') \supseteq I_j^*(\mathbf{w})$ for all $j \in \mathcal{J}$. Further, the assertions (25), (26) from the inductive hypotheses lead to the assertions (28), (29) for the original case. Therefore, by the preliminary observations also (25), (26) follow. Moreover, the assertion regarding strict inequalities for all t in case of a strictly concave K follow from the respective strict inequalities for the inductive hypotheses, noting that $I_{k-1}^*(\tilde{\mathbf{w}}) = I_{k-1}^*(\mathbf{w}) \cup I_k^*(\mathbf{w})$ can handle the necessary inequalities for both indices $k - 1$ and k , because these belong to the same index set \mathcal{I} or \mathcal{J} , and hence invoke inequalities in the same direction.

Case 2. Consider now the case when some of the partition sets \mathcal{I}, \mathcal{J} contain some neighboring indices $k - 1, k$, such that $w_{k-1} = w_k$ or $w_k = w_{k+1}$ holds, too. Then this index set cannot be \mathcal{I} , for indices in \mathcal{I} the respective arcs were supposed to be nondegenerate. So, $k - 1, k \in \mathcal{J}$.

Repeating the above argument in Case 1 then works. Let us detail, why.

The main point where we needed that $w_{k-1} < w_k < w_{k+1}$ was where we wanted to see that the new node system $\tilde{\mathbf{w}}'$, provided by the induction hypothesis, not only preserves cyclic ordering of nodes from $\tilde{\mathbf{w}}$, but even with re-inserting w_k and thus manufacturing \mathbf{w}' will still result in a point belonging to $\overline{L}^{(n)}$.

Observe that the inductive hypothesis, in view of $k - 1, k \in \mathcal{J}$, furnishes that $I_{k-1}^*(\tilde{\mathbf{w}})$ is subject to growth, so it will still contain the point w_k , that is, $w'_{k-1} := \tilde{w}'_{k-1} \leq \tilde{w}_{k-1} := w_{k-1} \leq w_k \leq w_{k+1} =: \tilde{w}_k \leq \tilde{w}'_k =: w'_{k+1}$. It follows immediately that we will thus have $\mathbf{w}' \in \overline{L}$ at least.

Moreover, as above, equality of other perturbed and original nodes w'_ℓ and w_ℓ ($\ell \neq k$) can occur only when they occurred also in $\tilde{\mathbf{w}}$, hence in \mathbf{w} , too. Checking the arising conditions for the indices can be done mutatis mutandis the above case, proving (23), on noting that for $\ell = k$ we already have $k - 1, k \in \mathcal{J}$ by assumption.

To prove (24), assume now the identity $w'_\ell = w'_{\ell+1}$. Again, we separate cases according to the size of ℓ and start with the case $\ell < k - 1$. This implies $\tilde{w}'_\ell =: w'_\ell = w'_{\ell+1} := \tilde{w}'_{\ell+1}$, hence also $\tilde{w}_\ell = \tilde{w}_{\ell+1}$ and $\ell - 1, \ell, \ell + 1 \in \tilde{\mathcal{J}}$ in view of the inductive hypothesis, so $w_\ell = w_{\ell+1}$ and $\ell - 1, \ell, \ell + 1 \in \mathcal{J}$, too. Similarly, if $\ell > k$ then the identity $w'_\ell = w'_{\ell+1}$ implies $\tilde{w}'_{\ell-1} =: w'_\ell = w'_{\ell+1} := \tilde{w}'_\ell$, hence also $\tilde{w}_{\ell-1} = \tilde{w}_\ell$ and $\ell - 2, \ell - 1, \ell \in \tilde{\mathcal{J}}$ in view of the inductive hypothesis, so $w_\ell = w_{\ell+1}$ and if $\ell > k + 1$, then $\ell - 1, \ell, \ell + 1 \in \mathcal{J}$, too, while if $\ell = k + 1$, then we get only $k - 1, k + 1, k + 2 \in \mathcal{J}$, but then again we remind to $k \in \mathcal{J}$ and get $\{k - 1, k, k + 1, k + 2\} \subset \mathcal{J}$, proving (24).

It remains to deal with $\ell = k - 1$ and $\ell = k$, which did not occur in Case 1 above. Now, $\ell = k - 1$ means that we have the identity $w'_{k-1} = w'_k$. Since $k - 1, k \in \mathcal{J}$ by assumption, we also have $k - 1 \in \tilde{\mathcal{J}}$, hence $I_{k-1}^*(\tilde{\mathbf{w}})$ cannot

shrink, and therefore $\tilde{w}'_{k-1} \leq \tilde{w}_{k-1}$. Moreover, if we had strict inequality here, then we would have to have $w'_{k-1} := \tilde{w}'_{k-1} < \tilde{w}_{k-1} := w_{k-1} \leq w_k =: w'_k$, although we supposed the contrary now. So, we must have $\tilde{w}'_{k-1} = \tilde{w}_{k-1}$, the inductive hypothesis applies, and we derive that both $k-2$ and $k-1$ belong to the same index set – that is, because of $k-1$, to \mathcal{J} . However, we already know by assumption also $k \in \mathcal{J}$, so altogether $\{k-2, k-1, k\} \subset \mathcal{J}$, as needed.

The case $\ell = k$ is similar. If $w'_k = w'_{k+1}$, then taking into account that $I_{k-1}^*(\tilde{\mathbf{w}})$ cannot shrink (as $k-1 \in \tilde{\mathcal{J}}$) we must have $w'_{k+1} := \tilde{w}'_k \geq \tilde{w}_k := w_{k+1} \geq w_k = w'_k$ entailing that all inequalities are in fact equalities, and in particular both $w_{k+1} = w_k$ and $\tilde{w}'_k = \tilde{w}_k$. Referring to the inductive hypothesis this furnishes $k-1, k \in \tilde{\mathcal{J}}$, that is, $k-1, k+1 \in \mathcal{J}$, whilst $k \in \mathcal{J}$ by the original condition, altogether yielding $\{k-1, k, k+1\} \subset \mathcal{J}$, as needed.

So, we proved (23) and (24) for this case, too. The proof for the remaining inequalities and strictness of them in case of a strictly concave kernel is identical to the argument in Case 1.

We thus conclude the proof of Case 2, whence the whole Lemma. \square

Remark 9. Although the formulation of the Lemma is a bit complicated, one may note that assuming $w_i < w_{i+1}$ for all $i \in \mathcal{I}$ is absolutely natural and minimal. Natural, because if we want to decrease the arcs $I_i^*(\mathbf{w})$ for all $i \in \mathcal{I}$, then these arcs must shrink, hence we cannot perform this change when they are already degenerate one point intervals. Minimal, because we do not assume similar conditions for any $I_j^*(\mathbf{w})$ with $j \in \mathcal{J}$, so even the less that $\mathbf{w} \in L$.

Remark 10. If we had $\mathbf{w} \in L$ from the outset, then we will as well have $\mathbf{w}' \in L$.

Corollary 11. *If we only want respective inequalities for the m_j^* , without requiring strict inclusions regarding the underlying intervals, then we can apply a further perturbation, now leading to $\mathbf{w}'' \in L$ (the point being that with different endpoint nodes!) and still satisfying the required strict inequalities between the m_j^* , provided that we had strict inequalities (so, e.g., K was strictly concave) and provided the m_j^* change continuously.*

Note that this latter condition of continuity of the m_j^* is satisfied if K is singular or if J is continuous, see Proposition 4 (a) and (c).

5 Minimax and maximin theorems

5.1 Minimax for strictly concave kernels

The following theorem contains, as a rather special case with the choice of the log-sine kernel $K(t) := \log |\sin(\pi t)|$, the above stated Theorem 1.

Theorem 12. *Let $n \in \mathbb{N}$ and $\nu_1, \dots, \nu_n > 0$, let K be a singular, strictly concave periodic kernel function, and let J be an arbitrary periodic n -field function.*

Then there exists a minimax point \mathbf{w} on \bar{L} , it belongs to the open “cyclic simplex” L , and it is an equioscillation point.

Proof. We already know that \overline{m}^* is continuous, thus it attains its infimum at a minimum point (where, in view of $\overline{m}^* : \mathbb{T}^n \rightarrow \mathbb{R}$, a finite minimax value is attained).

Now assume for a contradiction that the obtained minimax node system \mathbf{w} is not an equioscillating system. Then there are indices with $m_i^*(\mathbf{w}) = \overline{m}^*(\mathbf{w})$, but not all indices are such. So, take $\mathcal{I} := \{i : m_i^*(\mathbf{w}) = \overline{m}^*(\mathbf{w})\}$ and $\mathcal{J} := \{1, 2, \dots, n\} \setminus \mathcal{I} = \{j : m_j^*(\mathbf{w}) < \overline{m}^*(\mathbf{w})\}$. These index sets will define a nontrivial partition of the full set of indices from 1 to n .

In order to apply the Perturbation Lemma we will need that for $i \in \mathcal{I}$ the endpoint nodes are different: $w_i < w_{i+1}$. Given that $m_i^*(\mathbf{w}) = \overline{m}^*(\mathbf{w})$, this is certainly so in case K is singular, for then $m_\ell^*(\mathbf{w}) = -\infty < \overline{m}^*(\mathbf{w})$ for any degenerate arc $I_\ell^*(\mathbf{w})$. Let us now apply Lemma 8, which results in a new node system \mathbf{w}' , admitting the same cyclic ordering and lying arbitrarily close to \mathbf{w} , and with $m_i^*(\mathbf{w}') < \overline{m}^*(\mathbf{w})$ for all $i \in \mathcal{I}$. On the other hand, the $m_j^*(\mathbf{w}')$ may exceed $m_j^*(\mathbf{w})$ for $j \in \mathcal{J}$, but only by arbitrarily little, because \mathbf{w}' is sufficiently close to \mathbf{w} and the m_j^* 's are continuous on \overline{L} in view of the singularity of K , see Proposition 4 (a). So, in all, we will have $m_k^*(\mathbf{w}') < \overline{m}^*(\mathbf{w})$ for all $k = 1, \dots, n$, hence $\overline{m}^*(\mathbf{w}') < \overline{m}^*(\mathbf{w})$, contradicting to minimality of \mathbf{w} .

Noting that an equioscillation node system is necessarily nondegenerate for singular kernels, we conclude the proof. \square

5.2 Maximin for strictly concave kernels

Theorem 13. *Let $n \in \mathbb{N}$ and $\nu_1, \dots, \nu_n > 0$, let K be a singular, strictly concave periodic kernel function, and let J be an arbitrary periodic n -field function.*

Then there exists a maximin point \mathbf{z} on \overline{L} , it belongs to the open "cyclic simplex" L , and it is an equioscillation point.

Proof. Again, as all the m_k^* are continuous, so is their minimum. The maximum of that minimum is finite, because there are points with all m_k^* finite: e.g., take the above found minimax point \mathbf{w} . Given that K is assumed to be singular, that gives that such a point with $\underline{m}^*(\mathbf{z}) > -\infty$ cannot be degenerate, i.e., $\mathbf{z} \in L$. Also, by continuity of the m_k^* and hence of \underline{m}^* , there exists a maximin point $\mathbf{z} \in L$.

Assume for a contradiction that this point is not an equioscillation point. Consider $\mathcal{I} := \{i : m_i^*(\mathbf{z}) > \underline{m}^*(\mathbf{z})\}$ and $\mathcal{J} := \{1, \dots, n\} \setminus \mathcal{I} = \{j : m_j^*(\mathbf{z}) = \underline{m}^*(\mathbf{z})\}$. This is a nontrivial partition of the index set $\{1, \dots, n\}$, while $\mathbf{z} \in L$, hence the Perturbation Lemma 8 can be applied, and we are led to a new system $\mathbf{z}' \in L$, with $m_j^*(\mathbf{z}') > m_j^*(\mathbf{z}) = \underline{m}^*(\mathbf{z})$ for all $j \in \mathcal{J}$. However, we also have $m_i^*(\mathbf{z}') > m_i^*(\mathbf{z}) - \varepsilon > \underline{m}^*(\mathbf{z})$ for all $i \in \mathcal{I}$, if ε was chosen small enough and \mathbf{z}' close enough to \mathbf{z} , because m_i^* is continuous. So, in all, we find $\underline{m}^*(\mathbf{z}') > \underline{m}^*(\mathbf{z})$, and \mathbf{z} could not be a maximin point. The obtained contradiction proves the assertion. \square

5.3 Extension to concave kernel functions

To extend Theorem 12 to general concave kernels, we apply limiting arguments similar to [9], p. 18.

Theorem 14. *Let $n \in \mathbb{N}$ and $\nu_1, \dots, \nu_n > 0$, let K be a singular periodic kernel function, and let J be an arbitrary periodic n -field function.*

Then there exists a minimax point \mathbf{w}^ on \bar{L} , it belongs to the open “cyclic simplex” L , and it is an equioscillation point.*

Proof. Let K be a singular, 1-periodic kernel function which is not necessarily strictly concave. Let $K^{(\eta)}(t) := K(t) + \eta|\sin \pi t|$ where $\eta > 0$. Then $K^{(\eta)}$ is a strictly concave kernel function, which is also singular and 1-periodic. We will denote the corresponding maximum functions and minimax quantity by $m_j^*(\eta, \mathbf{y})$, $\bar{m}^*(\eta, \mathbf{y})$, and $M^*(\eta, \bar{L})$. Taking into account that $K^{(\eta)}$ converges to K uniformly, we find the same for $F(\eta, \cdot) \searrow F$ and hence even $\bar{m}^*(\eta, \mathbf{y}) \searrow \bar{m}^*(\mathbf{y})$ uniformly. Hence $M^*(\eta, L) \searrow M^*(L)$, too.

Let $\mathbf{e}(\eta) \in \bar{L}$ be a node system such that $\bar{m}^*(\eta, \mathbf{e}(\eta)) = M^*(\eta, \bar{L})$. By Theorem 12, $\mathbf{e}(\eta) \in L$ and it is an equioscillating node system: $m_1^*(\eta, \mathbf{e}(\eta)) = \dots = m_n^*(\eta, \mathbf{e}(\eta))$. Since \bar{L} is compact, there exists $\eta_k \searrow 0$ such that $\mathbf{e}(\eta_k) \rightarrow \mathbf{e}$ for some $\mathbf{e} \in \bar{L}$ as $k \rightarrow \infty$.

Then $\bar{m}^*(\mathbf{e}) = M^*(\bar{L})$. Indeed, $\bar{m}^*(\mathbf{e}) \geq M^*(\bar{L})$ and for the other direction, let $a > M^*(\bar{L})$. Then for all sufficiently large k we have $a \geq M^*(\eta_k, \bar{L})$, so we can conclude

$$a \geq M^*(\eta_k, \bar{L}) = \bar{m}^*(\eta_k, \mathbf{e}(\eta_k)) \geq \bar{m}^*(\mathbf{e}(\eta_k))$$

where we used $\bar{m}^*(\eta, \cdot) \geq \bar{m}^*(\cdot)$. Letting $k \rightarrow \infty$ we conclude, by the continuity of \bar{m}^* and by $\mathbf{e}(\eta_k) \rightarrow \mathbf{e}$, that $a \geq \bar{m}^*(\mathbf{e})$ and $M^*(\bar{L}) \geq \bar{m}^*(\mathbf{e})$ follows. The claim is proved.

Next we claim that \mathbf{e} is an equioscillation point. Indeed, assume for a contradiction that for some $j \in \{1, \dots, n\}$ we have $m_j^*(\mathbf{e}) < \bar{m}^*(\mathbf{e})$. Then there is $k_0 \in \mathbb{N}$ such that $m_j^*(\eta_{k_0}, \mathbf{e}) < \bar{m}^*(\mathbf{e})$. Since $m_j^*(\eta_{k_0}, \cdot)$ is continuous (the kernel functions are singular; see Proposition 4 (a)), there is $\delta > 0$ such that for every $\mathbf{y} \in \bar{L}$ with $\text{dist}_{\mathbb{T}^n}(\mathbf{y}, \mathbf{e}) < \delta$ one has $m_j^*(\eta_{k_0}, \mathbf{y}) < \bar{m}^*(\mathbf{e})$, too.

There is $n_0 \in \mathbb{N}$ such that for every $k \geq n_0$ we have $\text{dist}_{\mathbb{T}^n}(\mathbf{e}(\eta_k), \mathbf{e}) < \delta$. So for $k \geq \max\{k_0, n_0\}$ we can write

$$m_j^*(\eta_k, \mathbf{e}(\eta_k)) \leq m_j^*(\eta_{k_0}, \mathbf{e}(\eta_k)) < \bar{m}^*(\mathbf{e}) = M^*(\bar{L}) \leq \bar{m}^*(\mathbf{e}(\eta_k)) \leq \bar{m}^*(\eta_k, \mathbf{e}(\eta_k)).$$

This is a contradiction, since $m_i^*(\eta_k, \mathbf{e}(\eta_k)) = \bar{m}^*(\eta_k, \mathbf{e}(\eta_k))$ for each $i \in \{1, \dots, n\}$.

Therefore, \mathbf{e} is necessarily an equioscillation point. As such, it cannot have any degenerate subarcs, for any degenerate subarc $I_k^*(\mathbf{e}) = \{e_k\}$ would yield a singular value $m_k^*(\mathbf{y}) = -\infty$ according to the singularity of K . Hence $\mathbf{e} \in L$.

Choosing $\mathbf{w}^* := \mathbf{e}$ concludes the proof. \square

Now we extend Theorem 13 to general concave kernels, but this time we approximate the kernel from below.

Theorem 15. *Let $n \in \mathbb{N}$ and $\nu_1, \dots, \nu_n > 0$, let K be a singular periodic kernel function, and let J be an arbitrary periodic n -field function.*

Then there exists a maximin point \mathbf{z}^ on \bar{L} , it belongs to the open “cyclic simplex” L , and it is an equioscillation point.*

Proof. Let K be a singular, 1-periodic kernel function which is not necessarily strictly concave. Let $K^{(\eta)}(t) := K(t) + \eta(|\sin \pi t| - 1)$ where $\eta > 0$. Then $K^{(\eta)}$ is a strictly concave kernel function, which is also singular and 1-periodic and $K^{(\eta)} \nearrow K$ as $\eta \searrow 0$, moreover this convergence is uniform. Again, we denote the corresponding maximum functions and maximin quantity by $m_j^*(\eta, \mathbf{y})$, $\underline{m}^*(\eta, \mathbf{y})$ and $m^*(\eta, \bar{L})$. Due to the uniform convergence of $K^{(\eta)}$'s, we also have $\underline{m}^*(\eta, \mathbf{y}) \nearrow \underline{m}^*(\mathbf{y})$ uniformly, in the extended sense. Therefore $m^*(\eta, \bar{L}) \nearrow m^*(\bar{L})$, too.

Let $\mathbf{e}(\eta) \in \bar{L}$ be a node system such that $\underline{m}^*(\eta, \mathbf{e}(\eta)) = m^*(\eta, \bar{L})$. By Theorem 13, $\mathbf{e}(\eta) \in L$ and it is an equioscillating node system: $m_1^*(\eta, \mathbf{e}(\eta)) = \dots = m_n^*(\eta, \mathbf{e}(\eta))$. Since \bar{L} is compact, there exists $\eta_k \searrow 0$ such that $\mathbf{e}(\eta_k) \rightarrow \mathbf{e}$ for some $\mathbf{e} \in \bar{L}$ as $k \rightarrow \infty$.

Then $\underline{m}^*(\mathbf{e}) = m^*(\bar{L})$. Indeed, $\underline{m}^*(\mathbf{e}) \leq m^*(\bar{L})$ and for the other direction, let $b < m^*(\bar{L})$. Then for all sufficiently large k we have $b \leq m^*(\eta_k, \bar{L})$, so we can conclude

$$b \leq m^*(\eta_k, \bar{L}) = \underline{m}^*(\eta_k, \mathbf{e}(\eta_k)) \leq \underline{m}^*(\mathbf{e}(\eta_k))$$

where we used $\underline{m}^*(\eta, \cdot) \leq \underline{m}^*(\cdot)$. Letting $k \rightarrow \infty$ we conclude, by the extended continuity of \underline{m}^* and by $\mathbf{e}(\eta_k) \rightarrow \mathbf{e}$, that $b \leq \underline{m}^*(\mathbf{e})$ and $m^*(\bar{L}) \leq \underline{m}^*(\mathbf{e})$ follows. The claim is proved.

Next we claim that \mathbf{e} is an equioscillation point. Assume for a contradiction that for some $j \in \{1, \dots, n\}$ we have $m_j^*(\mathbf{e}) > \underline{m}^*(\mathbf{e})$. Then there is $k_0 \in \mathbb{N}$ such that $m_j^*(\eta_{k_0}, \mathbf{e}) > \underline{m}^*(\mathbf{e})$. Since $m_j^*(\eta_{k_0}, \cdot)$ is continuous (the kernel functions are singular; see Proposition 4 (a)), there is $\delta > 0$ such that for every $\mathbf{y} \in \bar{L}$ with $\text{dist}_{\mathbb{T}^n}(\mathbf{y}, \mathbf{e}) < \delta$ one has $m_j^*(\eta_{k_0}, \mathbf{y}) > \underline{m}^*(\mathbf{e})$, too.

There is $n_0 \in \mathbb{N}$ such that for every $k \geq n_0$ we have $\text{dist}_{\mathbb{T}^n}(\mathbf{e}(\eta_k), \mathbf{e}) < \delta$. So for $k \geq \max\{k_0, n_0\}$ we can write

$$m_j^*(\eta_k, \mathbf{e}(\eta_k)) \geq m_j^*(\eta_{k_0}, \mathbf{e}(\eta_k)) > \underline{m}^*(\mathbf{e}) = m^*(\bar{L}) \geq \underline{m}^*(\mathbf{e}(\eta_k)) \geq \underline{m}^*(\eta_k, \mathbf{e}(\eta_k)).$$

This is a contradiction, since $m_i^*(\eta_k, \mathbf{e}(\eta_k)) = \underline{m}^*(\eta_k, \mathbf{e}(\eta_k))$ for each $i \in \{1, \dots, n\}$.

Finally, if \mathbf{e} is an equioscillating node system and K is singular, then necessarily $\mathbf{e} \in L$, as before.

So we proved that there is a node system $\mathbf{z}^* := \mathbf{e}$, $\mathbf{z}^* \in L$ for which $\underline{m}^*(\mathbf{z}^*) = m^*(\bar{L})$ and \mathbf{z}^* is an equioscillating node system. \square

5.4 A counterexample for nonsingular kernels

Example 16. *If K is not a singular kernel, then there can be no equioscillation, and no maximin node systems. Moreover, all node systems can be solutions of the minimax problem.*

Let $K \equiv 0$, and $J(1/\ell) := 1 - 1/\ell$ ($\ell \in \mathbb{N}$), and $J(t) := 0$ otherwise. Let $n := 2$. Then there is no equioscillation. We consider the following cases. If $y_1 = y_2 = 0$, then $m_1^*(\mathbf{y}) = 0$ and $m_2^*(\mathbf{y}) = 1$. If $y_1 = 0$, $0 < y_2 < 1$, then $m_1^*(\mathbf{y}) = 1$ and $m_2^*(\mathbf{y}) = 1 - 1/\lfloor 1/y_2 \rfloor$. If $y_1 > 0$ and $y_2 < 1$, then $m_1^*(\mathbf{y}) = 1 - 1/\lfloor 1/y_1 \rfloor$ and $m_2^*(\mathbf{y}) = 1$. It is easy to see that there is no equioscillating node system. To verify the assertions about minimax node systems, note that $\overline{m}^*(\mathbf{y}) = \max(m_0^*(\mathbf{y}), m_1^*(\mathbf{y})) \equiv 1$ hence every node system is a node system with the minimax value. By considering node systems $\mathbf{y} = (0, 1/n)$, we get that $\underline{m}^*(\overline{L}) = 1$, but there is no node system with $\underline{m}^*(\mathbf{y}) = 1$.

6 A partial homeomorphism result

In our earlier papers [5, 8, 7] on the subject, an outstanding role was played by a new finding, not seen in earlier works of Bojanov [2] and Fenton [10]. We established, that in case of singular and strictly concave kernels a certain homeomorphism exists between admissible node systems and differences of the interval or arc maxima. Since the differences all being zero is equivalent to equioscillation, such a result immediately gives the existence and uniqueness of an equioscillating node system. This entails that proving e.g. that the minimax point is an equioscillating system can be strengthened to say that this equioscillation property *characterizes* the minimax node system. Similarly, if we further prove that a maximin node system is necessarily equioscillating, then it follows that the minimax equals to the maximin, and is attained at that unique point of equioscillation. Furthermore, from the homeomorphism result further consequences could be proved, most importantly about the intertwining of m_j , see [7].

Above we have proved that minimax and maximin node systems exist and that they are necessarily equioscillating node systems. Therefore it is most natural to try to complete the picture by a corresponding homeomorphism theorem. In fact, even for the torus setup such a homeomorphism theorem was already proved in [5], when there were no weights allowed, and where we assumed a few technical assumptions, too. In our present notation Corollary 9.3 of [5] runs as follows.

Theorem 17. *Suppose that for each $j = 0, \dots, n$ the kernel K_j belongs to $C^2(0, 1)$ with $K_j'' < 0$ and satisfies (∞) . Let $S := \{\mathbf{y} \in \mathbb{T}^n : 0 < y_1 < \dots < y_n < 1\}$ be the open simplex, while y_0 is understood as fixed at $y_0 = 0$.*

Then the difference mapping $\Phi(\mathbf{y}) := (m_1^(\mathbf{y}) - m_0^*(\mathbf{y}), \dots, m_n^*(\mathbf{y}) - m_{n-1}^*(\mathbf{y}))$ is a homeomorphism of S onto \mathbb{R}^n .*

Observe that here the $n + 1$ “to be translated” kernels admit a symmetry: if we rotate a node system $\mathbf{y} := (y_0, y_1, \dots, y_n) \in \mathbb{T}^{n+1}$ by say $t \in \mathbb{T}$, then for the new system $\mathbf{y}_t := (y_0 + t, y_1 + t, \dots, y_n + t) \in \mathbb{T}^{n+1}$ we get exactly the same vector of arc maxima $\mathbf{m}^*(\mathbf{y}_t) := (m_0^*(\mathbf{y}_t), m_1^*(\mathbf{y}_t), \dots, m_n^*(\mathbf{y}_t)) \in \mathbb{R}^{n+1}$, hence in particular also the differences of these maxima remain the same as before. Therefore it was natural to select one copy of these identical systems by fixing

the value of y_0 —also this was the only way the repetitions could be discarded and a homeomorphism could hold.

In our current settings, however, there is an outer field J , too. Once the weight, i.e., the field is not constant, we no longer have this rotational symmetry. The situation can be compared to the previous case regarding $K_0(\cdot - y_0)$ not as a kernel with fixed y_0 , but just as an outer field. Theorem 17 just says that we have a homeomorphism if the field is strictly concave and singular, and all the kernels satisfy the extra assumptions on differentiability etc. However, in this interpretation one thing constitutes a major difference: if we take only the n element node systems $(y_1, \dots, y_n) \in \mathbb{T}^n$ accompanied by a field (like e.g. $J := K_0(\cdot - y_0)$), then there will be only n arcs, determined by the nodes, so that the arc $I_n^*((y_1, \dots, y_n)) = [y_n, y_1]$ will be the union of the former two arcs $I_n^*(\mathbf{y}) = [y_n, y_0]$ and $I_0^*(\mathbf{y}) = [y_0, y_1]$. Similarly, the maximums will form an n -dimensional vector with $m_n^*(y_1, \dots, y_n)$ becoming the maximum of the former two maxima $m_n^*(\mathbf{y})$ and $m_0^*(\mathbf{y})$. The maximum differences then form an $n - 1$ -dimensional manifold, and we can no longer hope for a homeomorphism from the domain of our n -dimensional node systems to this manifold of differences.

To cure this, we may consider $K_0(\cdot - y_0)$ both a fixed kernel and also a field (say writing $\tilde{K}_0 := \frac{1}{2}K_0$ and $J := \frac{1}{2}K_0(\cdot - y_0)$). Then the result of Theorem 17 will refer to the original $n + 1$ arc maximums and their n differences, with a valid homeomorphism result.

With this in mind, we prove that an analogous “partial” homomorphism theorem remains in effect even if there is an arbitrary weight, i.e., field. In addition, we will surpass all the other technical conditions of Theorem 17 by the more advanced technology we have developed in [8], capable of handling even non-differentiable kernels. Instead of repeating the technical steps of that proof in the torus context, we directly reduce the statement to results of [8]. In fact there we have made a substantial effort to formulate and prove results which can potentially be used even in the periodic case—and that investment brings a profit here enabling us to refer back to them. Actually, we will use the following, proved for the case of the interval setup as Theorem 18 in [8].

Theorem 18. *Let K_1, \dots, K_n be strictly concave, singular kernel functions fulfilling condition (PM_0) and let J be a field function satisfying either $J(0) = \lim_{t \downarrow 0} J(t) = -\infty$ or $J(1) = \lim_{t \uparrow 1} J(t) = -\infty$ (or both).*

Then the difference function $\Phi(\mathbf{x}) := (m_1(\mathbf{x}) - m_0(\mathbf{x}), \dots, m_n(\mathbf{x}) - m_{n-1}(\mathbf{x}))$ is a locally bi-Lipschitz homeomorphism between $Y := \{(x_1, \dots, x_n) \in \overline{S}_{[0,1]} : m_k(\mathbf{x}) > -\infty \ (k = 0, 1, \dots, n)\} \subset [0, 1]^n$ and \mathbb{R}^n .

Note that here the ordering of nodes is fixed according to the simplex $\overline{S}_{[0,1]}$; and given the singularity condition, all degenerate node systems contain a degenerate arc with $m_k(\mathbf{x}) = -\infty$, so in fact the admissible set $Y \subset \overline{S}_{[0,1]}$, too. Also note that a non-admissible node system from $\overline{S}_{[0,1]} \setminus Y$ can never be an equioscillating node system, for $\overline{m}(\mathbf{x}) > -\infty$ excludes equioscillation at the $-\infty$ level. Thus in particular this entails existence and unicity of an equioscillating node system.

Theorem 19. Let $n \in \mathbb{N}$, let K_0, K_1, \dots, K_n be $n+1$ strictly concave 1-periodic kernel functions and let J be a 1-periodic, otherwise arbitrary $n+1$ -field function.

For any value $a \in \mathbb{T}$ denote $Y^* := Y^*(a) := \{\mathbf{y} = (y_0, y_1, \dots, y_n) \in \overline{L} \subset \mathbb{T}^{n+1} : y_0 = a, m_k(\mathbf{y}) > -\infty (k = 0, 1, \dots, n)\}$.

Then the difference function $\Phi^*(\mathbf{y}) := (m_1^*(\mathbf{y}) - m_0^*(\mathbf{y}), \dots, m_n^*(\mathbf{y}) - m_{n-1}^*(\mathbf{y}))$ is a locally bi-Lipschitz homeomorphism between $Y^*(a)$ and \mathbb{R}^n .

In particular, for each fixed value $y_0 = a$ there exists one unique equioscillating node system of the form $\mathbf{y}(a) = (a, y_1(a), \dots, y_n(a)) \in \overline{L}$.

Proof. First we introduce a new field function $J^* := J + K_0(\cdot - a)$. Obviously J^* will then be a 1-periodic upper bounded function which is nonsingular at more than n points mod 1 (while one finite value, if $J(a)$ was finite, could be “killed” by $K_0(\cdot - a)$). So, J^* is an n -field function, and in view of the singularity condition on K_0 it also satisfies the extra singularity equation $\lim_{t \rightarrow a} J^*(t) = -\infty$.

Interpreting the K_j and J^* as defined on the torus, we now transfer them to the interval $[0, 1]$ via the covering mapping π_a . We put $\tilde{J}(r) := J^*(\pi_a(r))$, which then becomes an n -field function on $[0, 1]$ satisfying $\lim_{r \downarrow 0} \tilde{J} = \lim_{r \uparrow 1} \tilde{J} = -\infty$. Also we put $\tilde{K}_j(r) := K_j(\pi_a(r) - a)$ for all $j = 1, \dots, n$. Obviously these are singular kernel functions, which are strictly concave, too.

We apply the above Theorem 18 to this new system. Let us see what are the arising node systems $\mathbf{x} \in Y$. First, $\mathbf{x} \in \overline{S}$ transfers to the cyclic ordering $a \preceq \pi_a(x_1) \preceq \dots \preceq \pi_a(x_n)$, so that writing $\mathbf{y} := (y_0, y_1, \dots, y_n)$ with $y_0 := a$ and $y_j := \pi_a(x_j)$ ($j = 1, \dots, n$), the ordering condition becomes $\mathbf{y} \in \overline{L}$.

Second, the non-singularity conditions for $\mathbf{x} \in Y$ translate to those for $\mathbf{y} \in Y^*(a)$, because $\pi_a : I_k(\mathbf{x}) \leftrightarrow I_k^*(\mathbf{y})$ and

$$\tilde{F}(\mathbf{x}, r) = \tilde{J}(r) + \sum_{k=1}^n \tilde{K}_k(r - x_k) = J(t) + K_0(t - a) + \sum_{k=1}^n K_j(t - y_k) \quad (t := \pi_a(r))$$

makes a one-to-one correspondence between sum of translates function values on corresponding points of any $I_k(\mathbf{x})$ resp. $I_k^*(\mathbf{y})$; in particular, we find $m_k(\mathbf{x}) = m_k^*(\mathbf{y})$, ($k = 0, \dots, n$), and an arc $I_k^*(\mathbf{y})$ will be singular if and only if the corresponding interval $I_k(\mathbf{x})$ was.

So we have seen that when \mathbf{y} runs $Y^*(a)$ then \mathbf{x} runs Y , and the correspondence is one-to-one. Moreover, degenerate points do not satisfy the nonsingularity condition, hence $Y \subset S_{[0,1]}$ and $Y^*(a) \subset L$. This means that between node systems \mathbf{x} and \mathbf{y} , the mapping π_a and even its inverse π_a^{-1} acts continuously, given that π_a^{-1} is applied only to y_1, \dots, y_n off a .

Summing up, $\Phi^*(\mathbf{y}) = \Phi(\pi_a^{-1}(\mathbf{y}))$, and even this composition mapping is continuous, moreover, it maps one-to-one to \mathbb{R}^n . Obviously its inverse $\Phi^{-1} \circ \pi_a$ is continuous, too, once Φ^{-1} was, so the mapping is a homeomorphism between $Y^*(a)$ and \mathbb{R}^n . We also get the bi-Lipschitz property from that of Φ .

Finally, uniqueness of an equioscillating node system with given fixed $y_0 = a$ follows from the homeomorphism result for $Y^*(a)$ (where we get exactly one system with all differences zero), and from the fact that outside $Y^*(a)$ all node

systems provide some singular value $m_k^*(\mathbf{y}) = -\infty$, while equioscillation cannot take place on that level, for $\overline{m}^*(\mathbf{y}) \in \mathbb{R}$, always. \square

Having proved the above, we may try to progress towards a Bojanov-type characterisation result. We already know that for the global minimax point there is equioscillation; and we now established that for each fixed value of y_0 there is exactly one equioscillating node system. So, writing $\varphi(a) = (a, y_1(a), \dots, y_n(a))$ for this unique equioscillation point with $y_0 = a$, it suffices to look for the global minimum of $\mu(a) := \overline{m}^*(\varphi(a))$ over $a \in \mathbb{T}$. The question is if these equioscillation values are always the same – as we have seen in [5] when $J \equiv 0$ – or if $\mu(a)$ is non-constant. Unicity of equioscillating *node systems* cannot hold (there is one for each fixed value a of the first coordinate y_0), but one may hope for unicity of the *equioscillation value*.

In Section 7, however, we will see that even this modified hope is deluded.

7 Counterexamples – equioscillation does not characterize minimax or maximin, majorization occurs, minimax can be smaller than maximin

Bojanov’s Theorem [2] in the classical algebraic polynomial setting included an important characterization statement, too: the extremal minimax system was *characterized* by the equioscillation property. That is, there was exactly one equioscillating node system, which was necessarily the minimax point.

Fenton’s classical theorem added another statement to the theory (in his context): he also proved that the unique maximin point equals to the (unique) minimax point (and hence the maximin and minimax values are equal, too).

In the weighted algebraic setting we proved similarly strong results in [7]. Also, we found the same for the unweighted trigonometric setting in [5], save the trivial free rotation (which was discarded by indexing from 0 and fixing the value of y_0 in \mathbb{T}).

In this section we explore examples which show that we cannot expect as many results as for the interval case or for the torus without weights. Basically, we will show that $m(L) > M(L)$ does occur.

Our examples will use a singular kernel, so by our above results minimax and maximin points exist, moreover, they are equioscillation points. Therefore, the examples also mean that there are different equioscillation values, hence strict majorization occurs even between equioscillating node systems. Furthermore, we will choose the kernel to be the standard log-sine kernel $K(t) := \log |\sin(\pi t)|$. This means that even for the classical trigonometric polynomial case one cannot expect any better results for general weights.

7.1 A counterexample with majorization

Example 20. Set $K(t) := \log |\sin(\pi t)|$, $n = 2$, $\nu_1 = \nu_2 = 1$, and $J(t) = 0$ on $\{0\} \cup [1/2, 1)$ and $J(t) = -\infty$ on $(0, 1/2)$.

Then, the minimax and maximin values on the “cyclic simplex” L are $m^*(\bar{L}) = -2 \log(2)$, $M^*(\bar{L}) = -\log(2)$, respectively.

Moreover, for any $\lambda \in [-2 \log(2), -\log(2)]$ there is an equioscillating node system $\mathbf{y} \in L$ with $\lambda = m_1^*(\mathbf{y}) = \dots = m_n^*(\mathbf{y})$.

In the following we will determine all equioscillation values.

We are to minimize $\bar{m}^*(\mathbf{y})$ (and maximize $\underline{m}^*(\mathbf{y})$) on $\mathbf{y} \in \mathbb{T}^2$. By relabeling, if necessary, we can assume $\mathbf{y} = (y_1, y_2)$ with $0 \leq y_1 \leq y_2 < 1$.

First, we make the following simple observations about the behavior of the pure sum of translates function. From the evenness of K it follows that $f(\mathbf{y}, \cdot)$ behaves symmetrically on the two intervals before and after the midpoint $\frac{y_1+y_2}{2}$ on $I_1^*(\mathbf{y}) \sim [y_1, y_2]$, and similarly for the other arc $I_2^*(\mathbf{y})$. In particular, $f(\mathbf{y}, \cdot)$ is strictly monotone increasing on $(y_1, \frac{y_1+y_2}{2})$ and decreasing on $(\frac{y_1+y_2}{2}, y_2)$. Similarly, it is strictly increasing on $(y_2, \frac{y_1+y_2}{2} + \frac{1}{2})$ and strictly decreasing on $(\frac{y_1+y_2}{2} + \frac{1}{2}, 1+y_1)$. Moreover, f is maximal on $I_i(\mathbf{y})$ at the midpoint, and if the length of this interval is $\ell := |I_i(\mathbf{y})|$, the length of the arc $I_i(\mathbf{y})$, then its value is $2K(\ell/2)$. As for $F(\mathbf{y}, \cdot)$, it follows that on any of the arcs $I_i^*(\mathbf{y})$ it attains its maximum on the point(s) of the arc which have $J(t) = 0$ and are closest to the midpoint among those. Also note that among the two midpoints, which are exactly of distance $1/2$, only one can belong to the singular set $X_J = (0, 1/2)$.

In the following we use the variables $x = y_1 + y_2$, $z = y_2 - y_1$, by which we can express $y_1 = \frac{x-z}{2}$, $y_2 = \frac{x+z}{2}$.

Note that we cannot have $0 \leq y_1 \leq y_2 \leq 1/2$, for then $m_1^*(\mathbf{y}) = -\infty$, which cannot be an equioscillation value, given that $\bar{m}^*(\mathbf{y}) > -\infty$. Thus $y_2 \geq 1/2$, and we have $x \geq y_2 \geq 1/2$, too.

So, let the first case be $1/2 \leq x < 1$. Then, the midpoint $x/2$ of the arc $I_1^*(\mathbf{y})$ lies in $[1/4, 1/2)$, which is in the singular set $(0, 1/2)$, hence the maximum will be attained at the closest nonsingular point of the arc, which is $1/2$. That is, $m_1^*(\mathbf{y}) = F(\mathbf{y}, 1/2) = f(\mathbf{y}, 1/2)$. Further, $m_2^*(\mathbf{y}) = F(\mathbf{y}, x/2 + 1/2) = f(\mathbf{y}, x/2 + 1/2)$.

We will use the identity

$$\begin{aligned} f(\mathbf{y}, t) &= f\left(\left(\frac{x-z}{2}, \frac{x+z}{2}\right), t\right) = \log \left| \sin \pi \left(t - \frac{x-z}{2}\right) \sin \pi \left(t - \frac{x+z}{2}\right) \right| \\ &= -\log(2) + \log |\cos(\pi z) - \cos \pi(2t - x)| \quad (t \in \mathbb{R}). \end{aligned} \quad (33)$$

With this, the equation $f(\mathbf{y}, 1/2) = f(\mathbf{y}, x/2 + 1/2)$ can be rewritten as

$$\left| \cos(\pi z) - \cos \pi \left(2\frac{1}{2} - x\right) \right| = \left| \cos(\pi z) - \cos \pi \left(2\left(\frac{x}{2} + \frac{1}{2}\right) - x\right) \right|.$$

The right hand side is $1 + \cos(\pi z)$. For the sign of the left hand side we observe $x + z = 2y_2 \geq 1$, hence $1 - x \leq z \leq x \leq 1$, so by monotonicity of $\cos(\pi s)$ for $0 \leq s \leq 1$, we get $|\cos(\pi z) - \cos \pi(2\frac{1}{2} - x)| = \cos(\pi(1 - x)) - \cos(\pi z)$.

So we are led to the equation

$$\cos \pi(1 - x) - \cos(\pi z) = \cos(\pi z) + 1,$$

and solving it for $\cos(\pi z)$ yields

$$\cos(\pi z) = \frac{\cos \pi(1-x) - 1}{2}.$$

The condition $1-x \leq z$ is satisfied, since $\cos \pi(1-x) \geq (\cos \pi(1-x) - 1)/2$ in general. By simple steps, the condition $z \leq x$ is equivalent to $\cos(\pi x) \leq -1/3$. So if $1/2 \leq x \leq \beta_0 := \arccos(-1/3)/\pi \approx 0.608$, then z does not satisfy $z \leq x$.

The arising equioscillation value is

$$\begin{aligned} m_2^* \left(\left(\frac{x-z}{2}, \frac{x+z}{2} \right) \right) &= f(\mathbf{y}, x/2 + 1/2) \\ &= -\log(2) + \log \left| \frac{\cos \pi(1-x) - 1}{2} + 1 \right| \\ &= -2\log(2) + \log(1 - \cos(\pi x)) \end{aligned}$$

for $\beta_0 \leq x \leq 1$.

The next case is $1 \leq x < 3/2$. Since $x+z = y_1+y_2 \leq 2$, we have $0 \leq z \leq 2-x$, too. Now $x/2 \notin X_J$, while $x/2+1/2 \in X_J$ (except for $x=1$ and $x/2+1/2=1$), with the closest nonsingular point from $I_2^*(\mathbf{y})$ being 1 (remaining valid even in case $x=1$). Hence, $m_1^*(\mathbf{y}) = f(\mathbf{y}, x/2)$ and $m_2^*(\mathbf{y}) = f(\mathbf{y}, 1)$. Now, again by (33), $f(\mathbf{y}, x/2) = f(\mathbf{y}, 1)$ is equivalent to

$$\left| \cos(\pi z) - \cos \pi \left(2\frac{x}{2} - x \right) \right| = |\cos(\pi z) - \cos \pi(2-x)|. \quad (34)$$

Here the left hand side is $|\cos(\pi z) - 1| = 1 - \cos(\pi z)$. Also, $0 \leq z \leq 2-x \leq 1$, so $\cos(\pi z) \geq \cos \pi(2-x)$, and (34) can be written as

$$1 - \cos(\pi z) = \cos(\pi z) - \cos \pi(2-x).$$

Solving it for $\cos(\pi z)$ we are led to

$$\cos(\pi z) = \frac{1 + \cos \pi x}{2}.$$

The condition $0 \leq z \leq 2-x \leq 1$ on z is equivalent to $1 \geq \cos(\pi z) \geq \cos \pi(2-x) = \cos(\pi x)$ and hence is obviously satisfied.

The equioscillation value, again depending only on x , is found again to be

$$\begin{aligned} m_2^* \left(\left(\frac{x-z}{2}, \frac{x+z}{2} \right) \right) &= f(\mathbf{y}, x/2) \\ &= -\log(2) + \log \left| 1 - \frac{1 + \cos(\pi x)}{2} \right| \\ &= -2\log(2) + \log(1 - \cos(\pi x)). \end{aligned}$$

Finally, let $3/2 \leq x < 2$. This means that $y_1 \geq 1/2$, for $x = y_1 + y_2 \leq y_1 + 1$. Then again, $m_1^*(\mathbf{y}) = f(\mathbf{y}, x/2)$, for $x/2 \in [1/2, 1]$. However, in $I_2^*(\mathbf{y})$ the closest

non-singular point to the midpoint $x/2+1/2 \equiv x/2-1/2 \pmod{1}$ will be $1/2$, and we will get $m_2^*(\mathbf{y}) = f(\mathbf{y}, 1/2)$. Therefore, the equioscillation equation becomes $f(\mathbf{y}, x/2) = f(\mathbf{y}, 1/2)$ and, by (33), it is

$$|\cos(\pi z) - \cos \pi \left(2\frac{x}{2} - x\right)| = |\cos(\pi z) - \cos \pi \left(2\frac{1}{2} - x\right)|.$$

The left hand is $1 - \cos(\pi z)$ and the right hand side is $|\cos(\pi z) - \cos \pi x|$. Since $3/2 \leq x < 2$ and $z + x = 2y_2 \leq 2$ implies $0 \leq z \leq 2 - x < 1/2$, here both terms are nonnegative, and the equation becomes

$$1 - \cos(\pi z) = \cos(\pi z) + \cos \pi x.$$

Solving it for $\cos(\pi z)$ we get

$$\cos(\pi z) = \frac{1 - \cos(\pi x)}{2}.$$

Again, we verify that $0 \leq z \leq 2 - x$, which is equivalent to $1 \geq \cos(\pi z) \geq \cos \pi(2 - x)$. The second inequality is equivalent to $\cos(\pi x) \leq 1/3$, and it holds for $x \in [3/2, 2]$ if and only if $x \leq 2 - \arccos(1/3)/\pi = 1 + \beta_0$, $1 + \beta_0 \approx 1.608$. So $z \leq 2 - x$ if and only if $x \in [3/2, 1 + \beta_0]$.

The equioscillation value is

$$\begin{aligned} m_2^* \left(\left(\frac{x-z}{2}, \frac{x+z}{2} \right) \right) &= f(\mathbf{y}, 1) \\ &= -\log(2) + \log \left| 1 - \frac{1 - \cos(\pi x)}{2} \right| \\ &= -2\log(2) + \log(1 + \cos(\pi x)). \end{aligned}$$

Now we can collect the obtained equioscillation values coming from all the three cases:

$$m_1^*(\mathbf{y}) = m_2^*(\mathbf{y}) = -2\log 2 + \begin{cases} \log(1 - \cos(\pi x)) & (\beta_0 \leq x < 3/2) \\ \log(1 + \cos(\pi x)) & (3/2 \leq x < 1 + \beta_0) \end{cases}$$

It is minimal when $x = 3/2$ (in this case $z = 1/3$) and it is maximal when $x = 1$ (in this case $z = 1/2$). The corresponding values are $-2\log(2) \approx -1.386$ and $-\log(2) \approx -0.693$.

Hence we get $M(\bar{L}) = -2\log(2)$ and $m(\bar{L}) = -\log(2)$.

7.2 A modified counterexample with continuous field function

In this subsection we sketch a counterexample which is a modification of the previous one, but with a continuous field function.

First, we set a new external field function \tilde{J} , and then we compute the arc maxima for the two extremal node systems from the previous counterexample. We will find that the arc maxima $m_i^*(\mathbf{y})$ of those two node systems will not change when we replace \tilde{J} for J , whence they will still be equioccillating with the same equioscillation values $-\log 2$ and $-2\log 2$, respectively.

Let $\alpha > 4\pi$ be fixed and let \tilde{J} be 0 on $[1/2, 1]$, $-\alpha t$ on $[0, 1/4]$ and $\alpha(t - 1/2)$ on $[1/4, 1/2]$; further, let \tilde{J} be extended 1-periodically to \mathbb{R} .

Regarding the $x = 1, z = 1/2$ maximin configuration, we have $y_1 = 1/4, y_2 = 3/4$. It is easy to check that $f(\mathbf{y}, t)$ is strictly monotone decreasing on $(0, 1/4]$ and on $[1/2, 3/4]$ and it is strictly monotone increasing on $(1/4, 1/2]$ and on $(3/4, 1]$. Hence, $\tilde{F}(\mathbf{y}, t) = \tilde{J}(t) + f(\mathbf{y}, t)$ is the same monotone in these intervals. Therefore $\tilde{m}_1^*(\mathbf{y}) = \tilde{F}(\mathbf{y}, 1/2) = F(\mathbf{y}, 1/2) = f(\mathbf{y}, 1/2) = -\log(2)$ and $\tilde{m}_2^*(\mathbf{y}) = \tilde{F}(\mathbf{y}, 1) = F(\mathbf{y}, 1) = f(\mathbf{y}, 1) = -\log(2)$.

Regarding the $x = 3/2, z = 1/3$ minimax configuration, we have $y_1 = 7/12, y_2 = 11/12$. It is easy to check that $f(\mathbf{y}, t)$ is strictly monotone increasing (and concave) on $(-1/12, 1/2]$ and $f'(\mathbf{y}, 0) = 4\pi$. Adding \tilde{J} to it, we see that $\tilde{F}(\mathbf{y}, \cdot)$ is strictly monotone increasing on $(-1/12, 0]$ and strictly monotone decreasing on $[0, 1/4]$. Using the symmetry of $\tilde{F}(\mathbf{y}, \cdot)$ with respect to $1/4$, $\tilde{F}(\mathbf{y}, \cdot)$ is strictly monotone increasing on $(1/4, 1/2]$ and strictly monotone decreasing on $[1/2, 7/12]$. Moreover, $\tilde{m}_1^*(\mathbf{y}) = \tilde{F}(\mathbf{y}, 0) = \tilde{F}(\mathbf{y}, 1/2) = f(\mathbf{y}, 0) = -2\log(2)$. Computing $\tilde{m}_2^*(\mathbf{y})$ is simpler: for $t \in I_2^*(\mathbf{y}) = (7/12, 11/12) \subset [1/2, 1]$, $\tilde{F}(\mathbf{y}, t) = f(\mathbf{y}, t) = F(\mathbf{y}, t)$, and $\sup f(\mathbf{y}, \cdot)$ remains the same as before. Therefore, $\tilde{m}_2^*(\mathbf{y}) = \tilde{F}(\mathbf{y}, 3/4) = f(\mathbf{y}, 3/4) = -2\log(2)$.

Summing up, we obtained that $\tilde{m}(\bar{L}) \leq -2\log(2)$ and $\tilde{M}(\bar{L}) \geq -\log(2)$. Also, taking into account that $\tilde{J} \geq J$, hence $\tilde{F} \geq F$, we also know that $\tilde{m}(\bar{L}) \geq m(\bar{L})$ and $\tilde{M}(\bar{L}) \geq M(\bar{L})$. Therefore, $\tilde{M}(\bar{L}) = -2\log(2)$, too.

Even if we don't proceed to compute the exact maximin value, too, it is clear from what has already been done that in this example $\tilde{m}(\bar{L}) \geq -\log(2) > \tilde{M}(\bar{L}) = -2\log 2$, hence the same phenomenon takes place for the continuous, finite kernel \tilde{J} as before for the kernel J .

Acknowledgment

This research was supported by project TKP2021-NVA-09. Project no. TKP2021-NVA-09 has been implemented with the support provided by the Ministry of Innovation and Technology of Hungary from the National Research, Development and Innovation Fund, financed under the TKP2021-NVA funding scheme.

The work of Sz. Gy. Révész was supported in part by Hungarian National Research, Development and Innovation Fund project # K-119528.

References

- [1] C. Berge, *Topological spaces*. Translated from the French original by E. M. Patterson. Reprint of the 1963 translation Dover Publications, Inc., Mineola, NY, 1997. xiv+270 pp.
- [2] Bojanov, B. D. *A generalization of Chebyshev polynomials*. J. Approx. Theory **26** (1979), no.4, 293–300.
- [3] P. Borwein, and T. Erdélyi, *Polynomials and polynomial inequalities*. Grad. Texts in Math., 161 Springer-Verlag, New York, 1995. x+480 pp. ISBN:0-387-94509-1
- [4] B. Farkas and B. Nagy, *Transfinite diameter, Chebyshev constant and energy on locally compact spaces*, Potential Anal. **28** (2008), no. 3, 241–260.
- [5] B. Farkas, B. Nagy, and Sz. Gy. Révész, *A minimax problem for sums of translates on the torus*, Trans. London Math. Soc. **5** (2018), no. 1, 1–46.
- [6] B. Farkas, B. Nagy, and Sz. Gy. Révész, *On intertwining of maxima of sum of translates functions with nonsingular kernels*, Tr. Inst. Mat. Mekh. **28** (2022), no. 4, 262–272.
- [7] B. Farkas, B. Nagy, and Sz. Gy. Révész, *On the weighted Bojanov-Chebyshev problem and the sum of translates method of Fenton*, (in Russian), Sbornik Math. **214** (2023), no. 8, 119–150.
- [8] B. Farkas, B. Nagy, and Sz. Gy. Révész, *A homeomorphism theorem for sums of translates*, Rev. Mat. Complut. pp. 49, DOI: 10.1007/s13163-023-00461-6.
- [9] B. Farkas, B. Nagy, and Sz. Gy. Révész, *Fenton type minimax problems for sum of translates functions*, pp. 27, arXiv:2210.04348.
- [10] P. C. Fenton, *A min-max theorem for sums of translates of a function*, J. Math. Anal. Appl. **244** (2000), no. 1, 214–222.
- [11] P. C. Fenton, *$\cos \pi \lambda$ again*, Proc. Amer. Math. Soc. **131** (2003), no. 6, 1875–1880.
- [12] R. A. Rankin, *On the closest packing of spheres in n dimensions*, Ann. of Math. (2) **48** (1947), 1062–1081.
- [13] E. B. Saff and V. Totik, *Logarithmic potentials with external fields*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 316, Springer-Verlag, Berlin, 1997, Appendix B by Thomas Bloom.
- [14] Tatiana Nikiforova, *On the weighted Bojanov-Chebyshev problem on an infinite interval*, manuscript.

Béla Nagy
Department of Analysis,
Bolyai Institute, University of Szeged
Aradi vértanúk tere 1
6720 Szeged, Hungary

Szilárd Gy. Révész
Alfréd Rényi Institute of Mathematics
Reáltanoda utca 13-15
1053 Budapest, Hungary