

# On the Extremal Functions of Acyclic Forbidden 0–1 Matrices\*

Seth Pettie  
University of Michigan

Gábor Tardos  
HUN-REN Rényi Institute

## Abstract

The extremal theory of forbidden 0–1 matrices studies the asymptotic growth of the function  $\text{Ex}(P, n)$ , which is the maximum weight of a matrix  $A \in \{0, 1\}^{n \times n}$  whose submatrices avoid a fixed pattern  $P \in \{0, 1\}^{k \times l}$ . This theory has been wildly successful at resolving problems in combinatorics [Kla00, MT04, CK12], discrete and computational geometry [Für90, Agg15, ES96, PS91, Mit92, BG91], structural graph theory [GM14, BGK<sup>+</sup>21, BKTW22] and the analysis of data structures [Pet10, KS20], particularly corollaries of the dynamic optimality conjecture [CGK<sup>+</sup>15b, CGK<sup>+</sup>15a, CGJ<sup>+</sup>23, CPY24].

All these applications use *acyclic* patterns, meaning that when  $P$  is regarded as the adjacency matrix of a bipartite graph, the graph is acyclic. The biggest open problem in this area is to bound  $\text{Ex}(P, n)$  for acyclic  $P$ . Prior results [Pet11a, PS13] have only ruled out the strict  $O(n \log n)$  bound conjectured by Füredi and Hajnal [FH92]. At the two extremes, it is *consistent* with prior results that  $\forall P. \text{Ex}(P, n) \leq n \log^{1+o(1)} n$ , and also consistent that  $\forall \epsilon > 0. \exists P. \text{Ex}(P, n) \geq n^{2-\epsilon}$ .

In this paper we establish a stronger lower bound on the extremal functions of acyclic  $P$ . Specifically, for any  $t \geq 1$  we give a new construction of relatively dense 0–1 matrices with  $\Theta(n(\log n / \log \log n)^t)$  1s that avoid a certain acyclic pattern  $X_t$ . Pach and Tardos [PT06] have conjectured that this type of result is the best possible, i.e., no acyclic  $P$  exists for which  $\text{Ex}(P, n) \geq n(\log n)^{\omega(1)}$ .

## 1 Introduction

The theory of forbidden 0–1 matrices subsumes or generalizes many problems in extremal combinatorics, such as Davenport-Schinzel sequences [HS86, ASS89, Niv10, Pet15a, WP18] and their generalizations [Pet11b, Pet15b, FH92], Zarankiewicz’s problem [KST54], and bipartite Turán-type subgraph avoidance. Forbidden 0–1 matrices have been applied to problems in discrete and computational geometry, the amortized analysis of data structures, and in other areas of extremal combinatorics. Some highlights in geometry include bounding the number of unit-distances in a convex point set [Für90, Agg15], the number of critical placements of an  $n$ -gon in a hippodrome [ES96], an analysis of the Bentley-Ottman line sweeping algorithm [PS91], and an analysis of Mitchel’s algorithm for obstacle-avoiding shortest paths in the plane [Mit92, BG91]. In data structures, forbidden 0–1 matrices have been used to analyze data structures based on binary search trees and path-compression [Pet10], and more recently, to several corollaries of Sleator and Tarjan’s [ST85] *dynamic optimality* conjecture [CGK<sup>+</sup>15b, CGK<sup>+</sup>15a, CGJ<sup>+</sup>23, KS20, CPY24]. The most well-known application of forbidden 0–1 matrices is probably Marcus and Tardos’s proof [MT04] of the Stanley-Wilf conjecture, via Klazar’s reduction [Kla00] to a Füredi-Hajnal conjecture [FH92]. They have also been used to bound Stanley-Wilf limits [Cib09, Fox13, CK17], and to bound the size of sets of permutations with some fixed VC-dimension [CK12]. Most recently, results on searching for forbidden patterns [GM14] inspired the definition of *twin-width* for graphs and other binary structures [BKTW22, BGK<sup>+</sup>21].

If  $P \in \{0, 1\}^{k \times l}$  and  $A \in \{0, 1\}^{n \times n}$ , we say  $A$  *contains*  $P$ , written  $P < A$ , if there are rows  $r_1 < \dots < r_k$  and columns  $c_1 < \dots < c_l$  such that  $P(i, j) = 1 \rightarrow A(r_i, c_j) = 1$ . If  $P \not< A$  then  $A$  *avoids*  $P$  or is  $P$ -free. The basic extremal function is defined as follows

$$\text{Ex}(P, n) = \max\{\|A\|_1 \mid A \in \{0, 1\}^{n \times n} \text{ and } P \not< A\},$$

\*S. Pettie is supported by NSF Grant CCF-2221980. G. Tardos is supported by the National Research, Development and Innovation Office projects K-132696 and SNN-135643 and by the ERC Advanced Grants “ERMiD” and “GeoScape.” This research was performed while S. Pettie was visiting Rényi Institute, Budapest.

and may be generalized in various ways, e.g., to avoid *sets* of forbidden patterns [Tar05] or rectangular  $A$  [FH92, Pet11b], or to  $d$ -dimensional matrices/patterns [KM06, MP17, Gen19]. Observe that  $P$  and  $A$  can be viewed as incidence matrices of bipartite graphs, where the two vertex sets are implicitly *ordered*; let  $G(P)$  be the *unordered* undirected graph corresponding to  $P$ .

Füredi and Hajnal [FH92] attempted to systematically classify small forbidden patterns by their extremal functions, and managed to do this for most weight-4 patterns,<sup>1</sup> with the last holdouts being classified by Tardos [Tar05]. For any pattern  $P$ , the extremal function  $\text{Ex}(P, n)$  is at least as large as the unordered (Turán) extremal function for  $G(P)$ . A natural question is to determine when they are the same, asymptotically, and the maximum factor by which they can differ.

Füredi and Hajnal [FH92] concluded their article with several influential conjectures. They first conjectured that when  $P$  is a  $k \times k$  permutation matrix, that  $\text{Ex}(P, n) \leq c_k n$ , i.e., it is asymptotically the same as the Turán number of  $G(P)$ . Klazar [Kla00] proved that this conjecture implies the Stanley-Wilf conjecture, and Marcus and Tardos [MT04] proved both conjectures. See [Gen09, Cib09, CK17, Fox13, KM06] for generalizations and sharper analyses of the leading constants. Füredi and Hajnal next conjectured that  $\text{Ex}(P, n)$  would never be more than a  $\log n$ -factor larger than the (unordered) Turán number of  $G(P)$ . Perhaps doubting this conjecture, they immediately asked whether it held for *acyclic* patterns  $P$ , i.e., if  $G(P)$  is a forest, is  $\text{Ex}(P, n) = O(n \log n)$ ?

Pach and Tardos [PT06] refuted the second Füredi-Hajnal conjecture, by exhibiting arbitrarily large  $P$  for which  $G(P) = C_{2k}$  is a  $2k$ -cycle but  $\text{Ex}(P, n) = \Omega(n^{4/3})$ . This implies the gap between the ordered and unordered extremal functions is  $n^{1/3-\epsilon}$ , where  $\epsilon = 1/k$  can be made arbitrarily small. However, this refutation did not imply anything about the gap for *acyclic* matrices. Understanding acyclic patterns is important, as *every* application to geometry, data structures, and combinatorics mentioned in the first paragraph uses only acyclic patterns. Pettie [Pet11a] disproved the last Füredi-Hajnal conjecture by exhibiting a specific acyclic pattern  $X$  for which  $\text{Ex}(X, n) = \Omega(n \log n \log \log n)$ . An unpublished manuscript of Park and Shi [PS13] extended this lower bound to a set of patterns  $\{X_m\}$  for which  $\text{Ex}(X_m, n) = \Omega(n \log n \log \log n \log \log \log n \cdots \log^{(m)} n)$ .

The constructions of [Pet11a, PS13] refuted the *letter* of Füredi and Hajnal's conjecture, but certainly not its *spirit*. Consider several non-trivial possibilities for the extremal function of an acyclic  $P$ .

**Absolute Polylog( $n$ ).** There is an absolute constant  $c \geq 1$  such that for any acyclic  $P$ ,  $\text{Ex}(P, n) \leq n \log^{c+o(1)} n$ .

**Variable Polylog( $n$ ).** For any acyclic  $P$ , there is a constant  $c = c(P)$  such that  $\text{Ex}(P, n) \leq n \log^c n$ .

**Subpolynomial.** For every acyclic  $P$ , there is some  $\epsilon(n) = o(1)$  depending on  $P$  such that  $\text{Ex}(P, n) \leq n^{1+\epsilon(n)}$ .

**Polynomial.** For some  $c < 2$ , every acyclic  $P$  has  $\text{Ex}(P, n) \leq O(n^c)$ .

None of these upper bounds have been established or ruled out. In particular, prior work [Pet11a, PS13] does not preclude the possibility that **Absolute Polylog( $n$ )** holds even with  $c = 1$ , and it is also possible the **Polynomial** fails, i.e., for every  $\epsilon > 0$ , there exists an acyclic  $P$  for which  $\text{Ex}(P, n) = \Omega(n^{2-\epsilon})$ . Pach and Tardos [PT06] conjectured broadly that the **Variable Polylog( $n$ )** upper bound is true, and conjectured more specifically that  $\text{Ex}(P, n) = O(n \log^{\|P\|_1 - 3} n)$ .

The biggest open problem in the theory of forbidden 0–1 matrices is to understand *acyclic patterns*. On the upper bound side, we have a perfect classification of all patterns with four 1s [FH92, Tar05], and a good classification for those with five 1s [PT06], up to a  $\log n$  factor. For example,  $\text{Ex}(R_1, n)$  and  $\text{Ex}(R_2, n)$  are known to be  $\Omega(n \log n)$  and  $O(n \log^2 n)$  [PT06].

$$R_1 = \begin{pmatrix} \bullet & \bullet & & \\ & & \bullet & \\ \bullet & & & \bullet \end{pmatrix} \qquad R_2 = \begin{pmatrix} \bullet & \bullet & & \bullet \\ \bullet & & \bullet & \\ & & & \bullet \end{pmatrix}$$


Korándi, Tardos, Tomon, and Weidert [KTTW19] defined a pattern  $P$  to be *class- $s$  degenerate* if it can be written  $P = \begin{pmatrix} P' \\ P'' \end{pmatrix}$ , where at most one column has a non-zero intersection with both  $P'$  and  $P''$ , and  $P', P''$  are class- $(s-1)$  degenerate. (In the diagrams of  $S_1, S_2$ , a valid row partition cuts at most one edge.) Any  $P$  with a single

<sup>1</sup>The *weight* of  $A$  is  $\|A\|_1$ , i.e., the number of 1s in  $A$ .


row is class-0 degenerate. They proved that every class- $s$  degenerate  $P$  has

$$\text{Ex}(P, n) \leq n \cdot 2^{O(\log^{1-\frac{1}{s+1}} n)} = n^{1+o(1)}.$$


For example,  $\text{Ex}(S_1, n), \text{Ex}(S_2, n) \leq n \cdot 2^{O(\log^{2/3} n)}$  as  $S_1, S_2$  are class-2 degenerate.



Clearly, a pattern and its transpose have the same extremal function. The smallest non-degenerate acyclic pattern whose transpose is also non-degenerate is the “pretzel”  $T$ ; we have no non-trivial upper bounds on  $\text{Ex}(T, n)$ .



**1.1 New Result** Our main result is a *proper* refutation of the Füredi-Hajnal conjecture for acyclic matrices that rules out the **Absolute Polylog**( $n$ ) world. For any  $t \geq 2$ , we give a new construction of 0–1 matrices containing  $\Omega(n(\log n/\log \log n)^t)$  1s, and prove that they avoid a particular  $2t \times (2t + 1)$  acyclic pattern  $X_t$  with  $4t$  1s.



**THEOREM 1.1.** *For every  $t \geq 2$ , there exists a  $2t \times (2t + 1)$  acyclic pattern  $X_t$  such that*

$$\text{Ex}(X_t, n) = \begin{cases} \Omega(n(\log n/\log \log n)^t), \\ O(n \log^{4t-3} n). \end{cases}$$

**1.2 Organization** Section 2 presents the construction of 0–1 matrices and some of its properties unrelated to forbidden substructures. Section 3 analyzes the forbidden substructures, culminating in a proof of Theorem 1.1. We conclude in Section 4 with a concise survey of open problems in 0–1 matrices and related problems in *ordered graphs*.

**2 A Construction of 0–1 Matrices**

For any positive integer  $k$ , let  $[k] = \{1, \dots, k\}$ . Fix a constant integer  $t \geq 2$ . The rows and columns of  $A_t$  are indexed by length- $(tk)$  strings in the set

$$\mathcal{I} = [k^t]^{tk}.$$

An element of  $\mathcal{I}$  is partitioned into  $t$  *blocks*, each of length  $k$ . If  $a \in \mathcal{I}$ , let  $a(p) \in [k^t]^k$  be its  $p$ th block, and  $a(p, q) \in [k^t]$  be the  $q$ th coordinate in block  $p$ . We use angular brackets to denote any injective mapping from  $[k]^r$  to  $[k^r]$ , e.g.,  $\langle j_1, j_2, j_3 \rangle = (j_1 - 1)k^2 + (j_2 - 1)k + j_3$ . For  $(j_1, \dots, j_t) \in [k]^t$ , define  $\mathbf{v} = \mathbf{v}[j_1, \dots, j_t] \in \mathcal{I}$  to be

the vector that is 0 in all coordinates except:

$$\begin{aligned} \mathbf{v}(1, j_1) &= \langle \cdot \rangle = 1, \\ \mathbf{v}(2, j_2) &= \langle j_1 \rangle = j_1, \\ &\dots \\ \mathbf{v}(r, j_r) &= \langle j_1, \dots, j_{r-1} \rangle \\ &\dots \\ \mathbf{v}(t, j_t) &= \langle j_1, \dots, j_{t-1} \rangle. \end{aligned}$$

Define  $\mathcal{S}$  to be the set of eligible vectors,

$$\mathcal{S} = \{\mathbf{v}[j_1, \dots, j_t] \mid (j_1, \dots, j_t) \in [k]^t\}.$$

Letting  $n = |\mathcal{I}|$ ,  $A_t$  is an  $n \times n$  0-1 matrix whose row and column sets are both indexed by  $\mathcal{I}$ , ordered lexicographically. It is defined as follows:

$$A_t(a, b) = \begin{cases} 1 & \text{if } b - a \in \mathcal{S}, \\ 0 & \text{otherwise.} \end{cases}$$

LEMMA 2.1.  $\|A_t\|_1 = \Theta(n(\log n / \log \log n)^t)$ .

*Proof.*  $A_t$  is an  $n \times n$  0-1 matrix, where  $n = k^{t^2k} = |\mathcal{I}|$ . Pick a uniformly random row  $a \in \mathcal{I}$ , and a uniformly random vector  $\mathbf{v} = \mathbf{v}[i_1, \dots, i_t] \in \mathcal{S}$ . The probability that  $a + \mathbf{v} \in \mathcal{I}$  is legal is the probability that for all  $r \in [t]$ ,  $a(r, i_r) + \langle i_1, \dots, i_{r-1} \rangle \leq k^t$ , which is at least  $1 - k^{-t+(r-1)}$  since  $\langle i_1, \dots, i_{r-1} \rangle \leq k^{r-1}$ . We have

$$\Pr(a + \mathbf{v} \in \mathcal{I}) \geq \prod_{r=1}^t (1 - k^{-t+(r-1)}) \geq 1 - \sum_{r=1}^t k^{-r} > 1 - (k-1)^{-1}.$$

Therefore the number of 1s in  $A_t$  is at least  $(1 - (k-1)^{-1})nk^t = \Theta(n(\log n / \log \log n)^t)$ .  $\square$

### 3 Forbidden Substructures

If  $a, b \in \mathcal{I}$  are distinct vectors, their *type* is the first block where they differ, i.e.,

$$\text{type}(a, b) = \min\{r \mid a(r) \neq b(r)\}.$$

LEMMA 3.1. 1. For  $a < b < c$ ,  $a, b, c \in \mathcal{I}$ ,  $\text{type}(a, c) \leq \text{type}(b, c)$ .

2. Suppose  $A_t(a, c) = A_t(b, c) = 1$ , with  $a < b$ . Let  $c - a = \mathbf{v}[i_1, \dots, i_t]$  and  $c - b = \mathbf{v}[j_1, \dots, j_t]$ . If  $\text{type}(a, b) = r$ , then  $i_q = j_q$  for  $q < r$ ,  $i_r < j_r$ , and the first coordinate where  $a$  and  $b$  differ is  $(r, i_r)$ .

$$a \begin{pmatrix} c \\ \bullet \\ \bullet \end{pmatrix}$$

3. Suppose  $A_t(a, c) = A_t(a, d) = 1$ , with  $c < d$ . Let  $c - a = \mathbf{v}[i_1, \dots, i_t]$  and  $d - a = \mathbf{v}[j_1, \dots, j_t]$ . If  $\text{type}(c, d) = r$ , then  $i_q = j_q$  for  $q < r$ ,  $i_r > j_r$ , and the first coordinate where  $c$  and  $d$  differ is  $(r, j_r)$ .

$$a \begin{pmatrix} c & d \\ \bullet & \bullet \end{pmatrix}$$

4. Suppose  $A_t(b, c_1) = A_t(a, c_2) = A_t(b, d) = A_t(a, d) = 1$ , where  $a < b$  and  $c_1 < c_2 < d$ . Then it is not possible that  $\text{type}(a, b) = \text{type}(c_1, d) = \text{type}(c_2, d)$ .

$$a \begin{pmatrix} c_1 & c_2 & d \\ \bullet & \bullet & \bullet \end{pmatrix}$$

5. Suppose  $A_t(a, c_0) = A_t(b, c_1) = A_t(a, c_2) = A_t(b, d) = A_t(a, d) = 1$ , where  $a < b$  and  $c_0 < c_1 < c_2 < d$ . If  $\text{type}(a, b) \leq \text{type}(c_0, d)$ , then  $\text{type}(c_0, d) < \text{type}(c_2, d)$ .

$$\begin{matrix} a \\ b \end{matrix} \begin{pmatrix} c_0 & c_1 & c_2 & d \\ \bullet & & \bullet & \bullet \\ & \bullet & & \bullet \\ & & & \bullet \end{pmatrix}$$

*Proof.* Part 1 holds because  $\mathcal{I}$  is ordered lexicographically.

Part 2. Let  $s$  be the first index where  $i_s \neq j_s$ . Clearly, the first two coordinates where  $\mathbf{v}[i_1, \dots, i_t]$  and  $\mathbf{v}[j_1, \dots, j_t]$  differ is  $(s, i_s)$  and  $(s, j_s)$ . This makes  $a = c - \mathbf{v}[i_1, \dots, i_t]$  and  $b = c - \mathbf{v}[j_1, \dots, j_t]$  also differ first at the same two coordinates, making  $\text{type}(a, b) = s$ , so we have  $r = s$ . At coordinate  $(s, j_s)$ ,  $\mathbf{v}[j_1, \dots, j_t]$  is  $\langle j_1, \dots, j_{s-1} \rangle > 0$  but  $\mathbf{v}[i_1, \dots, i_t]$  is zero there, so we have  $a(s, j_s) > b(s, j_s)$ . As  $a < b$ ,  $(s, j_s)$  cannot be the first coordinate where  $a$  and  $b$  differ, so we conclude that  $i_s < j_s$  and  $(s, i_s) = (r, i_r)$  is the coordinate of first difference between  $a$  and  $b$ .

Part 3 can be proved the same way as Part 2.

Part 4. Suppose, for the purpose of obtaining a contradiction, that  $\text{type}(a, b) = \text{type}(c_1, d) = \text{type}(c_2, d) = r$ . Both vectors  $d - a$  and  $d - b$  are in  $\mathcal{S}$ , so let

$$d - a = \mathbf{v}[i_1, \dots, i_t],$$

$$d - b = \mathbf{v}[j_1, \dots, j_t].$$

Applying Part 3 to row  $b$ , we know  $c_1$  and  $d$  first differ at coordinate  $(r, j_r)$ , and applying Part 3 to row  $a$ , we know  $c_2$  and  $d$  first differ at coordinate  $(r, i_r)$ . Applying Part 2 to column  $d$ , we have  $i_r < j_r$ . However, since  $c_2 < d$  we have

$$c_1(r, i_r) = d(r, i_r) > c_2(r, i_r),$$

implying  $c_1 > c_2$ , contradicting the originally defined order  $c_1 < c_2$ .

Part 5. We have  $\text{type}(c_0, d) \leq \text{type}(c_2, d)$  by Part 1. Suppose, for the purpose of obtaining a contradiction, that  $\text{type}(c_0, d) = \text{type}(c_2, d) = r$ , which implies, again by Part 1, that  $\text{type}(c_1, d) = r$  as well. We have assumed  $\text{type}(a, b) \leq r$  and  $\text{type}(a, b) = r$  would contradict Part 4, so we have  $\text{type}(a, b) < r$ . With the notation introduced in Part 4, both pairs  $(c_0, d)$  and  $(c_2, d)$  first differ at coordinate  $(r, i_r)$ , while  $(c_1, d)$  first differ at coordinate  $(r, j_r)$ . We have  $c_0 < c_1 < c_2 < d$  lexicographically, so we must also have  $i_r = j_r$ . We further know that

$$\begin{aligned} d(r, i_r) - c_0(r, i_r) &= d(r, i_r) - c_2(r, i_r) = \mathbf{v}[i_1, \dots, i_t](r, i_r) = \langle i_1, \dots, i_{r-1} \rangle, \\ d(r, j_r) - c_1(r, j_r) &= \mathbf{v}[j_1, \dots, j_t](r, j_r) = \langle j_1, \dots, j_{r-1} \rangle. \end{aligned}$$

This implies

$$c_0(r, i_r) = c_2(r, i_r) \neq c_1(r, i_r)$$

since Part 2 and  $\text{type}(a, b) < r$  implies that  $\langle i_1, \dots, i_{r-1} \rangle \neq \langle j_1, \dots, j_{r-1} \rangle$ . Now  $(r, i_r)$  is the first coordinate where  $c_1$  differs from either  $c_0$  or  $c_2$ , but  $c_0$  and  $c_2$  agree on this coordinate, which contradicts the ordering  $c_0 < c_1 < c_2$ . This confirms Part 5 of the lemma.  $\square$

Define the alternating patterns  $P_t$  and  $Q_t$ , where  $Q_t$  is a reflection of  $P_t$  across the minor diagonal.

$$P_t = \begin{pmatrix} \bullet & & & & & & \\ & \bullet & & & & & \\ & & \bullet & & & & \\ & & & \cdots & & & \\ & & & & \bullet & & \\ & & & & & \bullet & \\ & & & & & & \bullet \end{pmatrix} \quad Q_t = \begin{pmatrix} \bullet & \bullet \\ & \bullet \\ & \vdots \\ & \bullet \\ & \bullet \end{pmatrix}$$

When  $t \geq 2$ ,  $P_t$  and  $Q_t$  appear in  $A_t$ , and in fact  $P_{t'}, Q_{t'}$  appear in  $A_t$  for every constant  $t' \geq t$ . Lemmas 3.2 and 3.3 give useful constraints on how  $P_t, Q_t$  can be embedded in  $A_t$ .

LEMMA 3.2. Consider an occurrence of  $P_t$  in  $A_t$ , where  $a, b \in \mathcal{I}$  are the indices of the two rows and  $c, d \in \mathcal{I}$  are the indices of the first and last columns.

$$\begin{matrix} a \\ b \end{matrix} \begin{pmatrix} c & & & & d \\ \bullet & \bullet & \cdots & \bullet & \bullet \\ & \bullet & & \bullet & \bullet \end{pmatrix}$$

If  $\text{type}(a, b) \leq \text{type}(c, d)$ , then  $\text{type}(a, b) = \text{type}(c, d) = 1$ .

*Proof.* For  $i \in [t]$ , let  $c_i \in \mathcal{I}$  be the index of the  $(2i - 1)^{\text{th}}$  column in this occurrence of  $A_t$ , so  $c = c_1 < c_2 < \cdots < c_t < d$ . We have  $\text{type}(a, b) \leq \text{type}(c_1, d) \leq \cdots \leq \text{type}(c_t, d)$ , where the first inequality is assumed and rest follow from Lemma 3.1(1). Part 5 of Lemma 3.1 applies and implies that these latter inequalities are *strict*, i.e.,

$$\text{type}(a, b) \leq \text{type}(c_1, d) < \text{type}(c_2, d) < \cdots < \text{type}(c_t, d).$$

All types are from  $[t]$ , therefore both  $\text{type}(a, b)$  and  $\text{type}(c, d) = \text{type}(c_1, d)$  must be 1.  $\square$

LEMMA 3.3. Consider an occurrence of  $Q_t$  in  $A_t$ , where  $a, b \in \mathcal{I}$  are the indices of the first and last rows and  $c, d \in \mathcal{I}$  are the indices of the two columns. If  $\text{type}(a, b) \geq \text{type}(c, d)$ , then  $\text{type}(a, b) = \text{type}(c, d) = 1$ .

*Proof.* First one has to establish the analogues of Lemma 3.1(4,5) for patterns that are reflected across the minor diagonal, depicted below, then follow the proof of Lemma 3.2.

$$\begin{pmatrix} \bullet & \bullet \\ & \bullet \\ \bullet & \end{pmatrix} \quad \begin{pmatrix} \bullet & \bullet \\ & \bullet \\ \bullet & \bullet \end{pmatrix}$$

For both parts, the original proofs work *mutatis mutandis*.  $\square$

Define  $X_t$  to be the following  $2t \times (2t + 1)$  pattern.

$$X_t = \begin{pmatrix} \bullet & \bullet & \cdots & \bullet & \bullet & \bullet \\ & \bullet & & & & \bullet \\ & & & & & \bullet \\ & \bullet & & & & \vdots \\ & \vdots & & & & \bullet \\ & \bullet & & & & \bullet \\ \bullet & \bullet & \cdots & \bullet & \bullet & \bullet \end{pmatrix}$$

Alternatively,  $X_t$  is defined to be the 0–1 matrix whose first and last rows with the first column removed form  $P_t$ , while its second and last column form  $Q_t$  and has a single 1 entry outside these submatrices in the first column and last row.

LEMMA 3.4.  $A_t$  avoids  $X_t$ .

*Proof.* Suppose there is an occurrence of  $X_t$  in  $A_t$ . Let  $a, b \in \mathcal{I}$  be the indices of the first and last rows of the  $X_t$  instance, and let  $c', c, d \in \mathcal{I}$  be the indices of its first, second and last columns. We either have  $\text{type}(a, b) \leq \text{type}(c, d)$  or  $\text{type}(a, b) \geq \text{type}(c, d)$ . We must have  $\text{type}(a, b) = \text{type}(c, d) = 1$  in both cases by Lemmas 3.2 and 3.3, respectively. But then  $\text{type}(c', d)$  is also 1 by Lemma 3.1(1) and then the rows  $a, b$  and columns  $c', c, d$  contradict Lemma 3.1(4). The contradiction proves our lemma.  $\square$

Lemma 3.4 lets us obtain a lower bound on  $\text{Ex}(X_t, n)$ . We use a Lemma of Pach and Tardos [PT06] to get a nearly matching upper bound.



problems [AKN<sup>+</sup>08]. A pattern is *light* if it contains exactly one 1 per column. All light patterns are in  $\mathcal{P}_{\text{qlin}}$  [Kla92, Niv10, Pet15b] and all patterns known to be in  $\mathcal{P}_{\text{qlin}} \setminus \mathcal{P}_{\text{lin}}$  are either light, or composed of light or linear patterns via Keszegh's [Kes09] joining operation.<sup>3</sup> It is an open question whether there are infinitely many *minimal* non-linear patterns in  $\mathcal{P}_{\text{qlin}}$ . It is consistent with known results that if  $P$  is light, then  $P \in \mathcal{P}_{\text{lin}}$  iff it avoids the following patterns (or their reflections), which correspond to order-3 Davenport-Schinzel sequences.

$$\begin{pmatrix} \cdot & & \\ & \cdot & \cdot \\ & & \cdot \end{pmatrix} \quad \begin{pmatrix} \cdot & \cdot & \cdot \\ & \cdot & \cdot \\ & & \cdot \end{pmatrix}$$

It would also be of interest to characterize, for each  $t \geq 1$ , (light) patterns  $P \in \mathcal{P}_{\text{qlin}}$  for which  $\text{Ex}(P, n) \geq n^{2\Omega(\alpha^t(n))}$ .

**Acyclic Patterns.** All acyclic  $P$  for which the bound  $\text{Ex}(P, n) \leq n(\log n)^{O(1)}$  is known can be proved via the Pach-Tardos reductions [PT06, Lemmas 2, 3, and 4] (see Lemma 3.5 for one), together with Keszegh's [Kes09] joining operation.<sup>4</sup> If  $P$  is *degenerate* then  $\text{Ex}(P, n) \leq n^{1+o(1)}$  [KTTW19]. Upper bounding  $\text{Ex}(P, n)$  by  $n(\log n)^{O_P(1)}$ ,  $n^{1+o(1)}$ , or even  $n^{2-\epsilon}$  for *all* acyclic patterns  $P$  is the main open problem in this area. A more subtle problem is to determine which extremal functions are possible. Tardos [Tar05] gave examples of *pairs* of patterns with  $\text{Ex}(\{P, P'\}, n) = \Theta(n \log \log n)$  and  $\text{Ex}(\{P'', P'''\}, n) = \Theta(n \log n / \log \log n)$ , but it is unknown whether these extremal functions can be achieved by a *single* forbidden pattern.

**Arbitrary Patterns.** The Kővári-Sós-Turán theorem [KST54] implies that if  $P \in \{0, 1\}^{k \times l}$ , then  $\text{Ex}(P, n) = O\left(n^{2 - \frac{1}{\min\{k, l\}}}\right)$ . Pach and Tardos [PT06] constructed  $\Theta(n^{4/3})$ -weight matrices that avoid some arbitrarily long ordered cycles. See Timmons [Tim12] and Győri et al. [GKM<sup>+</sup>18] for more results on ordered cycles. Methuku and Tomon [MT22] defined a matrix  $P$  to be *row  $t$ -partite* if it can be cut along rows into  $t$  light matrices, and  *$t \times t$ -partite* if both  $P$  and  $P^T$  are row  $t$ -partite. They proved that if  $P$  is row  $t$ -partite and  $t \times t$ -partite, that  $\text{Ex}(P, n)$  is at most  $n^{2-1/t+1/t^2+o(1)}$  and  $n^{2-1/t+o(1)}$ , respectively.

A 0–1 matrix can be viewed as an *ordered bipartite* graph, where the two parts of the bipartition are given independent linear orders. Forbidden 0–1 matrix theory has been extended to other types of ordered subgraph containment.

**Vertex-ordered graphs** It is natural to drop the requirement that the forbidden graph and host graph be bipartite and just consider the extremal theory of arbitrary *vertex-ordered graphs*: these are simple graphs with a linear order on their vertices. Containment between vertex-ordered graphs must preserve the ordering. The extremal function of such a (forbidden) graph  $H$  was introduced in [PT06]:  $\text{Ex}(H, n)$  is the maximum number of edges of an  $n$ -vertex vertex-ordered graph that does not contain  $H$ . The connection to the extremal theory is 0–1 patterns is very close. A vertex-ordered graphs  $H$  is called *ordered bipartite* (or of interval chromatic number 2) if it is a bipartite graph with one partite class of vertices preceding the other in the vertex-order. If  $H$  is not ordered bipartite, then  $\text{Ex}(H, n) = \Theta(n^2)$ . If  $H$  is ordered bipartite,

<sup>3</sup>Keszegh [Kes09] observed that if  $A$  has a 1 in its southeast corner and  $B$  has a 1 in its northwestern corner, that by *joining* them at their corners, the resulting pattern  $A \oplus B$  has extremal function  $\text{Ex}(A \oplus B, n) \leq \text{Ex}(A, n) + \text{Ex}(B, n)$ .

$$A \oplus B = \begin{pmatrix} \boxed{\begin{matrix} & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \end{matrix}} & \begin{matrix} \\ \\ \bullet \\ \\ \end{matrix} \\ \begin{matrix} \\ \\ \\ \\ \end{matrix} & \boxed{\begin{matrix} & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \end{matrix}} \end{pmatrix}$$

If  $A$  is light, and  $B$  is the transpose of a light pattern, then  $A \oplus B \in \mathcal{P}_{\text{qlin}}$ , but neither it nor its transpose is light.  
<sup>4</sup>For example, the following pattern has extremal function  $O(n \log^3 n)$ , but it is not subject to any of the Pach-Tardos reductions. It must first be decomposed into two patterns via Keszegh [Kes09], each of which has extremal function  $O(n \log^3 n)$  by [PT06].

$$\begin{pmatrix} \bullet & & & & \\ \bullet & \bullet & & & \\ & \bullet & \bullet & & \\ & & \bullet & & \\ & & & \bullet & \\ & & & & \bullet \end{pmatrix}$$



let  $P(H)$  be its bipartite 0–1 adjacency matrix (ordering the rows and columns consistent with the given vertex-order) and we have

$$\text{Ex}(P(H), n/2) \leq \text{Ex}(H, n) = O(\text{Ex}(P(H), n) \cdot \log n).$$

This implies that the vertex-ordered graph  $H$  for which  $P(H) = X_t$  is an example of an ordered bipartite tree whose extremal function is  $\Omega(n(\log n/\log \log n)^t)$ . Previously no ordered bipartite tree was known whose extremal function was not  $n \log^{1+o(1)} n$ .

Although the connection between the 0–1 matrices and vertex-ordered graphs is not close enough to directly translate questions about the *linearity* of extremal functions, the situation was similar in the two theories: although characterization of forbidden vertex-ordered graphs  $H$  with  $\text{Ex}(H, n) = O(n)$  is currently beyond reach, the paper [FKMV20] provides such a characterization for *connected*  $H$ .

**Edge-ordered graphs** Rather than extend Turán-type extremal graph theory by adding a total order on *vertices*, we could instead add a total order on *edges*. This yields the extremal theory of *edge-ordered graphs* as introduced by [GMN<sup>+</sup>23]. A rich theory starts to form but it is not as closely related to the extremal theory of 0–1 patterns as the vertex-ordered variant is. Nevertheless, many results and problems have analogues in the two theories. The analogue of the interval chromatic number is the *order chromatic number*: an edge-ordered graph has order chromatic number 2 if it is contained in the lexicographically ordered complete bipartite graph. The extremal function of an edge-ordered graph is  $\Theta(n^2)$  if and only if its order chromatic number is not 2. While the characterization of edge-ordered graphs with linear extremal functions seems to also be beyond reach in general, such a characterization is given for connected edge-ordered graphs [KT23b]. The extremal function of acyclic edge-ordered graphs of order-chromatic number 2 was conjectured to be  $n^{1+o(1)}$  in [GMN<sup>+</sup>23] and this has recently been established in [KT23a], where the stronger (and still open) conjecture was formulated that these extremal functions are all of the form  $O(n \log^c n)$ , where  $c$  may depend on the forbidden edge-ordered graph. In contrast to the main result of this paper, it is still possible that the above bound holds with  $c = 1$  for *all* acyclic edge-ordered graphs of order chromatic number 2.

## References

- [Agg15] Amol Aggarwal. On unit distances in a convex polygon. *Discret. Math.*, 338(3):88–92, 2015.
- [AKN<sup>+</sup>08] Noga Alon, Haim Kaplan, Gabriel Nivasch, Micha Sharir, and Shachar Smorodinsky. Weak  $\epsilon$ -nets and interval chains. *J. ACM*, 55(6), 2008.
- [ASS89] Pankaj K. Agarwal, Micha Sharir, and Peter W. Shor. Sharp upper and lower bounds on the length of general Davenport-Schinzel sequences. *J. Combin. Theory Ser. A*, 52:228–274, 1989.
- [BG91] Daniel Bienstock and Ervin Györi. An extremal problem on sparse 0–1 matrices. *SIAM J. Discr. Math.*, 4(1):17–27, 1991.
- [BGK<sup>+</sup>21] Édouard Bonnet, Colin Geniet, Eun Jung Kim, Stéphan Thomassé, and Rémi Watrigant. Twin-width II: small classes. In *Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1977–1996, 2021.
- [BKTW22] Édouard Bonnet, Eun Jung Kim, Stéphan Thomassé, and Rémi Watrigant. Twin-width I: tractable FO model checking. *J. ACM*, 69(1):3:1–3:46, 2022.
- [CGJ<sup>+</sup>23] Parinya Chalermsook, Manoj Gupta, Wanchote Jiamjitrak, Nidia Obscura Acosta, Akash Pareek, and Sorrachai Yingchareonthawornchai. Improved pattern-avoidance bounds for greedy BSTs via matrix decomposition. In *Proceedings of the 34th ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 509–534, 2023.
- [CGK<sup>+</sup>15a] Parinya Chalermsook, Mayank Goswami, László Kozma, Kurt Mehlhorn, and Thatchaphol Saranurak. Greedy is an almost optimal deque. In *Proceedings 14th Symposium on Algorithms and Data Structures*, volume 9214 of *Lecture Notes in Computer Science*, pages 152–165, 2015.
- [CGK<sup>+</sup>15b] Parinya Chalermsook, Mayank Goswami, László Kozma, Kurt Mehlhorn, and Thatchaphol Saranurak. Pattern-avoiding access in binary search trees. In *Proceedings of the 56th IEEE Symposium on Foundations of Computer Science (FOCS)*, pages 410–423, 2015.
- [Cib09] Josef Cibulka. On constants in the Füredi-Hajnal and the Stanley-Wilf conjecture. *J. Comb. Theory, Ser. A*, 116(2):290–302, 2009.
- [CK12] Josef Cibulka and Jan Kynčl. Tight bounds on the maximum size of a set of permutations with bounded VC-dimension. *J. Combin. Theory Ser. A*, 119(7):1461–1478, 2012.

- [CK17] Josef Cibulka and Jan Kynčl. Better upper bounds on the Füredi-Hajnal limits of permutations. In *Proceedings of the 28th ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 2280–2293, 2017.
- [CPY24] Parinya Chalermsook, Seth Pettie, and Sorrachai Yingchareonthawornchai. Sorting pattern-avoiding permutations via 0–1 matrices forbidding product patterns. In *Proceedings 35th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, 2024.
- [ES96] Alon Efrat and Micha Sharir. A near-linear algorithm for the planar segment-center problem. *Discrete & Computational Geometry*, 16(3):239–257, 1996.
- [FH92] Zoltán Füredi and Péter Hajnal. Davenport-Schinzel theory of matrices. *Discrete Mathematics*, 103(3):233–251, 1992.
- [FKMV20] Zoltán Füredi, Alexandr V. Kostochka, Dhruv Mubayi, and Jacques Verstraëte. Ordered and convex geometric trees with linear extremal function. *Discret. Comput. Geom.*, 64(2):324–338, 2020.
- [Fox13] Jacob Fox. Stanley-Wilf limits are typically exponential. *CoRR*, abs/1310.8378, 2013.
- [Ful09] Radoslav Fulek. Linear bound on extremal functions of some forbidden patterns in 0–1 matrices. *Discrete Mathematics*, 309:1736–1739, 2009.
- [Für90] Zoltán Füredi. The maximum number of unit distances in a convex  $n$ -gon. *J. Combin. Theory Ser. A*, 55(2):316–320, 1990.
- [Gen09] Jesse Geneson. Extremal functions of forbidden double permutation matrices. *J. Combin. Theory Ser. A*, 116(7):1235–1244, 2009.
- [Gen19] Jesse Geneson. Forbidden formations in multidimensional 0–1 matrices. *Eur. J. Comb.*, 78:147–154, 2019.
- [GKM<sup>+</sup>18] Ervin Györi, Dániel Korándi, Abhishek Methuku, István Tomon, Casey Tompkins, and Máté Vizer. On the Turán number of some ordered even cycles. *Eur. J. Comb.*, 73:81–88, 2018.
- [GM14] Sylvain Guillemot and Dániel Marx. Finding small patterns in permutations in linear time. In *Proceedings 25th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 82–101, 2014.
- [GMN<sup>+</sup>23] Dániel Gerbner, Abhishek Methuku, Dániel T. Nagy, Dömötör Pálvölgyi, Gábor Tardos, and Máté Vizer. Turán problems for edge-ordered graphs. *J. Comb. Theory, Ser. B*, 160:66–113, 2023.
- [HS86] Sergiu Hart and Micha Sharir. Nonlinearity of Davenport-Schinzel sequences and of generalized path compression schemes. *Combinatorica*, 6(2):151–177, 1986.
- [Kes09] Balázs Keszegh. On linear forbidden submatrices. *J. Combin. Theory Ser. A*, 116(1):232–241, 2009.
- [Kla92] Martin Klazar. A general upper bound in extremal theory of sequences. *Comment. Math. Univ. Carolin.*, 33(4):737–746, 1992.
- [Kla00] Martin Klazar. The Füredi-Hajnal conjecture implies the Stanley-Wilf conjecture. In *Formal power series and algebraic combinatorics (Moscow, 2000)*, pages 250–255. Springer, Berlin, 2000.
- [KM06] Martin Klazar and Adam Marcus. Extensions of the linear bound in the Füredi-Hajnal conjecture. *Adv. in Appl. Math.*, 38(2):258–266, 2006.
- [KS20] László Kozma and Thatchaphol Saranurak. Smooth heaps and a dual view of self-adjusting data structures. *SIAM J. Comput.*, 49(5), 2020.
- [KST54] Tamás Kövari, Vera T. Sós, and Pál Turán. On a problem of K. Zarankiewicz. *Colloquium Math.*, 3:50–57, 1954.
- [KT23a] Gaurav Kucheriya and Gábor Tardos. A characterization of edge-ordered graphs with almost linear extremal functions. *CoRR*, abs/2206.12979, 2023. *Combinatorica*, to appear.
- [KT23b] Gaurav Kucheriya and Gábor Tardos. On edge-ordered graphs with linear extremal functions. In *Proceedings of EuroComb23*, 2023.
- [KTTW19] Dániel Korándi, Gábor Tardos, István Tomon, and Craig Weidert. On the Turán number of ordered forests. *J. Comb. Theory, Ser. A*, 165:32–43, 2019.
- [KV94] Martin Klazar and Pavel Valtr. Generalized Davenport-Schinzel sequences. *Combinatorica*, 14(4):463–476, 1994.
- [Mit92] Joseph S. B. Mitchell.  $L_1$  shortest paths among polygonal obstacles in the plane. *Algorithmica*, 8(1):55–88, 1992.
- [MP17] Abhishek Methuku and Dömötör Pálvölgyi. Forbidden hypermatrices imply general bounds on induced forbidden subposet problems. *Comb. Probab. Comput.*, 26(4):593–602, 2017.
- [MT04] Adam Marcus and Gábor Tardos. Excluded permutation matrices and the Stanley-Wilf conjecture. *J. Combin. Theory Ser. A*, 107(1):153–160, 2004.
- [MT22] Abhishek Methuku and István Tomon. Bipartite Turán problems for ordered graphs. *Comb.*, 42(6):895–911, 2022.
- [Niv10] Gabriel Nivasch. Improved bounds and new techniques for Davenport-Schinzel sequences and their generalizations. *J. ACM*, 57(3), 2010.
- [Pet10] Seth Pettie. Applications of forbidden 0–1 matrices to search tree- and path compression-based data structures. In *Proceedings 21st ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1457–1467, 2010.
- [Pet11a] Seth Pettie. Degrees of nonlinearity in forbidden 0–1 matrix problems. *Discrete Mathematics*, 311:2396–2410, 2011.
- [Pet11b] Seth Pettie. Generalized Davenport-Schinzel sequences and their 0–1 matrix counterparts. *J. Comb. Theory Ser. A*, 118(6):1863–1895, 2011.

- [Pet11c] Seth Pettie. On the structure and composition of forbidden sequences, with geometric applications. In *Proceedings 27th Annual Symposium on Computational Geometry (SoCG)*, pages 370–379, 2011.
- [Pet15a] Seth Pettie. Sharp bounds on Davenport-Schinzel sequences of every order. *J. ACM*, 62(5):36, 2015.
- [Pet15b] Seth Pettie. Three generalizations of Davenport-Schinzel sequences. *SIAM J. Discrete Mathematics*, 29(4):2189–2238, 2015.
- [PS91] János Pach and Micha Sharir. On vertical visibility in arrangements of segments and the queue size in the Bentley-Ottmann line sweeping algorithm. *SIAM J. Comput.*, 20(3):460–470, 1991.
- [PS13] Sung Gi Park and Qinru Shi. New bounds on extremal numbers in acyclic ordered graphs. <https://math.mit.edu/news/summer/SPURprojects/2013ParkShi.pdf>, 2013.
- [PT06] János Pach and Gábor Tardos. Forbidden paths and cycles in ordered graphs and matrices. *Israel J. Math.*, 155:359–380, 2006.
- [ST85] Daniel D. Sleator and Robert E. Tarjan. Self-adjusting binary search trees. *J. ACM*, 32(3):652–686, 1985.
- [Tar05] Gábor Tardos. On 0–1 matrices and small excluded submatrices. *J. Combin. Theory Ser. A*, 111(2):266–288, 2005.
- [Tim12] Craig Timmons. An ordered turán problem for bipartite graphs. *Electron. J. Comb.*, 19(4):43, 2012.
- [WP18] Julian Wellman and Seth Pettie. Lower bounds on Davenport-Schinzel sequences via rectangular Zarankiewicz matrices. *Discret. Math.*, 341(7):1987–1993, 2018.