

## *Polybrominated diphenyl ethers' exposure causes changes in the physiology of zebra mussel, *Dreissena polymorpha* (Pallas, 1771)*

Vesela Yancheva<sup>1</sup>, Iliana Velcheva<sup>1</sup>, Elenka Georgieva<sup>2</sup>, Krisztán Nyeste<sup>3,4</sup>,  
László Antal<sup>3,4</sup>, Borislava Todorova<sup>1</sup>, Stela Stoyanova<sup>2\*</sup>

<sup>1</sup> - University of Plovdiv, Faculty of Biology, Department of Ecology and Environmental Conservation, Tzar Assen Str. 24, Plovdiv, 4000, BULGARIA

<sup>2</sup> - University of Plovdiv, Faculty of Biology, Department of Developmental Biology, Tzar Assen Str. 24, Plovdiv, 4000, BULGARIA

<sup>3</sup> - University of Debrecen, Department of Hydrobiology, Egyetem Sqr. 1, Debrecen, 4032, HUNGARY

<sup>4</sup> - University of Debrecen, National Laboratory for Water Science and Water Security, Egyetem Sqr. 1, Debrecen, 4032, HUNGARY

\*Corresponding author: stela.stoyanova@uni-plovdiv.bg

**Abstract.** The present laboratory study (96 hours and 30 days) aimed to provide for the first time the possible adverse effects of different concentrations of PBDEs congeners (PBDE 28, PBDE 47, PBDE 99, PBDE 100, PBDE 153, PBDE 154) based on Water Framework Directive 2000/60/EC (WFD) in zebra mussel (*Dreissena polymorpha* Pallas, 1771). Therefore, we analysed the gill histochemical structure by applying the Periodic acid–Schiff staining method to assess the exposed individuals' physiological status and linked them to the tested PBDEs concentrations in the contaminated water.

**Key words:** mussels, pollution, organic contaminants, PBDEs, biomarkers.

### Introduction

Persistent organic pollutants (POPs) are well-known compounds that are toxic to humans and wildlife because they are lipophilic, have low degradation rates, biomagnify through the food chain, and can travel long distances far from the releasing use source (Ahrend et al., 2023). POPs of most significant concern fall into several categories, including polychlorinated biphenyls (PCBs), organochlorine pesticides (OCPs), perfluorinated chemicals (PFCs), and polybrominated diphenyl ethers (PBDEs) (Medina, 2024). According to Yang et al. (2023), PBDEs are a class of typical POPs with 209 congeners.

Three types of commercial PBDE mixtures, namely penta-, octa, and deca-BDE, have been ubiquitously used as flame retardants in various industrial products, such as plastic products, electronics, building materials, and textiles (Portet-Koltalo et al., 2021). However, the widespread usage of PBDE-containing products, combined with their long-distance migration, high lipophilic ability, and accumulation in fat tissues, has resulted in global environmental problems (Abbasi et al., 2019). For instance, among the PBDE congeners, tetrabromodiphenyl ether (PBDE 47) is the dominant congener found in the environment, and numerous studies have already indica-

ted that PBDEs are endocrine-disrupting chemicals in both, animal and human bodies (Al-Harbi et al., 2021; Li et al., 2024).

Mussels are widely used as bioindicators and probably the most commonly used organisms in field or laboratory pollution experiments and for aquatic monitoring along with fish because they are filter-feeders and sessile, they have a wide geographical distribution and availability throughout the year and due to their efficiency in accumulating POPs, and tolerance to various environmental conditions (Yancheva et al., 2018a; Gecheva et al., 2020; Georgieva et al., 2022). The freshwater zebra mussel is suggested as an equivalent of *Mytilus* spp., which is notably used in large-scale programs, such as the Mussel Watch (Binelli et al., 2015; Beyer et al., 2017). This type of monitoring was first applied in a marine environment using blue mussels (Goldberg, 1975), but later in freshwaters with zebra mussels (Secor et al., 1993; Faria et al., 2010; Lepom et al., 2012; Louis et al., 2020; Hani et al., 2021; Baratange et al., 2023). However, the zebra mussel is a wide-spread invasive bivalve species commonly found in freshwaters of the northern hemisphere and identified by the IUCN as one of the 100 of the World's worst invasive alien species, but it has also been used actively as a model organism for freshwater toxicology since the late 1970s (Stoyanova et al., 2020; Yancheva et al., 2016, 2017a,b, 2018b, 2019a, 2020; 2021, 2022). In this regard, the zebra mussel is also widespread in most European freshwaters, including Bulgaria. In Bulgaria, this species was reported initially in the Danube River, the lowest reaches of its tributaries, some Black Sea lagoons, and the Black Sea coastal lakes and rivers (Valkanov, 1957; Russev et al., 1994; Angelov, 2000; Hubenov, 2005), including the Veleka River.

This study aimed to evaluate the zebra mussels' health status after acute (96 hours) and subchronic (30 days) exposure to different PBDEs concentrations under laboratory conditions. To achieve this, we applied the histochemical biomarker approach.

## **Materials and Methods**

### ***Experimental setup***

The experimental setup was conducted by national and international guidelines of

the European Parliament and the Council on protecting animals used for scientific purposes (Directive 2010/63/EU).

### ***Mussel collection***

About 100 zebra mussels were hand-collected and transported on the same day in polyethylene containers, thoroughly cleaned with distilled water, and then filled with water from the water source. The mussels were placed in a 50 L aquarium, pre-filled with dechlorinated water, and equipped with oxygen pumps in the vivarium of the Faculty of Biology at the University of Plovdiv to acclimatize for two weeks.

### ***PBDEs' exposure***

Acute (96 h) and subchronic (30 days) test was carried out. The zebra mussels (n=30 in each water tank) were exposed to two different test concentrations of a PBDEs mixture (PBDE 28, PBDE 47, PBDE 99, PBDE 100, PBDE 153, and PBDE 154) based on the maximum allowable concentrations according to the EU and national legislation – 0.14 µg/L and 0.0085 µg/kg. A water tank was used for control with PBDEs-free tap water. The water physicochemical parameters including pH (ISO 10523:2008), temperature (BDS: 17.1.4.01:1977), electrical conductivity (µS/cm, ISO 7888:1985), and dissolved oxygen (mg/L, ISO 17289:2014) were read three times per day using a multi-device (WTW, Germany).

### ***Histochemical assessment***

Mussel dissection was performed according to the methodology for bioaccumulated pollutants described in the EMERGE protocol of Rosseland et al. (2003), which we adapted for mussels. Each mussel was weighed with an analytical balance (KERN, Germany) (g) and measured with a caliper (cm) to calculate different condition factors further. The dissected-out gills were placed in sterile polyethylene plastic bags and deep-frozen.

Frozen gill tissue samples were fixed on pre-cooled aluminum chucks for cryostat sectioning. Deep-frozen multiple zebra mussel's gill sections (5 µm) were prepared using a cryotome (Leica CM 1520, Wetzlar, Germa-

ny) at a chamber temperature of  $-25^{\circ}\text{C}$ . The sections were stored after that at  $-80^{\circ}\text{C}$  for further staining procedures following a standard PAS methodology for polysaccharides according to McManus (1948) and Culing (1974) with slight modifications as previously described in our studies (Georgieva et al., 2013; Yancheva et al., 2019b; Kovacheva et al., 2022). The samples were fixed in absolute alcohol, rinsed quickly with distilled water, hydrolyzed with periodic acid solution, rinsed with tap and distilled water, stained with Schiff's reagent at room temperature, rinsed with warm tap water ( $35^{\circ}\text{C}$  minimum), then rinsed quickly with distilled water, counterstained with hemalum solution, colored in blue in flowing tap water, dehydrated by ascending alcohol series (50, 70, 80, 90 and 100%), cleared with clearing agent, dried at room temperature and capped with Canada balsam. Five slides per treated specimen ( $n=15$  from each tank) were prepared.

The gill histochemical changes, including the controls, were evaluated blindly and appraised individually and semi-quantitatively using the grading system of Mishra & Mohanty (2008) adopted for this study. A positive PAS reaction was presented in pink to violet staining, and the cell nuclei were colored in blue, respectively. Each grade represented specific histochemical characteristics and was categorized as follows: (0) - negative reaction of the histochemical staining; (1) - very weak positive reaction of the histochemical staining; (2) - weak positive reaction of the histochemical staining; (3) - moderate positive reaction of the histochemical staining; (4) - strong positive reaction of the histochemical staining.

#### *Statistical analysis*

The results from the conducted experiment were presented as mean $\pm$ SD for all evaluated individuals per experimental concentration. Past 3.03 (Hammer et al., 2001) and GraphPad Prism 7 for Windows (USA) were used to assess the data statistically. The normality of data was tested by the Shapiro-Wilk test. The homogeneity of variances was tested with the Levene's test. The results were also analyzed for the significance of

differences among the control and the treated mussel groups by the Kruskal-Wallis test, followed by the Mann-Whitney test (medians comparison). The levels of statistical significance were set at  $p < 0.05$ .

#### **Results and Discussion**


The physicochemical parameters of the water contaminated with PBDEs stayed relatively constant throughout the experiment. No significant changes between the control and the tested aquaria ( $p > 0.05$ ) were observed; hence, they will not be further discussed as we did not link the changes in the glycogen amount with their values, but rather with the PBDEs' toxicity.

The degree of intensity of the applied histochemical PAS reaction in the zebra mussel gills after the acute and subchronic exposure to both experimental PBDEs concentrations is presented in Figures 1 and 2. Overall, the results showed a decrease in the amount of glycogen compared to the control group. There were no detectable differences in the amount of glycogen between the two experimental PBDEs concentrations administered after the 96 hours' exposure. Such were not found on day 30, either. The only difference was found in the glycogen amount when comparing the results between the 96 hours' and 30 days' exposure, but they were not statistically significant ( $p > 0.05$ ). We consider that this data could indicate that, in our case, the PBDE toxicity depends on the time of action of the toxicant, but not on the administered dose, which, from our perspective, requires further research.


Our results are in line with Ansaldo et al. (2006), that the glycogen levels are negatively correlated with the level of chemical stress, and with Hamed et al. (2023), who found that after seven days of exposure to several concentrations of ZnO nanoparticles, the glycogen levels exhibited a significant reduction in the gill tissues. As explained by Arrighetti et al. (2019), the gills of bivalves play an essential role in the feeding process, and they are also the primary organ for the entry of pollutants. In addition, glycogen is the molecule that functions as secondary long-term energy storage in animal and fungi cells, it is found in the

form of granules in the cytosol and plays a vital role in the glucose cycle (Brenner et al., 2014). Therefore, glycogen is the primary energy reserve in bivalves. According to Brenner et al. (2014), glycogen forms an energy reserve that can be quickly mobilized to meet a sudden need for glucose. When living in a stressful environment affected by various anthropogenic pollution, the glycogen levels tend to decrease to supply different energetic requirements due to elevated metabolism in the exposed organisms (Hyötyläinen et al., 2002). We agree that changes in the amount of glycogen in the cells indicate changes in the carbohydrate metabolism of the organism under the adverse effect of various pollutants

and that the reduced glycogen storage represents a non-specific reaction of the zebra mussels against the PBDE exposure in our case. We also express the same opinion of Au (2004) that such histochemical biomarkers are unspecific and easy to determine, and their advantage is that they could be directly related to the health status of the organism. Also, as an early warning tool for assessing environmental risk, histochemical methods can be used in combination with biomarkers at other organizational levels, such as different enzyme activities (cellular level), condition factors, and shell analysis (organismal level), etc. (Brooks et al., 2015; Rementeria et al., 2017; Bebianno et al., 2018).



**Fig. 1.** Glycogen levels (PAS-reaction) decrease in the gills of zebra mussels (n=15 individuals per treatment) after exposure to two different concentrations of PBDEs (maximum allowable concentrations, MAC in water and biota) after 96 hours and 30 days of exposure (average±standard deviation).



**Fig. 2.** Glycogen levels (PAS-reaction) decrease in outer lamella and middle mantle of the gills of zebra mussels after 96 hours' exposure to PBDEs (maximum allowable concentrations, MAC in biota): A) control and B) treated individuals, x 400.

## Conclusions

In summary, we can conclude that the changes in the intracellular glycogen levels and storage capacities correlate with the duration of chemical stress. We also suggest that the PAS reaction can be used as a general tissue biomarker of nutritional status and health, and the effects of exposure of freshwater mussels to PBDEs, along with other biological tools in terms of the multi-biomarker approach.

**Acknowledgments:** This study is financed by the European Union-NextGenerationEU, through the National Recovery and Resilience Plan of the Republic of Bulgaria, project № BG-RRP-2.004-0001-C01.

## References

- Abbasi, G., Li, L., & Breivik, K. (2019). Global historical stocks and emissions of PBDEs. *Environmental Science and Technology*, 53(11), 6330–6340. doi: [10.1021/acs.est.8b07032](https://doi.org/10.1021/acs.est.8b07032)
- Ahrendt, C., Galbán-Malagón, C., Gómez, V., Torres, M., Mattar, C., DeCoite, M., Guida, Y., Příbylová, P., & Pozo, K. (2023). Marine debris and associated organic pollutants in surface waters of Chiloé in the Northern Chilean Patagonia (42°–44°S). *Marine Pollution Bulletin*, 187, 114558. doi: [10.1016/j.marpolbul.2022.114558](https://doi.org/10.1016/j.marpolbul.2022.114558)
- Al-Harbi, M., Al-Enzi, E., Al-Mutairi, H., & Whalen, J. K. (2021). Human health risks from brominated flame retardants and polycyclic aromatic hydrocarbons in indoor dust. *Chemosphere*, 282, 131005. doi: [10.1016/j.chemosphere.2021.131005](https://doi.org/10.1016/j.chemosphere.2021.131005)
- Angelov, A. (2000). Catalogus faunae bulgaricae. 4. Mollusca: Gastropoda et Bivalvia aquae dulcis. Sofia, Bulgaria: Pensoft, 57 p.
- Ansaldo, M., Nahabedian, D. E., Holmes-Brown, E., Agote, M., Ansay, C. V., Guerrero, N. R. V., & Wider, E. A. (2006). Potential use of glycogen level as a biomarker of chemical stress in *Biomphalaria glabrata*. *Toxicology*, 224(1–2), 119–127. doi: [10.1016/j.tox.2006.04.037](https://doi.org/10.1016/j.tox.2006.04.037)
- Arrighetti, F., Landro, S. M., Lambre, M. E., Penchaszadeh, P. E., & Teso, V. (2019). Multiple-biomarker approach in the assessment of the health status of a novel sentinel mussel *Brachidontes rodriguezii* in a harbor area. *Marine Pollution Bulletin*, 140, 451–461. doi: [10.1016/j.marpolbul.2019.02.002](https://doi.org/10.1016/j.marpolbul.2019.02.002)
- Au, D. W. T. (2004). The application of histocytopathological biomarkers in marine pollution monitoring: A review. *Marine Pollution Bulletin*, 48, 817–834. doi: [10.1016/j.marpolbul.2004.02.032](https://doi.org/10.1016/j.marpolbul.2004.02.032)
- Baratange, C., Baali, H., Gaillet, V., Bonnard, I., Delahaut, L., Gaillard, J. C., Grandjean, D., Sayen, S., Gallorini, A., Le Bris, N., Renault, D., Breider, F., Loizeau, J. L., Armengaud, J., & Cosio, C. (2023). Bioaccumulation and molecular effects of carbamazepine and methylmercury co-exposure in males of *Dreissena polymorpha*. *Science of the Total Environment*, 897, 165379. doi: [10.1016/j.scitotenv.2023.165379](https://doi.org/10.1016/j.scitotenv.2023.165379)
- BDS 17.1.4.01:1977 - Nature protection. Hydrosphere. Water quality indicators. A method for determining smell, color and temperature (In Bulgarian).
- Bebianno, M. J., Cardoso, C., Gomes, T., Blasco, J., Santos, R. S., & Colaço, A. (2018). Metal interactions between the polychaete *Branchipolynoe seepensis* and the mussel *Bathymodiolus azoricus* from Mid-Atlantic-Ridge hydrothermal vent fields. *Marine Environmental Research*, 135, 70–81. doi: [10.1016/j.marenvres.2018.01.017](https://doi.org/10.1016/j.marenvres.2018.01.017)
- Beyer, J., Green, N. W., Brooks, S., Allan, I. J., Ruus, A., Gomes, T., Bråte, I. L. N., & Schøyen, M. (2017). Blue mussels (*Mytilus edulis* spp.) as sentinel organisms in coastal pollution monitoring: A review. *Marine Environmental Research*, 130, 338–365. doi: [10.1016/j.marenvres.2017.07.024](https://doi.org/10.1016/j.marenvres.2017.07.024)
- Binelli, A., Della Torre, C., Magni, S., & Parolini, M. (2015). Does zebra mussel (*Dreissena polymorpha*) represent the freshwater counterpart of *Mytilus* in ecotoxicological studies? A critical review. *Environmental Pollution*, 196, 386–403. doi: [10.1016/j.envpol.2014.10.023](https://doi.org/10.1016/j.envpol.2014.10.023)
- Brenner, M., Broeg, K., Frickenhaus, S., Buck, B. H., & Koehler, A. (2014). Multi-biomarker approach using the blue mussel (*Myti-*

- lus edulis* L.) to assess the quality of marine environments: Season and habitat-related impacts. *Marine Environmental Research*, 95, 13–27. doi: [10.1016/j.marenvres.2013.12.009](https://doi.org/10.1016/j.marenvres.2013.12.009)
- Brooks, S. J., Harman, C., Hultman, M. T., & Berge, J. A. (2015). Integrated biomarker assessment of the effects of tailing discharges from an iron ore mine using blue mussels (*Mytilus* spp.). *Science of the Total Environment*, 524–525, 104–114. doi: [10.1016/j.scitotenv.2015.03.135](https://doi.org/10.1016/j.scitotenv.2015.03.135)
- Culling, C. F. A., Allison, R. T., & Barr, W. T. (1985). *Cellular pathology technique*. (4<sup>th</sup> Edition, pp. 216–220). Oxford, United Kingdom: Butterworth-Heinemann.
- EC. (2010). Directive 2010/63/EU of the European Parliament and of the Council on the protection of animals used for scientific purposes. *Official Journal of the European Union*, L276, 33–79. Retrieved from <https://eur-lex.europa.eu/>
- Faria, M., Huertas, D., Soto, D. X., Grimalt, J. O., Catalan, J., Riva, M. C., & Barata, C. (2010). Contaminant accumulation and multi-biomarker responses in field collected zebra mussels (*Dreissena polymorpha*) and crayfish (*Procambarus clarkii*), to evaluate toxicological effects of industrial hazardous dumps in the Ebro river (NE Spain). *Chemosphere*, 78(3), 232–240. doi: [10.1016/j.chemosphere.2009.11.003](https://doi.org/10.1016/j.chemosphere.2009.11.003)
- Gecheva, G., Yancheva, V., Velcheva, I., Georgieva, E., Stoyanova, S., Arnaudova, D., Stefanova, V., Georgieva, D., Genina, V., Todorova, B., & Mollov I. (2020). Integrated monitoring with moss-bag and mussel transplants in reservoirs. *Water*, 12, 1800. doi: [10.3390/w12061800](https://doi.org/10.3390/w12061800)
- Georgieva, E., Antal, L., Stoyanova, S., Arnaudova, D., Velcheva, I., Iliev, I., Vasileva, T., Bivolarski, V., Mitkovska, V., Chassovnikarova, T., Todorova, B., Uzochukwu, I. E., Nyeste, K., & Yancheva, V. (2022). Biomarkers for pollution in caged mussels from three reservoirs in Bulgaria: a pilot study. *Heliyon*, 8(3), e09069. doi: [10.1016/j.heliyon.2022.e09069](https://doi.org/10.1016/j.heliyon.2022.e09069)
- Georgieva, E., Atanasova, P., Velcheva, I., Stoyanova, S. & Yancheva, V. (2013). Histochemical effects of “Verita WG” on glycogen and lipid storage in common carp (*Cyprinus carpio* L.) liver. *Ecologia Balkanica*, 5(2), 91–97.
- Goldberg, E. D. (1975). The mussel watch – A first step in global marine monitoring. *Marine Pollution Bulletin*, 6(7), 111. doi: [10.1016/0025-326X\(75\)90271-4](https://doi.org/10.1016/0025-326X(75)90271-4)
- Hamed, E. S. A. E., Said, S. M., Abd El-Baky, R. F., & Mohammad, W. A. (2023). Assessment the toxicological perturbations of zinc oxide nanoparticles on histopathology, biochemistry and protein profile in marine mussels *Brachidontes pharaonis*. *The Egyptian Journal of Aquatic Research*, in press. doi: [10.1016/j.ejar.2023.08.004](https://doi.org/10.1016/j.ejar.2023.08.004)
- Hani, Y. M. I., Prud’Homme, S. M., Nuzillard, J. M., Bonnard, I., Robert, C., Nott, K., Ronkart, S., Dedourge-Geffard, O., & Geffard, A. (2021). 1H-NMR metabolomics profiling of zebra mussel (*Dreissena polymorpha*): A field-scale monitoring tool in ecotoxicological studies. *Environmental Pollution*, 270, 116048. doi: [10.1016/j.envpol.2020.116048](https://doi.org/10.1016/j.envpol.2020.116048)
- Hubenov, Z. (2005). *Dreissena* (Bivalvia: Dreissenidae) – systematics, autochthonous and anthropogenic areas. *Acta Zoologica Bulgarica*, 57(3), 259–268.
- Hyötyläinen, T., Karels, A., & Oikari, A. (2002). Assessment of bioavailability and effects of chemicals due to remediation actions with caging mussels (*Anodonta anatina*) at a creosote-contaminated lake sediment site. *Water Research*, 36(18), 4497–504. doi: [10.1016/s0043-1354\(02\)00156-2](https://doi.org/10.1016/s0043-1354(02)00156-2)
- ISO 10523:2008 - Water quality. Determination of pH.
- ISO 7888:1985 - Water quality. Determination of electrical conductivity.
- ISO 17289:2014 - Water quality. Determination of dissolved oxygen.
- Kovacheva, E., Georgieva, E., Velcheva, I., Iliev, I., Vasileva, T., Bivolarski, V., Nikolova, M., Todorova, B., Todorova-Bambaldokova, D., Yancheva, V., & Tomov, S. (2022). Histochemical and biochemical changes in common carp (*Cyprinus carpio* Linnaeus, 1785) liver after cypermethrin and chlorpyrifos exposure. *Ecologia Balkanica*, 14(2), 123–141.
- Lepom, P., Irmer, U., & Wellnitz, J. (2012). Mercury levels and trends (1993–2009) in

- breem (*Abramis brama* L.) and zebra mussels (*Dreissena polymorpha*) from German surface waters. *Chemosphere*, 86(2), 202–211. doi: [10.1016/j.chemosphere.2011.10.021](https://doi.org/10.1016/j.chemosphere.2011.10.021)
- Li, Y., Liu, C., Yang, H., He, W., Li, B., Zhu, X., Liu, S., Jia, S., Li, R., & Tang, K. H. D. (2024). Leaching of chemicals from microplastics: A review of chemical types, leaching mechanisms and influencing factors. *Science of the Total Environment*, 906, 167666. doi: [10.1016/j.scitotenv.2023.167666](https://doi.org/10.1016/j.scitotenv.2023.167666)
- Louis, F., Rocher, B., Barjhoux, I., Bultelle, F., Dedourge-Geffard, O., Gaillet, V., Bonnard, I., Delahaut, L., Pain-Devin, S., Geffard, A., Paris-Palacios, S., & David, E. (2020). Seasonal monitoring of cellular energy metabolism in a sentinel species, *Dreissena polymorpha* (bivalve): Effect of global change? *Science of the Total Environment*, 725, 138450. doi: [10.1016/j.scitotenv.2020.138450](https://doi.org/10.1016/j.scitotenv.2020.138450)
- McManus, J. F. A. (1948). Histological and histochemical uses of periodic acid. *Stain Technology*, 23, 99–108. doi: [10.3109/10520294809106232](https://doi.org/10.3109/10520294809106232)
- Medina, G. (2024). Persistent organic pollutants. *Encyclopedia of Toxicology*, 409–418. doi: [10.1016/B978-0-12-824315-2.01149-0](https://doi.org/10.1016/B978-0-12-824315-2.01149-0)
- Mishra, A. K., & Mohanty, B. (2008). Acute toxicity impacts of hexavalent chromium on behavior and histopathology of gill, kidney and liver of the freshwater fish, *Channa punctatus* (Bloch). *Environmental Toxicology and Pharmacology*, 26, 136–141. doi: [10.1016/j.etap.2008.02.010](https://doi.org/10.1016/j.etap.2008.02.010)
- Portet-Koltalo, F., G., N., Morin, C., de Mengin-Fondragon, F., & Frouard, A. (2021). Evaluation of polybrominated diphenyl ether (PBDE) flame retardants from various materials in professional seating furnishing wastes from French flows. *Waste Management*, 131, 108–116. doi: [10.1016/j.wasman.2021.05.038](https://doi.org/10.1016/j.wasman.2021.05.038)
- Rementeria, A., Mikolaczyk, M., Peña, A., Lancelleur, L., Blanc, G., Soto, M., Schäfer, J., & Zaldibar, B. (2017). Assessment of health status of oysters (*Crassostrea gigas*) exposed to environmentally relevant concentrations of Ag and Cu in brackish waters. *Journal of Sea Research*, 130, 229–238. doi: [10.1016/j.seares.2017.04.013](https://doi.org/10.1016/j.seares.2017.04.013)
- Rosseland, B. O., Massabuau, J. C., Grimalt, J., Hofer, R., Lackner, R., Raddum, G., Rognerud, S., & Vives, I. (2003). Fish ecotoxicology: European mountain lake ecosystems regionalisation, diagnostic and socio-economic evaluation (EMERGE). Fish sampling manual for live fish. Norwegian Institute for Water Research (NIVA), Oslo, pp. 8.
- Russev, B., Yaneva, I., Detcheva, R., & Karapetkova, M. (1994). Zusammensetzung der Hydrofauna. In: Russev B. (Ed.) *Limnologie der Bulgarischen Donauzuflüsse*. (pp. 130–174). Sofia, Bulgaria: Knizhen Tigar Publ. (In Bulgarian, German summary).
- Secor, C. L., Mills, E. L., Harshbarger, J., Thomas Kuntz, H., Gutenmann, W. H., & Lisk, D. J. (1993). Bioaccumulation of toxicants, element and nutrient composition, and soft tissue histology of zebra mussels (*Dreissena polymorpha*) from New York State waters. *Chemosphere*, 26(8), 1559–1575. doi: [10.1016/0045-6535\(93\)90224-S](https://doi.org/10.1016/0045-6535(93)90224-S)
- Stoyanova, S., Mollov, I., Velcheva, I., Georgieva, E., & Yancheva, V. (2020). Cadmium and polyaromatic hydrocarbons exposure changes the condition indices in zebra mussel, *Dreissena polymorpha* (Pallas, 1771): a case study. *Acta Zoologica Bulgaria*, S15, 141–146.
- Valkanov, A. (1957). Katalog unserer Schwarzmeerfauna. Arbeiten aus der Biologischen Meeresstation in Varna, Bulgarien, 19, 1–62. (In Bulgarian, German summary)
- Yancheva, V., Mollov, I., Velcheva, I., Georgieva, E., & Stoyanova, S. (2016). Effects of cadmium (Cd) on the lysosomal membrane stability and respiratory rate of two freshwater mollusks under ex situ exposure: preliminary data. *South-Western Journal of Horticulture, Biology and Environment*, 7(1), 27–34.
- Yancheva, V., Mollov, I., Georgieva, E., Stoyanova, S., Tsvetanova, V., & Velcheva, I. (2017a). Ex situ effects of chlorpyrifos on the lysosomal membrane stability and respiration rate in Zebra mussel (*Dreissena polymorpha* Pallas, 1771). *Acta Zoologica Bulgarica*, S8, 85–90.

*Polybrominated diphenyl ethers' exposure causes changes in the physiology of zebra mussel, Dreissena polymorpha (Pallas, 1771)*

- Yancheva, V., Mollov, I., Velcheva, I., Stoyanova, S., Todorova, K. & Georgieva, E. (2017b). Lysosomal membrane stability and respiration rate in zebra mussel (*Dreissena polymorpha* Pallas, 1771) as biomarkers for *ex situ* heavy metal exposure. *Periodicum Biologorum*, 119(4), 229-237.
- Yancheva, V., Stoyanova, S., Georgieva, E. & Velcheva, I. (2018a). Mussels in ecotoxicological studies - are they better than fish?. *Ecologia Balkanica*, 10(1), 57-84.
- Yancheva, V., Georgieva, E., Stoyanova, S., Tsvetanova, V., Todorova, K., Mollov, I. & Velcheva, I. (2018b). Short and long-term toxicity of cadmium (Cd) and polyaromatic hydrocarbons (PAHs) on zebra mussel (*Dreissena polymorpha* Pallas, 1771). *Acta Zoologica Bulgarica*, 70(4), 557-564.
- Yancheva, V., Velcheva, I., Georgieva, E. & Stoyanova, S. (2019a). Bioaccumulation of polyaromatic hydrocarbons (PAHs) and cadmium (Cd) and its toxic effects on zebra mussel *Dreissena polymorpha* (Pallas, 1771) (Bivalvia: Dreissenidae). *Acta Zoologica Bulgarica*, 71(4), 567-574.
- Yancheva, V., Velcheva, I., Georgieva, E. & Stoyanova, S. (2019b). Periodic Acid - Schiff (PAS) reaction in fish liver exposed to fungicide contamination: A possible histochemical biomarker. *Ecologia Balkanica*, 11(1), 1-10.
- Yancheva, V., Velcheva, I., Iliev, I., Vasileva, T., Bivolarski, V., Georgieva, E. & Stoyanova, S. (2020). Histochemical and biochemical alterations in zebra mussel, *Dreissena polymorpha* (Pallas, 1771) after cadmium and polyaromatic hydrocarbons exposure. *Acta Zoologica Bulgarica*, S15, 155-164.
- Yancheva, V., Mollov, I., Stoyanova, S., Todorova, B., Velcheva, I. & Georgieva, E. (2021). Toxic pesticides effects on the respiration rate in *Dreissena polymorpha* (Pallas, 1771). *Zoonotes*, 173, 1-4.
- Yancheva, V., Stoyanova, S., Todorova, B., Georgieva, E., & Velcheva, I. (2022). Zebra mussel (*Dreissena polymorpha* Pallas, 1771): the invasive bioindicator for freshwater quality? *Zoonotes*, 197, 1-4.
- Yang, C., Jiang, Y., Zhao, W., Peng, J., Liu, Y., Lin, X., & Zhang, J. (2023). Characterization and distribution of polybrominated diphenyl ethers in shellfish in Shenzhen coastal waters and assessment of human health risks. *Marine Pollution Bulletin*, 191, 114957. doi: [10.1016/j.marpolbul.2023.114957](https://doi.org/10.1016/j.marpolbul.2023.114957)

Received: 30.10.2023  
Accepted: 30.11.2023