


# Antihyperlipidemic activity of myricetin

Z.Z. Yang, J.Q. Zhang, B.B. Shi, J.Q. Qian\*  and H. Guo

College of Pharmaceutical Science, Zhejiang University of Technology, 18 Chaowang Road, Hangzhou 310014, China

## ORIGINAL RESEARCH PAPER

Received: March 25, 2024 • Accepted: May 18, 2024

Published online: June 7, 2024

© 2024 The Author(s)



### ABSTRACT

Myricetin has been reported to have a wide variety of beneficial physiological functions. The present study investigated the antihyperlipidaemic activity of myricetin against hyperlipidaemia of high-fat diet-fed obese rats. The four-week antihyperlipidaemic activity was assayed by giving different doses of myricetin to hyperlipidaemic rats. Results showed that myricetin could reduce the harm caused by oxidative stress, decrease thiobarbituric acid reactive substances value, and decrease total cholesterol and triacylglycerol levels of hyperlipidaemic rats. Quantitative analysis of gene expression showed that myricetin's lipid-lowering activity can be activated by downregulating gene expression of fatty acid synthase (FAS) and peroxisome proliferator-activated receptor- $\gamma$  (PPAR $\gamma$ ) with upregulation of hormone-sensitive lipase (HSL) mRNA level. Thus, myricetin had significant health benefits and could be explored as a potentially promising dietary supplement for treating hyperlipidaemia.

### KEYWORDS

antihyperlipidaemic, myricetin, oxidative stress, gene expression

## 1. INTRODUCTION

Hyperlipidaemia is a disease characterised by a disorder in lipid metabolism, manifesting as abnormally elevated levels of plasma lipids (including triglycerides and cholesterol) or lipoproteins. Usually, this type of disease is closely associated with cardiovascular disease, fatty liver

\* Corresponding author. E-mail: qianjq757@163.com

disease, cancer, etc (Gao et al., 2013). The incidence of hyperlipidaemia and its complications has increased significantly in the last few decades. Statins (hydroxy methylglutaryl coenzyme A reductase inhibitors) are typical antihyperlipidaemia drugs, but they account for 61% of side effect types of all antihyperlipidaemic drugs (Jacquet et al., 1993). It can cause certain damage to the skin, digestive system, liver, and kidney. It may be accompanied by severe myolysis side effects and certain neurotoxicity, among which muscle pain caused by myolysis side effects is only triggered by statins. However, plants usually contain phytochemicals with various physiological activities, including some compounds with high activity and almost non-toxic side effects, such as some flavonoids, and appropriate supplementation helps prevent various diseases (O'Day, 2023).

Myricetin (3,3',4',5,5',7-hexahydroxyflavone) is a dietary flavonol without toxicity (Huwait et al., 2023). The compound exhibits many activities, including strong antioxidant, antineoplastic, antihyperlipidaemic, and antiinflammatory (Singh and Bast, 2015; Semwal et al., 2016). Myricetin can reduce blood lipids (He et al., 2016) and improve fatty liver caused by hyperlipidaemia (Xia et al., 2019). The study on myricetin's *in vivo* antihyperlipidaemic activity remains limited, nonetheless. In the present study, we investigated the activity of high-purity myricetin on a high-fat diet (HFD) feeding induced hyperlipidaemia rats to reveal whether myricetin could effectively improve hyperlipidaemia.

Lipogenesis and lipolysis are major factors affecting lipid accumulation in tissues. Fatty acid synthase (FAS) mainly exists in adipose tissues and is critical to lipogenesis, which is an important regulator of triacylglycerol homeostasis (Loftus et al., 2000). Hormone-sensitive lipases (HSL) in the human body mainly catalyse the hydrolysis of triacylglycerol and their metabolites to produce fatty acids, which then enter mitochondria to participate in oxidative reactions and energy supply. This step plays an important role in lipid metabolism (Zechner and Langin, 2014). Peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ), primarily located in adipose and immune tissues, plays an indispensable role in adipogenesis and lipid redistribution (Chen et al., 2014). In this study, we have shown that myricetin may prevent the development of hyperlipidaemia via down-regulating FAS and PPAR $\gamma$  expression and up-regulating HSL expression. The animal experiment shows that myricetin could improve hyperlipidaemia.

## 2. MATERIALS AND METHODS

### 2.1. Chemicals

The crude extract of myricetin (51.3% content of myricetin) was extracted from *Myrica rubra* in Ningbo, Zhejiang Province with 95% ethanol. High-purity myricetin was prepared by silica gel column chromatography with 70% ethanol as the eluent. The high-purity eluent was pooled, and most of the ethanol was removed by spin evaporation. High purity myricetin sample (98.1%, HPLC) was prepared by vacuum freeze drying of concentrated myricetin aqueous solution. The characterisation data obtained by liquid chromatography, nuclear magnetic resonance, and mass spectrometry were consistent with the literature (Qian et al., 2022).

The DNA Ladder and agarose were purchased from Dalian Takara (Dalian, China). Diethyl pyrocarbonate, TRIzol reagent, M-MLV reverse transcriptase, PCR kit (Cat No. YSP97313), and fluorescent quantitative PCR kit (Cat. No. YSP97316) were purchased from Sangon Biotech (Shanghai, China). Tris base and ethylene diamine tetraacetic acid were from Promega



(Madison, WI, USA). All other reagents were from Sigma–Aldrich (St. Louis, MO, USA), except where specified otherwise.

## 2.2. Quantitative detection and nuclear magnetic characterisation of myricetin

The High-Performance Liquid Chromatography (HPLC) conditions were: instrument Agilent 1200 (Santa Clara, USA), Eclipse XDB-C18 silica gel column ( $4.6 \times 150$  mm,  $5 \mu\text{m}$ ); mobile phase: methanol: water: acetic acid = 50:49:1 (v/v/v); injection volume:  $5 \mu\text{L}$ ; column temperature:  $30^\circ\text{C}$ ; flow rate:  $1.0 \text{ mL min}^{-1}$ ; UV absorption detector; detection wavelength: 292 nm. The myricetin sample obtained by purification was detected qualitatively using a standard substance of 99.5% purity. The purity of the myricetin sample was 98.1% by HPLC (Fig. 1).

The nuclear magnetic spectrum (Avance III 500 MHz, Bruker, Billerica, USA) of myricetin is shown in Fig. 2A ( $^{13}\text{C}$  NMR) and Fig. 2B ( $^1\text{H}$  NMR). The peaks around the  $\delta 40$  of the chemical shift were solvent DMSO- $d_6$  peaks, and  $\delta 93.1$  to  $175.7$  representing 15 carbons with 13 different chemical shifts (Fig. 2A). The  $\delta 2.51$  and  $3.36$  in Fig. 2B were solvent DMSO- $d_6$  and water peaks, respectively.  $\delta 6.18$  to  $12.51$  represent 10 hydrogens with 8 different chemical shifts, respectively. The NMR data are as follows:  $^{13}\text{C}$  NMR (500 MHz, DMSO- $d_6$ ): 93.1 (C-8), 98.1 (C-6), 102.9 (C-10), 107.1 (C-2', C-6'), 120.7 (C-1'), 135.8 (C-3, C-4'), 145.7 (C-3', C-5'), 146.8 (C-2), 156.0 (C-9), 160.7 (C-5), 163.8 (C-7), 175.7 (C-4).  $^1\text{H}$  NMR (500 MHz, DMSO- $d_6$ ): 6.18 (d,  $J = 1.85$  Hz, 1H, H-6), 6.37 (d,  $J = 1.80$  Hz, 1H, H-8), 7.25 (s, 2H, H-2', 6'), 8.84 (s, 1H, 4'-OH), 9.24 (s, 2H, 3', 5'-OH), 9.36 (s, 1H, 3-OH), 10.80 (s, 1H, 7-OH), 12.51 (s, 1H, 5-OH).

The molecular formula of myricetin is  $\text{C}_{15}\text{H}_{10}\text{O}_8$ . The molecular structure is 3,5,7,3',4',5' hydrogen atoms at the aromatic ring skeleton formed by hydroxyl substitution. The nuclear magnetic spectrum of the sample and the standard nuclear magnetic data (Devi et al., 2015) were compared and the structure of myricetin was qualitatively analysed to confirm the molecular structure of the myricetin sample.




Fig. 1. Chromatogram of myricetin sample





### 2.3. Animals and experimental design

For this experiment, male Wistar rats (weight  $180 \pm 10$  g) were purchased from the Laboratory Animal Center of Zhejiang Medical Sciences Academy. They were housed at ambient temperature ( $20 \pm 2$  °C) with an illumination period of 12 h light per day. Animals were fed standard rat food (Slaccas, Shanghai, China) with unlimited water intake. Food intake and body weight of rats were measured weekly during the experimental period.

After 5 days of adaptive observation, the hyperlipidaemia model in rats was established after confirming that the purchased rats had no abnormal state. The rats were randomly assigned to either the normal diet group (10 rats) fed with standard rat food or the high-fat diet group (50 rats) fed with high-fat forage (Slaccas, Shanghai, China) composed of standard rat food (73.8%, w/w), lard (15%, w/w), powdered egg yolk (10%, w/w), cholesterol (1%, w/w), and bile salt (0.2%, w/w).

Poor water solubility of myricetin results in excessive drug loss during animal gavage. Myricetin powder was weighed precisely and mixed evenly into 1% carboxymethyl cellulose sodium (CMC) to prepare CMC suspension of different drug concentrations to avoid this situation (Fukuchi et al., 2004).

After 2 weeks of feeding, blood samples of HDF group rats were collected by venipunctures of the posterior orbital nerve plexus to measure their lipid levels. The lipid levels of all HFD group rats were in line with the indicators of hyperlipidaemia, which proved that the hyperlipidaemia model was successfully established. The HFD-fed rats were randomly divided into 5 groups with 10 rats in each group (Chat et al., 2013). The model control group, positive control group, and three dosage groups were set up, respectively. The given simvastatin dosage for the positive control group was  $50 \text{ mg kg}^{-1} \text{ d}^{-1}$ . The dosage of myricetin in the experimental group was 100, 200, and  $400 \text{ mg kg}^{-1} \text{ d}^{-1}$  (He et al., 2016). Myricetin-admin suspensions of 50, 100, and  $200 \text{ mg mL}^{-1}$  were prepared and administered intragastrically at a volume of  $2 \text{ mL kg}^{-1}$  once a day. A  $25 \text{ mg mL}^{-1}$  suspension of Simvastatin-CMC was administered intragastrically at a fixed volume of drug liquid. During the administration period of the drug treatment group, the model control group and the normal group were treated with the same method by gavage of 1% CMC solution with the same volume as administration. In addition, except for the normal group, rats in the other groups were continuously fed with HFD. After 4 weeks of treatment, all animals were fasted for 12 h followed by collection of blood samples. At the end of the study, the animals were euthanised by cervical dislocation under ether anaesthesia. The liver and white adipose tissue (WAT) from the viscera and subcutaneous layer were taken out, rinsed with a physiological saline solution, weighed, and immediately stored at  $-70$  °C.

The experiment was approved by the Experimental Animal Ethics Committee of Zhejiang University of Technology (Ethical approval number: 20200511051).

### 2.4. Blood lipid detection

After 4 weeks of myricetin treatment, the animals were fasted for 12 h then blood samples were collected without anticoagulation. Serum samples were separated from the blood by clotting for 2 h at room temperature and centrifuging at  $1,500 \text{ g}$  for 15 min. To assess the serum levels of total cholesterol (TC), triglycerides (TG), and low-density lipoprotein cholesterol (LDL-C), detection kits of TC (Cat. No. A111-2-1, 100T/96 samples), TG (Cat. No. A110-2-1, 100T/96



samples), and LDL-C (Cat. No. A113-2-1, 100T/96 samples) were used (Nanjing Jiancheng Bioengineering Inst., Nanjing, China).

## 2.5. Lipid peroxidation, SOD, and GSH-Px assays

To assess the activity of two liver antioxidant enzymes, superoxide dismutase (SOD) and glutathione peroxidase (GSH-Px), specific detection kits of SOD (Cat. No. A001-1-2, 100T/96 samples) and GSH-Px (Cat. No. A005-1-2, 100T/48 samples) were used (Nanjing Jiancheng Bioengineering Inst., Nanjing, China). The results were expressed as U/mg protein. Lipid peroxidation was measured with the thiobarbituric acid reactive substances (TBARS) of commercially available kits (Cat. No. A106-1-3, 100T, Nanjing Jiancheng Bioengineering Inst., Nanjing, China) and expressed as nmol of malondialdehyde equivalents (MDAeq) per milligram of protein.

## 2.6. Gene expression studies

Total RNA was extracted from the WAT of the rat. The first-strand cDNA was synthesised using the Revert Aid First Strand cDNA synthesis kit (Cat. No. 04896866001, Roche, Germany). The expression of selected genes was analysed by real-time PCR (StepOnePlus, Thermo Fisher Scientific, Waltham, USA). PCR conditions were as follows: 1 cycle of 94 °C for 4 min, 36 cycles of 94 °C for 30 s, 55 °C for 45 s, and 72 °C for 60 s, followed by 1 cycle of 72 °C for 10 min. Relative gene expression was calculated by normalising cycle threshold (Ct) values for genes of interest with Ct values for  $\beta$ -actin using the delta-delta Ct method. Primer sequences were as follows: HSL (forward: 5'-GGAGCATACAAA CGCAAC-3'; reverse: 5'-TCCCGTAGGTCATAG-GAGAT-3'); FAS (forward: 5'-CAGA GGGTGGTTGTTAGAAAGAT-3'; reverse: 5'-ACTGCCG-CATCCTCTTCCTC-3'); PPAR $\gamma$  (forward: 5'-CACAAGAGCTGACCCAATGGT-3'; reverse: 5'-CGCAGATCAGCAGACT CTGG-3');  $\beta$ -actin as a reference gene (forward: 5'-ACTGCCG-CATCCTCT TCCTC-3'; reverse: 5'-CTCCTGCTTGCTGATCCACATC-3'). The PCR products were analysed by agarose gel electrophoresis using 1.5% gel.

## 2.7. Statistical analysis

All data are represented as the mean  $\pm$  SD. Statistical analysis was conducted with SPSS 20.0 (SPSS Inc., Chicago, USA). The data were analysed by *t*-test. *P*-values less than 0.05 or 0.01 were considered statistically significant.

# 3. RESULTS AND DISCUSSION

## 3.1. Effects of myricetin on blood lipid

Serum and liver lipid levels of rats at the end of the experiments showed that the HFD caused the elevation of serum TC, TG, and LDL-C concentrations in rats. Serum TC, TG, and LDL-C levels in all hyperlipidaemic rats were equivalent before being given myricetin. The test results four weeks after the administration are shown in [Table 1](#).

The present study demonstrates myricetin-modified serum lipid levels by reducing TC and TG contents in rats fed a high-fat diet for 4 weeks. This diet, enriched in lard (15%), egg yolk



Table 1. Detection results of blood lipids in 4 weeks of myricetin feeding ( $\bar{x} \pm SD$ )

Groups	TC (mmol L <sup>-1</sup> )	TG (mmol L <sup>-1</sup> )	LDL-C (mmol L <sup>-1</sup> )
Normal control group	5.05 ± 0.31	1.62 ± 0.57	0.41 ± 0.06
Model control group	9.27 ± 0.54 <sup>##</sup>	1.96 ± 0.15 <sup>#</sup>	0.82 ± 0.20 <sup>##</sup>
Positive control group	6.92 ± 0.56 <sup>**</sup>	1.43 ± 0.17 <sup>**</sup>	0.42 ± 0.09 <sup>**</sup>
100 mg kg <sup>-1</sup> d <sup>-1</sup> of myricetin	7.52 ± 0.55 <sup>*</sup>	1.54 ± 0.08 <sup>*</sup>	0.49 ± 0.21 <sup>**</sup>
200 mg kg <sup>-1</sup> d <sup>-1</sup> of myricetin	6.83 ± 0.51 <sup>**</sup>	1.32 ± 0.23 <sup>**</sup>	0.43 ± 0.09 <sup>**</sup>
400 mg kg <sup>-1</sup> d <sup>-1</sup> of myricetin	5.04 ± 0.42 <sup>**</sup>	1.22 ± 0.21 <sup>**</sup>	0.36 ± 0.15 <sup>**</sup>

TC: total cholesterol; TG: triglycerides; LDL-C: LDL-cholesterol. <sup>#</sup>*P* < 0.05, <sup>##</sup>*P* < 0.01 VS. normal control group. <sup>\*</sup>*P* < 0.05, <sup>\*\*</sup>*P* < 0.01 vs. model control group.

powder (10%), cholesterol (1%), and bile salt (0.2%), was used because the combination of these components in a diet is universally used to induce hyperlipidaemia in rats and mice (Zhang et al., 2013). The results showed that the rats fed a high-fat diet had a higher concentration of TC, TG, and LDL-C in their sera than those fed a standard diet, indicating that the hyperlipidaemia model was successfully established.

Chronic administration of myricetin for 28 days, at the doses of 100 mg kg<sup>-1</sup> d<sup>-1</sup>, 200 mg kg<sup>-1</sup> d<sup>-1</sup>, and 400 mg kg<sup>-1</sup> d<sup>-1</sup>, provoked a significant reduction in serum levels of TC, TG, and LDL-C compared with vehicle-treated hyperlipidaemic rats. In the study, we showed that myricetin administration lowered lipid profile in HFD-feeding induced hyperlipidaemic rats.

Many researches have demonstrated that HFD-induced hyperlipidaemia has tight relations with oxidative stress (Jenkins et al., 2008). Recently, some preclinical observations have demonstrated that hyperlipidaemia promotes the accumulation of oxidised low-density lipoprotein (Ox-LDL) in the arterial wall, which plays a major role in the initiation and progression of atherosclerosis associated with hyperlipidaemia (Shan et al., 2019). In this work, we investigated the activities of three major defensive antioxidant agents. Our data showed that myricetin exhibited excellent antioxidant capacity to improve the activities of antioxidant enzymes (GSH-Px and T-SOD) and suppress lipid peroxidation (TBARS values) in hyperlipidaemic rats. This result suggest that the potential health benefits associated with myricetin have been partially attributed to its antioxidative properties.

### 3.2. Effects of myricetin on HFD-induced oxidative stress in liver

The effect of myricetin on the level of MDA equivalents of the rats' liver was determined, and the result is presented in Fig. 3A. The liver MDA equivalents significantly increased from 1.53 ± 0.36 nmol mg<sup>-1</sup> protein of the normal rats to 2.94 ± 0.47 nmol mg<sup>-1</sup> protein of the HFD-fed rats. By the administration of myricetin, the increase of MDA equivalents was significantly inhibited compared with the vehicle-treated HFD-fed group. Figure 3B shows the results that GSH-Px activity significantly increased in myricetin-treated groups compared to the vehicle-treated HFD-fed group. Moreover, Fig. 3C shows that an HFD-fed-induced decrease in SOD activity was prevented by the administration of myricetin at moderate (200 mg kg<sup>-1</sup> d<sup>-1</sup>) and high doses (400 mg kg<sup>-1</sup> d<sup>-1</sup>).

Moreover, myricetin has a unique chemical structure. The combined contribution of hydroxyl groups at the 3 and 5 positions and three continuous hydroxyl groups at positions 3', 4',






Fig. 3. Effects of myricetin or simvastatin ( $50 \text{ mg kg}^{-1} \text{ d}^{-1}$ ) on liver TBARS (3A), GSH-Px (3B), and SOD (3C) in the rats fed a high-fat diet for 4 consecutive weeks.

All data are presented as the mean  $\pm$  SD. ##  $P < 0.01$  VS. normal group. \*  $P < 0.05$ , \*\*  $P < 0.01$  VS. HFD group

and 5' can increase the antioxidant effectiveness of myricetin (Ma et al., 2022). T-SOD and GSH-Px are the major antioxidant enzymes that play a pivotal role in scavenging oxygen free radicals that protect the pancreatic tissue against oxidative stress injury. MDA concentration reflected the degree of the lipid peroxides. In our study, STZ treatment caused significant oxidative stress, indicated by significantly high levels of MDA and low activities of T-SOD and GSH-Px in the diabetic rats relative to the normal rats. Whereas, administration of myricetin prevents hepatic oxidative stress damage *via* normalising MDA, T-SOD, and GSH-Px levels. Complementing our findings, earlier studies have reported that myricetin could have antioxidant activity (Baldissera et al., 2016).

### 3.3. Effects on gene expression

The effects of myricetin on gene expression of FAS, HSL, and PPAR $\gamma$  are shown in Fig. 4A, all doses of myricetin increased the expression level of HSL. The low ( $100 \text{ mg kg}^{-1} \text{ d}^{-1}$ ), moderate ( $200 \text{ mg kg}^{-1} \text{ d}^{-1}$ ), and high doses ( $400 \text{ mg kg}^{-1} \text{ d}^{-1}$ ) of myricetin significantly increased








Fig. 4. Effects of myricetin on HSL (A), PPAR $\gamma$  (B), and FAS (C) in the rats fed a high-fat diet for 4 consecutive weeks.

All data are presented as the mean  $\pm$  SD. <sup>##</sup> $P < 0.01$  vs. normal group. <sup>\*</sup> $P < 0.05$ , <sup>\*\*</sup> $P < 0.01$  vs. HFD group

expression levels of HSL (23.26%, 41.86%, and 53.49%, respectively). Figure 4B displays that the expression level of PPAR $\gamma$  significantly decreased in myricetin-treated groups compared to the vehicle-treated HFD-fed group. Furthermore, Fig. 4C shows that the expression level of FAS was significantly higher in HFD-fed rats than that in the normal group; and those levels were lower by 36% in the myricetin (400 mg kg<sup>-1</sup> d<sup>-1</sup>) treated group.

HSL, abundant in the fatty tissue and other cholesterol-producing tissues, is the catabolic rate-limiting enzyme of lipolysis in animals, and plays a role in many aspects of lipid metabolism, as it hydrolyses triacylglycerol to monoacylglycerol and free fatty acid (Haemmerle et al., 2002; Gong et al., 2010). Recently, an increasing number of natural active substances have been extracted from plant and animal sources with antihyperlipidaemic effects *via* affecting lipolysis in adipocytes and hyperlipidaemic rats (Sarnaizul et al., 2013). In our



study, the expression level of the HSL gene significantly increased in the adipose tissue at myricetin administration. The result showed that myricetin might reduce lipids through increasing HSL gene expression.

Additionally, lipogenesis-associated genes such as FAS and PPAR $\gamma$  have also been reported as key transcriptional factors for lipid accumulation (Mangelsdorf et al., 1995). FAS plays an important role in denoting lipogenesis by converting acetyl-CoA and malonyl-CoA to palmitate and is related to various diseases such as obesity, cardiovascular disease, and cancer (Yeh et al., 2003). FAS may facilitate the synthesis and cytoplasmic storage of large triacylglycerols. When the FAS level is too high, triglyceride synthesis can be enhanced considerably (Zechner et al., 2009). Mice that are genetically engineered to overexpress an exogenous FAS gene, display increased lipid accumulation and lipogenesis (Postic and Girard, 2008). We also assayed the effect of myricetin on the expression of PPAR $\gamma$ , which is a lipid-activated transcription factor and highly expressed in white adipose tissue (Braissant et al., 1996), plays a central role in adipogenesis and lipid repartition (Weigt et al., 2013). Over-expression of PPAR $\gamma$  in adipocytes could selectively induce the production of lipoprotein lipase, regulate the signalling transduction, slow lipolysis speed, and increase the triacylglycerol synthesis (Ying et al., 2013). In macrophages, PPAR $\gamma$ -dependent lipid uptake and storage were initially suggested to increase foam-cell formation and, potentially, atherosclerosis (Lehrke and Lazar, 2005). We found that the expression levels of PPAR $\gamma$  and FAS genes in white adipose tissue in myricetin-treated HFD-fed rats significantly decreased as compared with the vehicle-treated counterparts. The results indicated that the action of myricetin to decrease lipid accumulation in hyperlipidaemic rats involved preventing lipogenesis beside stimulating lipolysis.

## 4. CONCLUSIONS

Myricetin exhibited excellent antihyperlipidaemic effects *in vivo*, with the improvement of oxidative status. It significantly lowers the concentration of serum TC, TG, and LDL-C. These effects appear to be mediated through down-regulation of adipogenesis-related genes (PPAR $\gamma$ ) and downstream genes (FAS), and up-regulation of lipolysis-related genes (HSL). Taken together, our results suggest that myricetin can serve as a dietary supplement in the management of hyperlipidaemia.

*Author contributions:* Zhengze Yang: investigation, formal analysis, data curation, writing; Jinqiu Zhang: methodology, validation, writing; Bobo Shi: methodology, data curation, writing; Junqing Qian: funding acquisition, project administration, conceptualisation, methodology, validation, supervision, writing; Hui Guo: methodology, validation. All authors contributed to writing the manuscript. All authors read and approved the final manuscript.

*Conflict of interest:* The authors declare that there are no conflicts of interest.

*Funding:* This work was supported by the China National Key R&D Program during the 13th Five-year Plan Period [grant numbers 2016YFD0400803], Zhejiang University of Technology, China [grant numbers KYY-HX-20210856].



## ACKNOWLEDGEMENTS

This research was supported by the Experimental Animal Center of Zhejiang University of Technology, Zhejiang, China. Animal experiments to lower blood lipids were approved by the Ethics Committee of Zhejiang University of Technology.

## REFERENCES

- Baldissera, G., Sperotto, N.D.M., Rosa, H.T., Henn, J.G., Peres, V.F., Moura, D.J., Roehrs, R., Denardin, E.L.G., Dal Lago, P., Nunes, R.B., and Saffi, J. (2016). Effects of crude hydroalcoholic extract of *Syzygium cumini* (L.) Skeels leaves and continuous aerobic training in rats with diabetes induced by a high-fat diet and low doses of streptozotocin. *Journal of Ethnopharmacology*, 194: 1012–1021.
- Braissant, O., Foufelle, F., Scotto, C., Dauca, M., and Wahli, W. (1996). Differential expression of peroxisome proliferator-activated receptors (PPARs): tissue distribution of PPAR-alpha, -beta, and -gamma in the adult rat. *Endocrinology*, 137(1): 354–366.
- Chat, O.A., Najar, M.H., and Dar, A.A. (2013). Evaluation of reduction kinetics of 2,2-diphenyl-1-picrylhydrazyl radical by flavonoid glycoside rutin in mixed solvent based micellar media. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 436: 343–353.
- Chen, G., Gao, X., Zhen, K.-S., Yin, Z.-Y., and Zheng, X.-X. (2014). Extract of *Zanthoxylum bungeanum* Maxim seed oil reduces hyperlipidemia in hamsters fed high-fat diet via activation of peroxisome proliferator-activated receptor  $\epsilon$ . *Tropical Journal of Pharmaceutical Research*, 13(11): 1837–1843.
- Devi, K.P., Rajavel, T., Habtemariam, S., Nabavi, S.F., and Nabavi, S.M. (2015). Molecular mechanisms underlying anticancer effects of myricetin. *Life Sciences*, 142: 19–25.
- Fukuchi, Y., Kato, Y., Okunishi, I., Matsutani, Y., Osawa, T., and Naito, M. (2004). 6-Methylsulfinylhexyl isothiocyanate, an antioxidant derived from *Wasabia japonica* MATUM, ameliorates diabetic nephropathy in type-2 diabetic mice. *Food Science and Technology Research*, 10(3): 290–295.
- Gao, H., Long, Y., Jiang, X., Liu, Z., Wang, D., Zhao, Y., Li, D., and Sun, B.-l. (2013). Beneficial effects of Yerba Mate tea (*Ilex paraguariensis*) on hyperlipidemia in high-fat-fed hamsters. *Experimental Gerontology*, 48(6): 572–578.
- Gong, J., Peng, C., Chen, T., Gao, B., and Zhou, H. (2010). Effects of theabrownin from pu-erh tea on the metabolism of serum lipids in rats: mechanism of action. *Journal of Food Science*, 75(6): H182–H189.
- Haemmerle, G., Zimmermann, R., Hayn, M., Theussl, C., Waeg, G., Wagner, E., Sattler, W., Magin, T.M., Wagner, E.F., and Zechner, R. (2002). Hormone-sensitive lipase deficiency in mice causes diglyceride accumulation in adipose tissue, muscle, and testis. *Journal of Biological Chemistry*, 277(7): 4806–4815.
- He, K., Li, X., Xiao, Y., Yong, Y., Zhang, Z., Li, S., Zhou, T., Yang, D., Gao, P., and Xin, X. (2016). Hypolipidemic effects of *Myrica rubra* extracts and main compounds in C57BL/6j mice. *Food and Function*, 7(8): 3505–3515.
- Huwait, E., Almassabi, R., Almowallad, S., Saddeek, S., Karim, S., Kalamegam, G., and Mirza, Z. (2023). Microarray expression profile of myricetin-treated THP-1 macrophages exhibits alterations in atherosclerosis-related regulator molecules and LXR/RXR pathway. *International Journal of Molecular Sciences*, 24(1): 278.



- Jacquet, A., Colomes, M., Ferrieres, J., Denat, M., Dousteblazy, P., and Monstastruc, J.L. (1993). A one-year prospective and intensive pharmacovigilance of antilipemic drugs in an hospital consultation for prevention of risk factors. *Therapie*, 48(5): 509–512.
- Jenkins, D.J.A., Kendall, C.W.C., Marchie, A., Josse, A.R., Nguyen, T.H., Faulkner, D.A., Lapsley, K.G., and Blumberg, J. (2008). Almonds reduce biomarkers of lipid peroxidation in older hyperlipidemic subjects. *Journal of Nutrition*, 138(5): 908–913.
- Lehrke, M. and Lazar, M.A. (2005). The many faces of PPAR gamma. *Cell*, 123(6): 993–999.
- Loftus, T.M., Jaworsky, D.E., Frehywot, G.L., Townsend, C.A., Ronnett, G.V., Lane, M.D., and Kuhajda, F.P. (2000). Reduced food intake and body weight in mice treated with fatty acid synthase inhibitors. *Science*, 288(5475): 2379–2381.
- Ma, X., Gong, H., Ogino, K., Yan, X., and Xing, R. (2022). Coordination-assembled myricetin nanoarchitectonics for sustainably scavenging free radicals. *Beilstein Journal of Nanotechnology*, 13: 284–291.
- Mangelsdorf, D.J., Thummel, C., Beato, M., Herrlich, P., Schutz, G., Umesono, K., Blumberg, B., Kastner, P., Mark, M., Chambon, P., and Evans, R.M. (1995). The nuclear receptor superfamily: the second decade. *Cell*, 83(6): 835–839.
- O'Day, D.H.H. (2023). Phytochemical interactions with calmodulin and critical calmodulin binding proteins involved in amyloidogenesis in Alzheimer's disease. *Biomolecules*, 13(4): 678.
- Postic, C. and Girard, J. (2008). Contribution of de novo fatty acid synthesis to hepatic steatosis and insulin resistance: lessons from genetically engineered mice. *Journal of Clinical Investigation*, 118(3): 829–838.
- Qian, J., Zhang, J., Chen, Y., Dai, C., Fan, J., and Guo, H. (2022). Hypoglycemic activity and mechanisms of myricetin. *Natural Product Research*, 36(23): 6177–6180.
- Sarnaizul, E., Borjihan, G., and Zhaorigetu, S. (2013). The preparation and antihyperlipidaemic assay of piperlonguminine *in vivo*. *Phytochemistry Letters*, 6(1): 101–105.
- Semwal, D.K., Semwal, R.B., Combrinck, S., and Viljoen, A. (2016). Myricetin: a dietary molecule with diverse biological activities. *Nutrients*, 8(2): 90.
- Shan, K., Wang, H., Xu, F., Luo, J., Li, Y., Han, A., Shao, X., and Wei, Y. (2019). Flavonoids from *Sonchus oleraceus* L. exert a hepatoprotective effect in hyperlipidemic mice by regulating blood lipid metabolism. (in Chinese with English abstract). *Food Science*, 40(19): 231–236.
- Singh, P. and Bast, F. (2015). Screening and biological evaluation of myricetin as a multiple target inhibitor insulin, epidermal growth factor, and androgen receptor; *in silico* and *in vitro*. *Investigational New Drugs*, 33(3): 575–593.
- Weigt, C., Hertrampf, T., Kluxen, F.M., Flenker, U., Huelsemann, F., Fritzemeier, K.H., and Diel, P. (2013). Molecular effects of ER alpha- and beta-selective agonists on regulation of energy homeostasis in obese female Wistar rats. *Molecular and Cellular Endocrinology*, 377(1–2): 147–158.
- Xia, S.-F., Qiu, Y.-Y., Chen, L.-M., Jiang, Y.-Y., Huang, W., Xie, Z.-X., Tang, X., and Sun, J. (2019). Myricetin alleviated hepatic steatosis by acting on microRNA-146b/thyroid hormone receptor b pathway in high-fat diet fed C57BL/6J mice. *Food and Function*, 10(3): 1465–1477.
- Yeh, C.W., Chen, W.J., Chiang, C.T., Lin-Shiau, S.Y., and Lin, J.K. (2003). Suppression of fatty acid synthase in MCF-7 breast cancer cells by tea and tea polyphenols: a possible mechanism for their hypolipidemic effects. *Pharmacogenomics Journal*, 3(5): 267–276.
- Ying, H.-Z., Zang, J.-N., Deng, L.-L., Wang, Z.-Y., and Yu, C.-H. (2013). Pentamethylquercetin reduces fat deposition via Sirt1-mediated pathways in male obese mice induced by a high fat diet. *Food and Chemical Toxicology*, 62: 463–469.
- Zechner, R., Kienesberger, P.C., Haemmerle, G., Zimmermann, R., and Lass, A. (2009). Adipose triglyceride lipase and the lipolytic catabolism of cellular fat stores. *Journal of Lipid Research*, 50(1): 3–21.



Zechner, R. and Langin, D. (2014). Hormone-sensitive lipase deficiency in humans. *Cell Metabolism*, 20(2): 199–201.

Zhang, W., Zhang, J., Jiang, Q., and Xia, W. (2013). The hypolipidemic activity of chitosan nanopowder prepared by ultrafine milling. *Carbohydrate Polymers*, 95(1): 487–491.

---

**Open Access statement.** This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium for non-commercial purposes, provided the original author and source are credited, a link to the CC License is provided, and changes – if any – are indicated.

