

Molecular profiles of small cell lung cancer subtypes: Therapeutic implications

Anna Schwendenwein,^{1,7} Zsolt Megyesfalvi,^{1,2,3,7} Nandor Barany,^{1,3,4} Zsuzsanna Valko,^{1,3} Edina Bugyik,³ Christian Lang,¹ Bence Ferencz,^{2,3} Sandor Paku,⁴ Andras Lantos,³ Janos Fillinger,^{2,3} Melinda Rezeli,⁵ Gyorgy Marko-Varga,⁵ Krisztina Bogos,³ Gabriella Galfy,⁶ Ferenc Renyi-Vamos,^{2,3} Mir Alireza Hoda,¹ Walter Klepetko,¹ Konrad Hoetzenecker,¹ Viktoria Laszlo,^{1,2,3,8} and Balazs Dome^{1,2,3,8}

¹Department of Thoracic Surgery, Comprehensive Cancer Center Vienna, Medical University of Vienna, 1090 Vienna, Austria; ²Department of Thoracic Surgery, National Institute of Oncology-Semmelweis University, 1122 Budapest, Hungary; ³National Koranyi Institute of Pulmonology, 1121 Budapest, Hungary; ⁴1st Department of Pathology and Experimental Cancer Research, Semmelweis University, 1085 Budapest, Hungary; ⁵Department of Biomedical Engineering, Lund University, 221 00 Lund, Sweden; ⁶Torokbalint County Institute of Pulmonology, 2045 Torokbalint, Hungary

Small cell lung cancer (SCLC; accounting for approximately 13%–15% of all lung cancers) is an exceptionally lethal malignancy characterized by rapid doubling time and high propensity to metastasize. In contrast to the increasingly personalized therapies in other types of lung cancer, SCLC is still regarded as a homogeneous disease and the prognosis of SCLC patients remains poor. Recently, however, substantial progress has been made in our understanding of SCLC biology. Advances in genomics and development of new preclinical models have facilitated insights into the intratumoral heterogeneity and specific genetic alterations of this disease. This worldwide resurgence of studies on SCLC has ultimately led to the development of novel subtype-specific classifications primarily based on the neuroendocrine features and distinct molecular profiles of SCLC. Importantly, these biologically distinct subtypes might define unique therapeutic vulnerabilities. Herein, we summarize the current knowledge on the molecular profiles of SCLC subtypes with a focus on their potential clinical implications.

Lung cancer, the leading cause of cancer-related deaths in the Western world, is classified into two major groups: small cell lung cancer (SCLC) and non-SCLC (NSCLC).¹ SCLC accounts for approximately 13%–15% of all lung cancers, and with a 5-year survival rate of less than 7%, it remains one of the most lethal forms of malignant diseases.^{2,3} It has a very aggressive course and is characterized by genomic instability, almost universal inactivation of the genes *TP53* and *RBI*, rapid tumor growth, increased vascularity, and high metastatic potential.^{4–6} Consequently, at the time of diagnosis, most SCLC patients already present with a metastatic spread outside the chest, which often leads to premature death.^{7,8} Most SCLC patients are current or former heavy smokers resulting in a high tumor mutational burden (TMB) (with C:G>A:T transversions being the most common type of base substitutions).^{9,10} Early detection strategies are mostly ineffective for SCLC even among high-risk populations, and there have been no significant improvements in survival and therapeutic approaches for more than 30 years, leading SCLC to be categorized as a “recalcitrant” cancer.^{5,11}

Platinum-based chemotherapy (CHT) in combination with etoposide and/or radiation therapy (RT) has been used in SCLC treatment and still remains the backbone for current combination strategies (Figure 1^{17–23}).^{16,24} Unlike NSCLC, which has an intrinsic tendency for CHT resistance, SCLC is initially highly sensitive to cytotoxic agents.²⁵ Even with response, however, SCLC frequently recurs within a short time span, and patients are seldom cured.²⁶ The most notable recent clinical progress in SCLC was the approval of the immune-checkpoint inhibitors (ICIs) atezolizumab, pembrolizumab, and nivolumab.^{27,28} Unfortunately, however, only 12.6% of patients remain progression-free at 1 year, and to date there are no reliable biomarkers predicting response to immune checkpoint blockade.^{27,29} With regard to other therapeutic approaches, due to the rapid doubling time of tumors and their high propensity to metastasize, surgery is rarely performed.³⁰ Accordingly, limited human SCLC tissue availability has increased the importance of preclinical models.^{5,31}

Unlike the increasingly personalized approach to clinical care of patients with other types of lung cancer, SCLC is still regarded as a “homogeneous” disease with a single morphological type.³² Consequently, current clinical study protocols for SCLC are generally based on disease stage with no consideration of predefining distinct molecular marker expressions that might have predictive or prognostic significance.^{5,33} However, recently, there has been a worldwide resurgence of studies on SCLC, including the development of new

<https://doi.org/10.1016/j.omto.2021.02.004>.

⁷These authors contributed equally

⁸Senior author

Correspondence: Zsolt Megyesfalvi, MD, Department of Thoracic Surgery, National Institute of Oncology-Semmelweis University, Rath Gyorgy utca 7–9, 1122 Budapest, Hungary.

E-mail: megyesfalvi.zsolt@semmelweis-univ.hu

Correspondence: Balazs Dome, MD, PhD, Department of Thoracic Surgery, Comprehensive Cancer Center, Medical University of Vienna, Waehringer Guertel 18–20, 1090 Vienna, Austria.

E-mail: balazs.dome@meduniwien.ac.at




Figure 1. Timeline of relevant therapeutic advances for small cell lung cancer (SCLC)

Initial therapeutic strategies for SCLC included surgery or radiotherapy alone. However, given the aggressive behavior and high metastatic potential of SCLC, systemic therapy with cytotoxic agents has rapidly become the cornerstone of management. In the 1940s, alkylating agents (such as nitrogen mustard) were used for the treatment of all bronchogenic carcinomas (including SCLC), resulting in tumor regression in more than 50% of patients. Yet, at that time, the true nature of SCLC was widely unknown, and all lung carcinomas were treated similarly. The first chemotherapeutic agent to show a statistically significant survival benefit for selected SCLC patients was cyclophosphamide, which doubled the survival when compared to best supportive care alone¹²⁻¹⁴. Despite the encouraging results with single-agent therapy, it became obvious early in the 1970s that combination therapy produces superior survival outcomes when compared to single-agent treatment^{14,15}. Therefore, during the late 1970s and early 1980s, cyclophosphamide was used in combination with other cytotoxic agents such as doxorubicin and vincristine (CAV). The basis of the currently used platinum-based combination chemotherapy (CHT) was defined during the mid-1980s when Evans et al.¹⁶ showed a clear survival benefit for patients treated with cisplatin plus etoposide. Since then, there have been no relevant advances in the standard-of-care CHT regimens, and the backbone for current combination strategies are still the platinum compounds. In more recent years, targeted therapy and immunotherapy have also been actively tested, leading to the approval of several immune-checkpoint inhibitors such as atezolizumab, pembrolizumab, and nivolumab.

preclinical models (such as patient-derived xenografts [PDXs] and genetically engineered mouse models [GEMMs]), comprehensive genomic profiling, and the identification of biologically distinct molecular subtypes.³⁴ Exploring the molecular profiles of SCLC subtypes might help to focus and accelerate therapeutic research. In this review, we systematically analyze the molecular heterogeneity of SCLC, mainly focusing on rationally targeted therapeutic implications and new treatment opportunities that may ultimately improve the clinical outcomes for patients with this devastating disease.

Tumor heterogeneity in SCLC

Neuroendocrine (NE) subtypes

Although clinically SCLC is still regarded as a single disease entity, preclinical studies from the past decades identified biologically different SCLC subgroups (Figure 2). Accordingly, SCLC can be classified today into NE-high and NE-low subtypes primarily based on the expression pattern of different NE markers such as chromogranin A (CHGA), synaptophysin (SYP), neural cell adhesion molecule 1 (NCAM1/CD56), and gastrin-releasing peptide (GRP).^{5,6,33,35} Additionally, some SCLCs lack NE differentiation and are termed as non-NE tumors.³³ The NE-high versus NE-low subtypes show major

differences in genetic alterations, morphology, growth properties, and immune infiltration.³⁵ NE-low SCLCs are furthermore associated with increased immune cell infiltration and referred to as “immune oasis” tumors, whereas NE-high SCLCs are characterized by low numbers of infiltrating immune cells and, consequently, have an “immune desert” phenotype.³⁶ This categorization has major clinical impact, as the two phenotypes are anticipated to respond differently to targeted therapeutics and ICIs.^{37,38}

Of note, another example of NE heterogeneity was described by Gazdar et al.^{39,40} in 1985 based on the *in vitro* and *in vivo* behavior of SCLC cells. The “classic” phenotype is associated with typical morphology, high expression of NE markers, and non-adherent growth pattern in cell cultures.^{5,39} In contrast, the “variant” phenotype is usually characterized by larger cells with prominent nucleoli, low expression of NE features, and an adherent or loosely adherent growth pattern *in vitro*.^{5,33,35}

Molecular subtypes

Recent SCLC profiling studies on cancer cell lines, PDXs, GEMMs, and primary human tumors suggest a model of distinct subtypes

SCLC subtype	Neuroendocrine				Non-neuroendocrine			
	ASCL1 "classic", NE-high		NEUROD1 "variant", NE-low		POU2F3		YAP1	
Marker expression	↑	↓	↑	↓	↑	↓	↑	↓
	BCL-2 DLL3 CD56 CHGA E-cadherin EZH2 GRP1 IGFBP5 INSM1 LSD1 L-MYC NFIB RET TTF-1 SOX2 SYP	MYC CREBBP Notch	ANTXR1 AURKA MYC HES6 INSM1 LSH1 NCAM NFIB TrkB	E-cadherin TTF1	AURKA ASCL2 AVIL MYC E-cadherin GFI1B IGF-1R SOX9 TRPM5 CHAT ATM	IGFBP5 INSM1 TTF1 SLFN11	AJUBA AURKA CDK4/6 Integrins Laminin mTOR PD-L1 PLK RB1 SOX9 TAZ	MYC INSM1
Potential therapeutic approaches	BCL-2 inhibitors DLL3 inhibitors HDAC inhibitors LSD1 inhibitors		AURKA inhibitors c-MYC inhibitors ADI-PEG 20 Seneca Valley virus		PARP inhibitors IGF-1R inhibitors Nucleoside analogues		Immune-checkpoint inhibitors mTOR inhibitors PLK inhibitors CDK4/6 inhibitors	

Figure 2. Tumor heterogeneity in SCLC with regard to neuroendocrine differentiation, molecular subtypes, and gene expression profile

Neuroendocrine (NE) differentiation can be defined by the expression pattern of different NE markers, including chromogranin A, synaptophysin, neural cell adhesion molecule 1, and gastrin-releasing peptide. However, a minority of SCLCs are negative for all standard NE markers. Additionally, SCLC can be subclassified according to the relative expression of four key transcriptional regulators: achaete-scute homolog 1 (ASCL1; also known as ASH1), neurogenic differentiation factor 1 (NEUROD1), yes-associated protein 1 (YAP1), and POU class 2 homeobox 3 (POU2F3). Various genetic alterations, biological properties, and thus potential therapeutic vulnerabilities are associated with these molecular subsets.

defined by the relative expression of four key transcriptional regulators: achaete-scute homolog 1 (ASCL1; also known as ASH1), neurogenic differentiation factor 1 (NEUROD1), yes-associated protein 1 (YAP1), and POU class 2 homeobox 3 (POU2F3).^{33,41} The association between these transcription factors (TFs) and the NE expression profile may provide subtype-specific therapeutic vulnerabilities (Figure 2).³³ SCLC-A (ASCL1) tumors show high expression of NE markers and classic morphology, compared to the NE-low SCLC-N (NEUROD1) subtype with variant morphology.³⁴ The transcription regulators ASCL1 and NEUROD1 both have been implicated as essential determinants of the developmental maturation of pulmonary NE cells (PNECs).⁴² In addition, SCLC-A and SCLC-N subtypes preferentially express the TF insulinoma-associated protein 1 (INSM1) as well.³³ By inhibiting the Notch signaling pathway, INSM1 also plays a key role in NE differentiation, and thus its expression levels are low in non-NE SCLCs.^{43–45} NEUROD1, ASCL1, and

INSM1 low-expressing or non-expressing SCLCs are classified either into SCLC-Y (YAP1) or SCLC-P (POU2F3) subtype, depending on their TF expression pattern.³³ YAP1 is a transcription regulator activated by the HIPPO signaling pathway.⁴⁵ Meanwhile, POU2F3 is required for the development of pulmonary tuft cells and chemosensory cells of the gastrointestinal epithelium.⁴⁶ Accordingly, recent data on high POU2F3 expression patterns detected in certain SCLCs suggest tuft cells as the origin of the SCLC-P subtype.^{33,46}

Genomic landscape of SCLC

SCLCs are likely to gain various genetic alterations as they evolve or as they metastasize outside the chest.⁴⁷ The metastatic potential of SCLC is often driven by the overexpression of nuclear factor I B (NFIB), which functions as an oncogene and is frequently amplified in metastases.⁴⁸ Likewise, lymph node (LN) metastases often undergo a change in NE expression patterns compared to the primary cancer,

thus enhancing the aggressive course of the disease.^{6,49} It is important to consider intratumoral heterogeneity in SCLC, as NE cancer cells are also capable of raising non-NE tumor cells that expedite the tumorigenesis and additionally contribute to CHT resistance.⁴⁷ Meanwhile, cancer stem cells (CSCs) are suspected to promote long-term tumor growth and affect the cellular heterogeneity of SCLC.⁴⁷ In this context, SOX2, the transcriptional regulator of pluripotent stem cells, is frequently overexpressed, especially in the SCLC-A subtype.⁵⁰

Notch signaling in SCLC is involved in NE differentiation.⁵¹ Tumor cells, which exhibit active Notch signaling, are slowly growing and often chemoresistant. Accordingly, Notch is considered as a tumor suppressor in SCLC.^{51,52} Based on the inhibitory activity of DLL3 on the Notch pathway or the inactivating mutations in Notch pathway genes, Notch is frequently inactivated in SCLCs with a high NE expression profile.⁶ Loss of NE differentiation and concomitant activation of Notch signaling is facilitated by the activation of the RE1 silencing transcription (REST) factor, a transcriptional repressor of NE and neuronal differentiation.⁵³ Notably, REST is absent in most NE SCLCs, also resulting in the inhibition of Notch signaling.⁵

NE-high SCLCs often harbor overexpression in one of the Myc family members, including c-MYC, l-MYC, and n-MYC.⁵⁴ The SCLC-A subtype is highly associated with the expression of l-MYC, whereas the upregulation of c-MYC is related to the SCLC-N subtype.³³ In addition, the SCLC-A subtype is suspected to lose ASCL1 expression as a result of c-MYC overexpression since high levels of c-MYC might contribute to the progression of SCLC cells from the classic phenotype to the NE-low variant phenotype.⁵⁵

It has also been hypothesized that MYC amplification appears during tumor progression and is connected to CHT resistance.^{54,56} Aurora kinase A and B (AURKA and AURKB) are serine/threonine kinases and have a main function in the regulation of mitosis.⁵⁷ The overexpression of AURKA promotes cell proliferation.⁵⁸ Meanwhile, AURKB phosphorylates RB1 and regulates the postmitotic checkpoints as well as prevents polyploidy after irregular mitosis.⁵⁵ The overexpression of aurora kinases in SCLC accompanied by Myc family amplification provides a growth advantage and causes polyploidy in SCLC.⁵⁹

The programmed cell death protein 1 (PD-1)/programmed death-ligand 1 (PD-L1) signaling pathway is a major therapeutic target in SCLC.⁶⁰ A considerable proportion of SCLCs exhibit aberrant PD-L1 expression on tumor cells that may be fundamental for moderate responses to immunotherapy.⁶¹ A subset of SCLCs is associated with the amplification of the fibroblast-growth factor receptor 1 (*FGFR1*) gene that displays a target for FGFR1 inhibitor therapy.⁶² Other unique features in SCLC include the expression of TFs SOX9 and ASCL2 and the receptor tyrosine kinase insulin-growth factor receptor 1 (IGF-1R) in POU2F3-expressing tumor cells.⁴⁶ Important aberrations in the SCLC-A subtype comprise amplifications of *BCL2*, *EZH2*, and the decrease of *CREBBP*, whereas SCLC-Y is associated

with mutations in the phosphatidylinositol 3-kinase (PI3K)/AKT/mTOR signaling pathway.⁵ Conclusively, intertumoral heterogeneity in SCLC has major therapeutic impacts.

Therapeutic implications

CHT

SCLC is initially highly sensitive to CHT and, accordingly, up to 75%–80% of all SCLCs first respond to platinum compounds.⁶³ However, the development of resistance is essentially universal, and patients are rarely cured. Chemorefractory tumor cells may arise due to the extensive TMB in SCLC and the coexisting subpopulations within a tumor.^{5,47,64} In addition, therapeutic outcomes might also be altered by the growth of resistant cell clones during disease progression.⁴⁷ Recent data suggest increased intratumoral heterogeneity after the onset of therapeutic resistance in SCLC.^{34,65} In contrast to NE-high SCLC cells (which are more sensitive to CHT because of their fast proliferation and high mitotic rate), the slower growing SCLC cells such as the previously mentioned Notch^{active} SCLC cells may be inherently chemoresistant.⁴⁷ Inactive Notch signaling triggers the NE cell differentiation during lung development, while active Notch signaling controls the non-NE cell fates.⁶⁶ Notch signaling further mediates the transition of NE to non-NE phenotype.⁶⁶ In SCLC, Notch signaling delivers context-dependent tumor-suppressive or oncogenic signals through its receptors.⁶⁶ MYC is also hypothesized to mediate NE plasticity in SCLC by activating Notch signaling.^{51,67} Furthermore, this interaction controls the dynamic behavior of cancer cells contributing to the co-existence of subtypes within a tumor.^{51,67}

The expression of Schlafen 11 (SLFN11) (a member of the Schlafen family involved in the control of cell proliferation and induction of the immune response) is observed to predict treatment responses to DNA-damaging agents such as cisplatin, etoposide, and poly(ADP-ribose) polymerase (PARP) inhibitors in SCLCs.^{34,68,69} SLFN11 expression, therefore, correlates with sensitivity to various DNA-damaging chemotherapeutics, but in many cases it is silenced in SCLC by methylation or acetylation.^{34,70} On the contrary, the upregulation of EZH2 mediates chemoresistance based on SLFN11 downregulation via histone methylation and modification.⁷¹ Preclinical data led to the hypothesis that therapeutic targeting of EZH2 might prolong and augment sensitivity to the CHT response.⁶⁰

Immunotherapy

ICIs have had a major impact on the clinical outcome of several solid tumors, including NSCLC, melanoma, and urothelial cancer.^{34,64,72,73} Epidemiological, biological, and clinical features of SCLC suggest that immunotherapy might be effective in this malignancy as well since many of the ICI susceptibility features in NSCLC are even more pronounced in SCLC.^{33,34,74} First, SCLC occurs almost exclusively in heavy smokers, and exposure to cigarette smoking is a predictive factor for responsiveness to ICIs in NSCLC.^{75,76} Second, compared to NSCLCs, which exhibit a lower TMB of 6.3–9 mutations/Mb, SCLCs display a higher median TMB of 9.9 mutations/Mb and lack the recurrent driver alterations in *EGFR* or *ALK* that correlate with poor response to immunotherapy.^{24,77} Third, SCLCs can also

Table 1. Completed clinical trials evaluating the safety and efficacy of immune-checkpoint inhibitors in SCLC patients

Study name	Study phase	Mechanism of action	Agent	Outcomes
First-line therapy				
IMpower133 ²⁷	I/III	PD-L1 inhibitor	atezolizumab	safety: well tolerated ORR: 60.2% versus 64.4% PFS: ^a 5.2 versus 4.3 (p = 0.02) OS: ^a 12.3 versus 10.3 (p = 0.007)
CASPIAN ⁸⁰	III	PD-L1 inhibitor	durvalumab	safety: well tolerated ORR: 79% versus 70% PFS: 5.1 versus 5.4 (p = ns) OS: 13.0 versus 10.3 (p = 0.004)
KEYNOTE-604 ¹³⁸	III	PD-1 inhibitor	pembrolizumab	safety: well tolerated ORR: 70.6% versus 61.8% 12-month PFS: 13.6% versus 3.1% (p = 0.0023) 24-month OS: 22.5% versus 11.2% (p = 0.0164)
REACTION ¹³⁹	II	PD-1 inhibitor	pembrolizumab	safety: well tolerated ORR: 61% PFS: 4.7 versus 5.4 OS: 12.3 versus 10.4
ClinicalTrials.gov: NCT01450761 ¹⁴⁰	III	CTLA-4 inhibitor	ipilimumab	safety: well tolerated ORR: 62% versus 62% PFS: 4.6 versus 4.4 (p = 0.016) OS: 11.0 versus 10.9 (p = 0.377)
Maintenance				
CheckMate 451 ¹⁴¹	III	PD-1 and CTLA-4 inhibitors	nivolumab plus ipilimumab	non-significant results
ClinicalTrials.gov: NCT02359019 ¹⁴²	II	PD-1 inhibitor	Pembrolizumab	safety: well tolerated ORR: 11.1% PFS: 1.4 OS: 9.6
Recurrent SCLC				
CheckMate-032 ¹⁴³	I/II	PD-1 and CTLA-4 inhibitors	nivolumab/ nivolumab plus ipilimumab ^b	safety: manageable safety profile ORR: 10% versus 23% versus 19% PFS: 1.4 versus 2.6 versus 1.4 OS: 4.4 versus 7.7 versus 6.0
BIOLUMA ^c ¹⁴⁴	II	PD-1 and CTLA-4 inhibitors	nivolumab plus ipilimumab	safety: high toxicity rates ORR: 38.8%
CheckMate-331 ¹⁴⁵	III	PD-1 inhibitor	nivolumab	safety: well tolerated ORR: 14% versus 16% PFS: 1.4 versus 3.8 OS: 7.5 versus 8.4
KEYNOTE-028 ¹⁴⁶	Ib	PD-1 inhibitor	pembrolizumab	safety: well tolerated ORR: 33.3% PFS: 1.9 OS: 9.7
KEYNOTE-158 ¹⁴⁷	II	PD-1 inhibitor	pembrolizumab	safety: well tolerated ORR: 18.7%

(Continued on next page)

Table 1. Continued

Study name	Study phase	Mechanism of action	Agent	Outcomes
				PFS: 2.0
				OS: 9.1
				safety: well tolerated
ClinicalTrials.gov: NCT02261220 ¹⁴⁸	I/II	PD-L1 and CTLA-4 inhibitors	durvalumab plus tremelimumab	ORR: 13.3%
				PFS: 1.8
				OS: 7.9

PD-1, programmed cell death protein 1; PD-L1, programmed death-ligand 1; CTLA-4, cytotoxic T lymphocyte-associated protein 4; ORR, objective response rate; PFS, progression-free survival; OS, overall survival.

^aIn months.

^bNivolumab plus ipilimumab combination consisted of nivolumab (1 mg/kg) + ipilimumab (3 mg/kg) OR nivolumab (3 mg/kg) + ipilimumab (1 mg/kg).

^cOnly patients with high tumor mutation burden were included.

spontaneously provoke a strong immune response based on the segregation of humoral or cellular components that are summarized as paraneoplastic syndromes (PNSs).⁷⁸ However, despite the above-mentioned susceptibility features, the latest clinical trials regarding the treatment of SCLC with ICIs have shown only modest improvement both in progression-free survival (PFS) and overall survival (OS) (Table 1).⁷⁴

The IMpower133 trial evaluated the efficacy and safety of the anti-PD-L1 antibody conjugate atezolizumab in combination with carboplatin and etoposide in treatment-naïve patients diagnosed with advanced SCLC.^{60,79} The investigators found that although the addition of atezolizumab to CHT significantly prolonged the PFS and OS of these patients (versus the placebo group; OS and PFS were 12.3 versus 10.3 months and 5.2 versus 4.3 months, respectively), the gain in survivals was relatively modest compared to those achieved in NSCLC.⁷⁹ Nevertheless, results of this trial represent the first significant improvement in systemic therapy for untreated SCLC patients in the last 30 years. Accordingly, the IMpower133 study has prompted the US Food and Drug Administration (FDA) and the European Medicines Agency (EMA) to approve atezolizumab in this setting.²⁴

Notably, similar results were observed in the CASPIAN study, where first-line durvalumab plus platinum-etoposide resulted in moderately improved OS and overall response rate (ORR) (versus a clinically relevant control group; OS rates were 13 versus 10.3 months, and ORRs were 79% versus 70%, respectively).⁸⁰ Importantly, however, there were no significant differences in PFS (5.1 versus 5.4 months).⁸⁰

Pembrolizumab is an anti-PD-1 antibody approved for metastatic SCLC patients with disease progression after platinum-based CHT and at least one other line of therapy.⁸¹ Results from the KEYNOTE-028 and KEYNOTE-158 studies revealed antitumor activity in a subset of patients with recurrent SCLC.⁸¹ The KEYNOTE-158 study enrolled patients regardless of their PD-L1 status, whereas the KEYNOTE-028 study only included patients with PD-L1-positive SCLCs. Based on the median OS of 9.7 (KEYNOTE-028) and 9.1

(KEYNOTE-158) months in patients treated with pembrolizumab, this agent constitutes a treatment option for recurrent SCLC as third- or subsequent-line therapy.⁸¹ The CheckMate 032 clinical trial evaluated the efficacy of nivolumab alone or in combination with ipilimumab.^{28,82} The patients eligible for the study received platinum-based CHT and at least one other agent, but they still showed a rapid disease progression. In addition, the study also demonstrated an improved cytotoxic activity for combined cytotoxic T lymphocyte-associated protein 4 (CTLA-4) (ipilimumab)- and PD-1 (nivolumab)-targeted antibodies.⁸² Of note, however, third-line therapy for metastatic SCLC patients with nivolumab monotherapy demonstrated a durable response in a subgroup of SCLC patients regardless of their PD-L1 status.^{28,82} Accordingly, in 2018, the FDA granted accelerated approval to nivolumab for patients with metastatic SCLC based on the results of the CheckMate 032 clinical trial.²⁸

All in all, although some data suggest that SCLC should be more vulnerable to immunotherapy than NSCLC, the available clinical results are less compelling. Therefore, there is an unmet need to identify the determinants of ICI activity in SCLC that might differ from those in NSCLC and other solid tumors.⁵⁷ Immune phenotypes as well as the NE and molecular subtypes may provide a better understanding of the underlying properties of SCLC patients. NE-high tumors with the immune desert phenotype are associated with decreased immune cell (CD8⁺ effector T cells primarily) infiltration compared to NE-low tumors.^{36,38} Expressions of indoleamine 2,3-dioxygenase (IDO) and poliovirus receptor (PVR), which are both important factors of the SCLC immune microenvironment, are also significantly lower in NE-high tumors (versus NE-low tumors).^{38,83–85} In addition, these tumors are associated with low T cell immunoglobulin and mucin domain-3 (TIM3) levels as well.^{38,86} Of note, TIM3 is a specific marker of lymphocyte exhaustion and is one of the most promising immune-checkpoint targets.⁸⁶ Altogether, patients with NE-high tumors are less likely to respond to ICIs.^{36,38} Another possible explanation for the poor response rates to immunotherapy in SCLC may be that PD-L1 expression is much lower in SCLC compared to other solid tumors, and cancer cell PD-L1 expression seemingly does not correlate with ICI efficacy.^{34,87} Furthermore, due to the suppressed

Table 2. Summary of clinical trials evaluating the safety and/or efficacy of targeted agents in advanced-stage SCLC patients

Study name	Study phase	Mechanism of action	Agent	Outcomes
ClinicalTrials.gov: NCT02289690 ⁹⁵	I/II	PARP inhibitor	veliparib	safety: well tolerated
ClinicalTrials.gov: NCT01286987 ⁹⁶	I	PARP inhibitor	talazoparib	safety: well tolerated
ClinicalTrials.gov: NCT01638546 ⁹⁷	II	PARP inhibitor	veliparib	ORR: 39% versus 14% (p = 0.016) PFS: ^a 3.8 versus 2.0 (p = 0.39) OS: ^a 82 versus 7.0 (p = 0.50)
ECOG-ACRIN 2511 ⁹⁸	II	PARP inhibitor	veliparib	ORR: 71.9% versus 65.6% (p = 0.57) PFS: 6.1 versus 5.5 (p = 0.06) OS: 10.3 versus 8.9 (p = 0.17)
TAHOE ⁹⁹	III	DLL3-targeted antibody drug conjugate	Rova-T	non-significant results
TRINITY ¹⁰⁰	II	DLL3-targeted antibody drug conjugate	Rova-T	non-significant results
ALTER 1202 ¹⁰¹	II	tyrosine kinase inhibitor	anlotinib	PFS: 4.1 versus 0.7 (p < 0.0001) OS: 7.3 versus 4.9 (p = 0.0210)
ClinicalTrials.gov: NCT00154388 ¹⁰²	II	tyrosine kinase inhibitor	imatinib	non-significant results
ClinicalTrials.gov: NCT01533181 ¹⁰³	II	IGF-R1 inhibitor	linsitinib	non-significant results
ClinicalTrials.gov: NCT00869752 ¹⁰⁴	I	IGF-R1 inhibitor	dalotuzumab	safety: well tolerated ORR: 22% versus 18% (p = 0.406)
ClinicalTrials.gov: NCT02038647 ¹⁰⁵	II	AURKA inhibitor	alisertib	PFS: 3.32 versus 2.17 (p = 0.113) OS: 6.86 versus 5.58 (p = 0.714)
SALUTE ¹⁰⁶	II	VEGF inhibitor	bevacizumab	ORR: 58% versus 48% PFS: 5.5 versus 4.4 OS: 9.4 versus 10.9
ClinicalTrials.gov: NCT02454972 ¹⁰⁷	II	RNA polymerase II inhibitor	lurbinectedin	ORR: 35.2% PFS: 3.5 OS: 9.3

PARP, poly(ADP-ribose) polymerase; DLL3, delta-like protein 3; IGF-R1, insulin-like growth factor 1 receptor; AURKA, aurora kinase A; VEGF, vascular endothelial growth factor.

^aIn months.

expression of major histocompatibility complex class I (MHC class I) in the tumor microenvironment, antigen presentation might also be defective.⁸⁸ Thus, the alterations affecting the antigen presentation by MHC molecules may contribute to escape from T cell recognition and destruction.^{34,87,88} Expressions of HLA-A, HLA-B, HLA-C, and β_2 -microglobulin are also significantly lower in SCLC cell lines (versus NSCLC), and, consequently, these cell lines are less immunogenic when injected into immunocompetent mice.^{34,88–90} Finally, the specific clinical features of SCLC might also affect the beneficial use of ICIs.⁷⁴ Most SCLC patients often require prolonged steroid therapy due to superior vena cava syndrome or brain metastases.⁷⁴ Chronic steroids, however, are a known limitation for immunotherapy.^{74,91}

Immunotherapy-related adverse events (irAEs) represent a major concern in SCLC.⁷⁴ irAEs mostly include inflammatory or autoimmune complications with sometimes severe sequelae for patients.^{74,92} Although the exact pathophysiological mechanism of irAEs has not been fully uncovered in SCLC, it is suspected that the genetic predisposition and the latent (i.e., clinically asymptomatic) PNSs might play a key role.⁹² One possible mechanism is a T cell-mediated reaction to shared antigens that are expressed both in tumors and inflammatory lesions.⁹² Additionally, the development of autoantibodies might also contribute to the appearance of irAEs, just as the overactivation of innate and adaptive immune cells, which lead to increased cytokine secretion.⁹² Lastly, the role of the gut mucosal immune system and the gut microbiome in irAEs has also been intensively investigated.^{93,94}

Targeted therapy

Several targeted agents are being tested for the treatment of SCLC (Table 2). However, to advance these treatment options, it is necessary to better understand the molecular alterations that have been already described in SCLCs.^{5,6,108}

Compared to other lung cancer subtypes and normal lung epithelial cells, SCLC cells show a high PARP expression profile and are highly sensitive to PARP inhibitors.^{98,109} In addition, PARP inhibitors also enhance the effects of CHT and ionizing radiation both *in vivo* and *in vitro*.^{109,110} Therefore, the use of PARP inhibitors might be a promising targeted therapeutic approach in SCLC patients. The phase II clinical trial ECOG-ACRIN 2511 investigated the efficacy of the PARP inhibitor veliparib in untreated, advanced-stage SCLC patients.⁹⁸ The investigators found that although patients treated with veliparib in combination with cisplatin and etoposide doublet had an improved OS compared to the control group (OSs were 10.3 versus 8.9 months), the results were not statistically significant.⁹⁸ PARP inhibitor combinations might thus be attractive therapeutic approaches for SCLC patients, but predictive biomarkers are required to maximize their clinical efficacy. Of note, expression of the already mentioned SLFN11 strongly correlates with veliparib efficacy and may represent a potential biomarker for these patients.⁹⁷

The delta-like ligand 3 (DLL3) is highly expressed in a subset of SCLCs with NE origin.²⁴ The antibody-drug conjugate rovalpituzumab




Figure 3. Potential novel therapeutic approaches in SCLC

Subtype-specific potential therapeutic targets for SCLC-A, SCLC-N, SCLC-P, and SCLC-Y subtypes are highlighted in yellow (BCL2 and DLL3), gray (c-MYC and AURKA), red (PARP and IGF-1R), and green (CDK4/6 and PD-L1), respectively. The association between the aforementioned molecular subtypes and the potentially targetable molecules highlighted in blue (PDGFR, VEGFR, FGFR, CTLA-4 and PD-1), is currently unknown. The figure was created with BioRender.

tesirine (Rova-T) binds DLL3 on these target-expressing cells to induce cell death.²⁴ Rova-T is the first targeted therapeutic agent in SCLC to use DLL3 as a novel biomarker.¹⁰⁰ The phase II TRINITY study investigated the efficacy and safety of Rova-T as a third-line agent in relapsed and refractory SCLC patients.¹⁰⁰ The study results revealed modest anti-tumor activity of Rova-T, which also caused toxicity.¹⁰⁰ The Rova-T development program with reference to the phase III MERU trial was considered ineffective in case of unselected patients and was terminated by the manufacturer.¹¹¹ Notably, although the results show a lack of survival benefit, Rova-T may still be a promising therapeutic approach for properly selected SCLC patients.

The phase II ClinicalTrials.gov: NCT01045421 study tested the activity and safety of alisertib in patients with relapsed or refractory SCLC or with other tumors.^{60,112} Patients eligible for the study had to have undergone two or fewer previous cytotoxic regimens. The phase II clinical trial of alisertib as a monotherapy or in combination with other agents in multiple tumor types showed antitumor activity and provides a therapeutic strategy in relapsed SCLCs.¹¹²

Surgery

Several retrospective observational studies and cancer registries have previously provided encouraging long-term results in patients who underwent surgical resection for early stage SCLC.^{113–117} Unfortunately, current screening and diagnostic approaches clearly fail to

identify early stage SCLC patients who might be eligible for surgery.¹¹⁸ Surgery is thus rarely performed in SCLC and approximately 80%–85% of SCLC patients are being diagnosed with extensive disease.¹¹⁹ Therefore, there is an urgent clinical need for novel biomarkers with the potential of enabling early SCLC diagnosis and of improved selection of surgical candidates. Recent data have demonstrated for example the predominance of the ASCL1 subtype in early SCLC lesions.^{42,55} Importantly, novel targeted drugs may also find their way to the armamentarium of SCLC therapies in the neoadjuvant setting. The development of such neoadjuvant approaches could even allow to offer surgery for SCLC patients with initially more advanced

disease. Thus, a better understanding of the molecular subtypes might revolutionize the role of surgery in SCLC and raise the hope for better outcomes in the future.

Targeted therapy with regard to NE and molecular subtypes

Targeted therapies for SCLC have so far failed, and the success of immunotherapy in NSCLC has not been reflected in SCLC.¹²⁰ One major reason for these relatively disappointing results lies behind the heterogeneity of SCLC.^{33,47} Unlike in NSCLC therapy and clinical studies, SCLC patients are still enrolled in clinical trials irrespective of their molecular background.³³ Accordingly, identification of subtype-specific molecular profiles and clinically meaningful biomarkers may contribute to novel targeted strategies in SCLC (Figure 3).

SCLC-A

The SCLC-A subtype is anticipated to respond to DLL3-targeted antibody drug conjugates due to the direct transcriptional interaction of DLL3 with *ASCL1* in Notch^{inactive} tumor cells.^{121,122} Accordingly, treatment with the DLL3-targeted antibody drug conjugate Rova-T defines a subtype-specific therapy for SCLC-A.

BCL2 is another direct transcriptional target of *ASCL1*, and the high expression levels are suggestive of potential therapeutic benefit from the *BCL2* inhibitor venetoclax.

The histone demethylase LSD1 activity has been described to be dependent on the disruption of INSM1, which is linked to the NE subtypes SCLC-A and SCLC-N.¹²³ Recent data also indicate that LSD1 inhibition leads to NOTCH1 activation, resulting in ASCL1 suppression in SCLC.¹²⁴

In addition, the SCLC-A subtype is also associated with CREBBP inactivation and thus with increased sensitivity to histone deacetylase (HDAC) inhibitors (e.g., pracinostat).¹²⁵

SCLC-N

The SCLC-N subtype is frequently associated with MYC amplification, which serves as a potential target for therapeutic agents.⁵⁶ Increased AURKA activity and arginine biosynthesis are also characteristic features of this particular subtype.⁵⁵ Accordingly, the combination therapy of both AURKA (e.g., alisertib) and c-MYC inhibitors is hypothesized to enhance the therapeutic efficacy.^{33,126} Additionally, the SCLC-N (i.e., “variant”) subtype exhibits a selective tropism for the oncolytic Seneca Valley virus (SVV),¹²⁷ which infects and eliminates the NE cancer cells via lysis. Therefore, with appropriate biomarker-guided patient selection, the SVV oncolytic virus may have selective efficacy either as single agent therapy or in combination with immunotherapy.^{33,127,128} In this context, the NEUROD1-to-ASCL1 ratio may function as a predictive biomarker.¹²⁷ Finally, based on recent *in vivo* studies, “NEUROD1-high” tumor cells are also suspected to be sensitive to arginine depletion caused by pegylated arginine deaminase (ADI-PEG 20), which leads to the inhibition of tumor cell growth.^{33,54}

SCLC-P

Based on the results of CRISPR screens, the SCLC-P subtype possesses a unique vulnerability to IGF-1R deficiency.⁴⁶ This leads to the hypothesis that IGF-1R inhibitors (e.g., dalotuzumab) may serve as potential specific therapeutic agents for these patients.⁴⁶ PARP inhibitors (e.g., veliparib) are also suspected to be most effective in this molecular subtype, although, to the best of our knowledge, SLFN11 expression does not correlate with the subtype-specific transcriptional regulators.¹²⁹ Lastly, we mention that the SCLC-P subtype might be sensitive to nucleoside analogues as well.¹³⁰

SCLC-Y

PD-1 or PD-L1 expressions are not subtype-specific but may preferentially be linked to the SCLC-Y subtype since YAP1 has been shown to upregulate PD-L1 transcripts and induce an immunosuppressive tumor microenvironment.^{131,132} In addition, SCLC-Y tumor cells have higher expression levels of both CD38 and LAG-3 and, consequently, they show a higher likelihood to respond to ICIs.¹³¹ Based on gene expression and recent *in silico* results, the SCLC-Y subtype is also the most sensitive to mTOR, PLK, and potentially to CDK4/6 inhibitors.^{45,133}

Metabolic pathways may also provide new targets for the treatment of SCLC. Recent data described an iron-dependent type of regulated necrosis called ferroptosis.¹³⁴ Non-NE SCLCs were demonstrated to be selectively sensitive to induced ferroptosis.¹³⁴ In contrast, high-NE

SCLCs are resistant to induced ferroptosis, but respond exquisitely to thioredoxin (TRX) pathway inhibition.¹³⁴ A combination of ferroptosis induction and TRX pathway inhibition may thus provide a treatment regimen for intratumoral NE/non-NE heterogeneous SCLC.¹³⁴

Open questions

What is the clinical relevance of intratumoral and intertumoral heterogeneity in SCLC?

The molecular characterization of SCLC is still of particular interest, although there have been major steps forward in identifying the exact pathogenesis and developing novel therapeutic approaches.^{6,33,47} SCLC is not a single entity as was thought before, but it displays heterogeneity in multiple ways. Significant intratumoral and intertumoral heterogeneity in SCLC was shown at the level of molecular diversity and NE differentiation. Bronchoscopic examination of the tumor may not be sufficient to determine the histological features and mutational landscape due to a possible presence of multiple subtypes within a tumor.^{51,135}

Do NE and molecular subtypes correlate with different prognosis and clinical implications?

Genetic alterations lead to the development of SCLC subtypes and therapeutic resistance. The classification of NE (NE-high, NE-low, and non-NE) and molecular subtypes (ASCL1, NEUROD1, YAP1, and POU2F3) in recent years improved our understanding of SCLC. Recent clinical data demonstrate that ASCL1 overexpression might be a negative prognostic indicator in early stage resected SCLC patients.¹²² With regard to therapeutic implications, however, the SCLC-A subtype is suspected to be more chemosensitive compared to the SCLC-N subtype, since most SCLC-A cell lines are derived from treatment-naïve patients whereas the SCLC-N cell lines mostly originate from post-treatment patients.⁴⁰ As for the SCLC-Y subtype, YAP1 expression is linked with poor prognosis and decreased survival plus increased chemoresistance.⁴⁵ These hypotheses are widely disputed, and the SCLC behavior in a clinical context is still not clarified.

What is the significance of biological plasticity between subtypes?

The presence of multiple subpopulations within a tumor and the biological plasticity between the subtypes might contribute to therapeutic outcomes. Recent preclinical studies suggest a possible hierarchy between subtypes, with SCLC-A being a necessary precursor of SCLC-N.^{33,42,55} This tumor evolution might greatly influence the response rates, as some tumor subpopulations may escape from therapy.³³ Of note, however, the success of targeted therapies in cancer treatment is impaired by other mechanisms of resistance as well. Increased TMB potentially leads to a higher chance of developing drug resistance just as the amplification of the transcriptional regulator NFIB, driving tumor initiation, progression, and metastasis of SCLC.⁵

Do metabolic pathways in each subtype have therapeutic impact?

Beside the molecular background in SCLC, metabolic pathways in each particular subtype also influence the tumor behavior and

therapeutic response. Ferroptosis and arginine depletion have recently been investigated to become targets for subtype-specific therapy.^{54,134} MYC-driven SCLC cells are dependent on different arginine-regulated pathways.⁵⁴ Non-NE SCLCs are reported to be specifically sensitive to induced ferroptosis.¹³⁴ Addressing selective cell death and metabolic pathways in SCLC subtypes may help to identify subtype-specific vulnerabilities for targeted therapies.¹³⁴

Do SCLC subtypes display different metastatic potential and organotropism?

The NE pattern of LN or organ metastases might not reflect that of the primary tumor.⁴⁹ Therefore, the resulting discordance between the primary tumor and metastases may result in the partial efficacy of therapeutic agents. Metastases of SCLC are observed to have a preference for certain organs. Very common sites for metastases comprise brain, bone, liver, and adrenal glands.¹³⁶ Additionally, SCLC cells are suspected to arise from different cells of origin. The definition of the distinct precursor cells may reveal biomarkers, which help to understand the early events of tumorigenesis and predict the tumor evolution.¹³⁷

Conclusions

In contrast to NSCLC, where genotype-based targeted therapies have dramatically improved the treatment outcomes in patients with advanced stage disease, the therapy options in SCLC are still limited and the survival rates are dismal. No significant progress has been made in the systemic treatment of SCLC in the last three decades, mainly due to the high plasticity of SCLC and also to the non-selected patient groups in clinical trials. Recent research defined the heterogeneity in SCLC, but further exploration of the nature of SCLC subtypes is needed to interpret their similarities, diversities, and respective behavior. Defining the distinct gene expression profiles (*ASCL1*, *NEUROD1*, *POU2F3*, and *YAP1*) of SCLC patients will be fundamental to choose the most effective therapy. Because immunotherapies, biomarker-directed therapies, and chemotherapies operate on different targets and mechanisms, a combined or synergistic treatment may increase the therapeutic effects. Conclusively, the development of new drugs and a combination of different subtype-specific therapies are substantial to fight this deadly disease.

ACKNOWLEDGMENTS

B.D. acknowledges funding from the Hungarian National Research, Development and Innovation Office (KH130356, K129065, and KNN121510). B.D. and V.L. were also supported by the Austrian Science Fund (FWF I3977 and FWF I4677 to B.D. and FWF I3522 to V.L.). V.L. is a recipient of Janos Bolyai Research Scholarship of the Hungarian Academy of Sciences and the New National Excellence Program of the Ministry for Innovation and Technology (UNKP-19-4). Z.M. was supported by the New National Excellence Program of the Ministry for Innovation and Technology (UNKP-20-3).

AUTHOR CONTRIBUTIONS

Conceptualization, A.S., Z.M., V.L., and B.D.; methodology, A.S., Z.M., S.P., A.L., J.F., M.R., G.M.-V., K.B., G.G., F.R.-V., M.A.H., K.H., and W.K.; investigation, A.S., Z.M., N.B., Z.V., E.B., C.L., B.F.,

V.L., and B.D.; writing – original draft, A.S., Z.M., V.L., and B.D.; writing – review & editing, all authors; funding acquisition, Z.M., V.L., and B.D.; resources, F.R.-V., W.K., V.L., and B.D.; supervision, L.V. and B.D.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

- Blandin Knight, S., Crosbie, P.A., Balata, H., Chudziak, J., Hussell, T., and Dive, C. (2017). Progress and prospects of early detection in lung cancer. *Open Biol.* 7, 170070.
- Kahnert, K., Kauffmann-Guerrero, D., and Huber, R.M. (2016). SCLC—state of the art and what does the future have in store? *Clin. Lung Cancer* 17, 325–333.
- Karachaliou, N., Pilotto, S., Lazzari, C., Bria, E., de Marinis, F., and Rosell, R. (2016). Cellular and molecular biology of small cell lung cancer: an overview. *Transl. Lung Cancer Res.* 5, 2–15.
- Siegel, R.L., Miller, K.D., and Jemal, A. (2020). Cancer statistics, 2020. *CA Cancer J. Clin.* 70, 7–30.
- Gazdar, A.F., Bunn, P.A., and Minna, J.D. (2017). Small-cell lung cancer: what we know, what we need to know and the path forward. *Nat. Rev. Cancer* 17, 725–737.
- George, J., Lim, J.S., Jang, S.J., Cun, Y., Ozretić, L., Kong, G., Leenders, F., Lu, X., Fernández-Cuesta, L., Bosco, G., et al. (2015). Comprehensive genomic profiles of small cell lung cancer. *Nature* 524, 47–53.
- Matthews, M.J., Kanhouwa, S., Pickren, J., and Robinette, D. (1973). Frequency of residual and metastatic tumor in patients undergoing curative surgical resection for lung cancer. *Cancer Chemother. Rep.* 3 4, 63–67.
- Carter, B.W., Glisson, B.S., Truong, M.T., and Erasmus, J.J. (2014). Small cell lung carcinoma: staging, imaging, and treatment considerations. *Radiographics* 34, 1707–1721.
- Pleasance, E.D., and Stephens, P.J. (2010). A small-cell lung cancer genome with complex signatures of tobacco exposure. *Nature* 463, 184–190.
- Alexandrov, L.B., Ju, Y.S., Haase, K., Van Loo, P., Martincorena, I., Nik-Zainal, S., Totoki, Y., Fujimoto, A., Nakagawa, H., Shibata, T., et al. (2016). Mutational signatures associated with tobacco smoking in human cancer. *Science* 354, 618–622.
- US Congress (2012). H.R.733—Recalcitrant cancer research act of 2012, <https://www.congress.gov/bill/112th-congress/house-bill/733>.
- Green, R.A., Humphrey, E., Close, H., and Patno, M.E. (1969). Alkylating agents in bronchogenic carcinoma. *Am. J. Med.* 46, 516–525.
- Chan, B.A., and Coward, J.I.G. (2013). Chemotherapy advances in small-cell lung cancer. *J. Thorac. Dis.* 5 (Suppl 5), S565–S578.
- Haddadin, S., and Perry, M.C. (2011). History of small-cell lung cancer. *Clin. Lung Cancer* 12, 87–93.
- Einhorn, L.H., Fee, W.H., Farber, M.O., Livingston, R.B., and Gottlieb, J.A. (1976). Improved chemotherapy for small-cell undifferentiated lung cancer. *JAMA* 235, 1225–1229.
- Evans, W.K., Osoba, D., Feld, R., Shepherd, F.A., Bazos, M.J., and DeBoer, G. (1985). Etoposide (VP-16) and cisplatin: an effective treatment for relapse in small-cell lung cancer. *J. Clin. Oncol.* 3, 65–71.
- Eckardt, J., Gralla, R., Palmer, M.C., Gandara, D., Laplante, J., Sandler, A., Fields, S.Z., Fitts, D., and Broom, C. (1996). Topotecan (T) as second-line therapy in patients (Pts) with small cell lung cancer (SCLC): a phase II study. *Ann. Oncol.* 7 (Suppl 5), 107.
- Aupérin, A., Arriagada, R., Pignon, J.P., Le Péchoux, C., Gregor, A., Stephens, R.J., Kristjansen, P.E., Johnson, B.E., Ueoka, H., Wagner, H., and Aisner, J.; Prophylactic Cranial Irradiation Overview Collaborative Group (1999). Prophylactic cranial irradiation for patients with small-cell lung cancer in complete remission. *N. Engl. J. Med.* 341, 476–484.
- Samantas, E., Skarlos, D.V., Pectasides, D., Nicolaides, P., Kalofonos, H., Mylonakis, N., Vardoulakis, T.h., Kosmidis, P., Pavlidis, N., and Fountzilas, G. (1999).

- Combination chemotherapy with low doses of weekly carboplatin and oral etoposide in poor risk small cell lung cancer. *Lung Cancer* 23, 159–168.
20. Okamoto, H., Watanabe, K., Nishiwaki, Y., Mori, K., Kurita, Y., Hayashi, I., Masutani, M., Nakata, K., Tsuchiya, S., Isobe, H., and Saijo, N. (1999). Phase II study of area under the plasma-concentration-versus-time curve-based carboplatin plus standard-dose intravenous etoposide in elderly patients with small-cell lung cancer. *J. Clin. Oncol.* 17, 3540–3545.
 21. 2007). FDA approves oral topotecan for relapsed small-cell lung cancer. *Oncology* 21. <https://www.cancernetwork.com/view/fda-approves-oral-topotecan-relapsed-small-cell-lung-cancer>.
 22. Eckardt, J.R., von Pawel, J., Pujol, J.L., Papai, Z., Quoix, E., Ardizzoni, A., Poulin, R., Preston, A.J., Dane, G., and Ross, G. (2007). Phase III study of oral compared with intravenous topotecan as second-line therapy in small-cell lung cancer. *J. Clin. Oncol.* 25, 2086–2092.
 23. Yao, J.C., Fazio, N., Singh, S., Buzzoni, R., Carnaghi, C., Wolin, E., Tomasek, J., Raderer, M., Lahner, H., Voi, M., et al.; RAD001 in Advanced Neuroendocrine Tumours, Fourth Trial (RADIANT-4) Study Group (2016). Everolimus for the treatment of advanced, non-functional neuroendocrine tumours of the lung or gastrointestinal tract (RADIANT-4): a randomised, placebo-controlled, phase 3 study. *Lancet* 387, 968–977.
 24. Zhang, C., Leigh, N.B., Wu, Y.L., and Zhong, W.Z. (2019). Emerging therapies for non-small cell lung cancer. *J. Hematol. Oncol.* 12, 45.
 25. Rossi, A., Di Maio, M., Chiodini, P., Rudd, R.M., Okamoto, H., Skarlos, D.V., Fröh, M., Qian, W., Tamura, T., Samant, E., et al. (2012). Carboplatin- or cisplatin-based chemotherapy in first-line treatment of small-cell lung cancer: the COCIS meta-analysis of individual patient data. *J. Clin. Oncol.* 30, 1692–1698.
 26. Gong, J., and Salgia, R. (2018). Managing patients with relapsed small-cell lung cancer. *J. Oncol. Pract.* 14, 359–366.
 27. Horn, L., Mansfield, A.S., Szczesna, A., Havel, L., Krzakowski, M., Hochmair, M.J., Huemer, F., Losonczy, G., Johnson, M.L., Nishio, M., et al.; IMPower133 Study Group (2018). First-line atezolizumab plus chemotherapy in extensive-stage small-cell lung cancer. *N. Engl. J. Med.* 379, 2220–2229.
 28. Ready, N., Farago, A.F., de Braud, F., Atmaca, A., Hellmann, M.D., Schneider, J.G., Spigel, D.R., Moreno, V., Chau, I., Hann, C.L., et al. (2019). Third-line nivolumab monotherapy in recurrent SCLC: CheckMate 032. *J. Thorac. Oncol.* 14, 237–244.
 29. Iams, W.T., Porter, J., and Horn, L. (2020). Immunotherapeutic approaches for small-cell lung cancer. *Nat. Rev. Clin. Oncol.* 17, 300–312.
 30. Inoue, M., Sawabata, N., and Okumura, M. (2012). Surgical intervention for small-cell lung cancer: what is the surgical role? *Gen. Thorac. Cardiovasc. Surg.* 60, 401–405.
 31. Zhao, G., Zhao, Z., and Jia, Y. (2019). Advances in preclinical models of small cell lung cancer. *Med One* 4, e190019.
 32. Travis, W.D., Brambilla, E., Nicholson, A.G., Yatabe, Y., Austin, J.H.M., Beasley, M.B., Chirieac, L.R., Dacic, S., Duhig, E., Flieder, D.B., et al. (2015). The 2015 World Health Organization classification of lung tumors: impact of genetic, clinical and radiologic advances since the 2004 classification. *J. Thorac. Oncol.* 10, 1243–1260.
 33. Rudin, C.M., Poirier, J.T., Byers, L.A., Dive, C., Dowlati, A., George, J., Heymach, J.V., Johnson, J.E., Lehman, J.M., MacPherson, D., et al. (2019). Molecular subtypes of small cell lung cancer: a synthesis of human and mouse model data. *Nat. Rev. Cancer* 19, 289–297.
 34. Drapkin, B.J., and Rudin, C.M. (2020). Advances in small-cell lung cancer (SCLC) translational research. *Cold Spring Harb. Perspect. Med.* a038240.
 35. Zhang, W., Girard, L., Zhang, Y.A., Haruki, T., Papari-Zareei, M., Stastny, V., Ghayee, H.K., Pacak, K., Oliver, T.G., Minna, J.D., and Gazdar, A.F. (2018). Small cell lung cancer tumors and preclinical models display heterogeneity of neuroendocrine phenotypes. *Transl. Lung Cancer Res.* 7, 32–49.
 36. Gazdar, A.F. (2018). Molecular phenotypes of SCLC. *J. Thorac. Oncol.* 13 (10, Suppl). S309. <https://doi.org/10.1016/j.jtho.2018.08.211>.
 37. Saito, M., Shiraiishi, K., Goto, A., Suzuki, H., Kohno, T., and Kono, K. (2018). Development of targeted therapy and immunotherapy for treatment of small cell lung cancer. *Jpn. J. Clin. Oncol.* 48, 603–608.
 38. Dora, D., Rivard, C., Yu, H., Bunn, P., Suda, K., Ren, S., Lueke Pickard, S., Laszlo, V., Harko, T., Megyesfalvi, Z., et al. (2020). Neuroendocrine subtypes of small cell lung cancer differ in terms of immune microenvironment and checkpoint molecule distribution. *Mol. Oncol.* 14, 1947–1965.
 39. Carney, D.N., Gazdar, A.F., Bepler, G., Guccion, J.G., Marangos, P.J., Moody, T.W., Zweig, M.H., and Minna, J.D. (1985). Establishment and identification of small cell lung cancer cell lines having classic and variant features. *Cancer Res.* 45, 2913–2923.
 40. Gazdar, A.F., Carney, D.N., Nau, M.M., and Minna, J.D. (1985). Characterization of variant subclasses of cell lines derived from small cell lung cancer having distinctive biochemical, morphological, and growth properties. *Cancer Res.* 45, 2924–2930.
 41. Baine, M.K., Hsieh, M.S., Lai, W.V., Egger, J.V., Jungbluth, A.A., Daneshbod, Y., Beras, A., Spencer, R., Lopardo, J., Bodd, F., et al. (2020). SCLC subtypes defined by ASCL1, NEUROD1, POU2F3, and YAP1: a comprehensive immunohistochemical and histopathologic characterization. *J. Thorac. Oncol.* 15, 1823–1835.
 42. Borromeo, M.D., Savage, T.K., Kollipara, R.K., He, M., Augustyn, A., Osborne, J.K., Girard, L., Minna, J.D., Gazdar, A.F., Cobb, M.H., and Johnson, J.E. (2016). ASCL1 and NEUROD1 reveal heterogeneity in pulmonary neuroendocrine tumors and regulate distinct genetic programs. *Cell Rep.* 16, 1259–1272.
 43. Neptune, E.R., Podowski, M., Calvi, C., Cho, J.H., Garcia, J.G., Tuder, R., Linnoila, R.I., Tsai, M.J., and Dietz, H.C. (2008). Targeted disruption of NeuroD, a proneural basic helix-loop-helix factor, impairs distal lung formation and neuroendocrine morphology in the neonatal lung. *J. Biol. Chem.* 283, 21160–21169.
 44. Borges, M., Linnoila, R.I., van de Velde, H.J., Chen, H., Nelkin, B.D., Mabry, M., Baylin, S.B., and Ball, D.W. (1997). An achaete-scute homologue essential for neuroendocrine differentiation in the lung. *Nature* 386, 852–855.
 45. McColl, K., Wildey, G., Sakre, N., Lipka, M.B., Behtaj, M., Kresak, A., Chen, Y., Yang, M., Velcheti, V., Fu, P., and Dowlati, A. (2017). Reciprocal expression of INSM1 and YAP1 defines subgroups in small cell lung cancer. *Oncotarget* 8, 73745–73756.
 46. Huang, Y.-H., Klingbeil, O., He, X.-Y., Wu, X.S., Arun, G., Lu, B., Somerville, T.D.D., Milazzo, J.P., Wilkinson, J.E., Demerdash, O.E., et al. (2018). POU2F3 is a master regulator of a tuft cell-like variant of small cell lung cancer. *Genes Dev.* 32, 915–928.
 47. Shue, Y.T., Lim, J.S., and Sage, J. (2018). Tumor heterogeneity in small cell lung cancer defined and investigated in pre-clinical mouse models. *Transl. Lung Cancer Res.* 7, 21–31.
 48. Denny, S.K., Yang, D., Chuang, C.H., Brady, J.J., Lim, J.S., Grüner, B.M., Chiou, S.H., Schep, A.N., Baral, J., Hamard, C., et al. (2016). Nf1b promotes metastasis through a widespread increase in chromatin accessibility. *Cell* 166, 328–342.
 49. Lohinai, Z., Megyesfalvi, Z., Suda, K., Harko, T., Ren, S., Moldvay, J., Laszlo, V., Rivard, C., Dome, B., and Hirsch, F.R. (2019). Comparative expression analysis in small cell lung carcinoma reveals neuroendocrine pattern change in primary tumor versus lymph node metastases. *Transl. Lung Cancer Res.* 8, 938–950.
 50. Rudin, C.M., Durinck, S., Stawiski, E.W., Poirier, J.T., Modrusan, Z., Shames, D.S., Bergbower, E.A., Guan, Y., Shin, J., Guillory, J., et al. (2012). Comprehensive genomic analysis identifies SOX2 as a frequently amplified gene in small-cell lung cancer. *Nat. Genet.* 44, 1111–1116.
 51. Lim, J.S., Ibaseta, A., Fischer, M.M., Cancilla, B., O'Young, G., Cristea, S., Luca, V.C., Yang, D., Jahchan, N.S., Hamard, C., et al. (2017). Intratumoural heterogeneity generated by Notch signalling promotes small-cell lung cancer. *Nature* 545, 360–364.
 52. Ntziachristos, P., Lim, J.S., Sage, J., and Aifantis, I. (2014). From fly wings to targeted cancer therapies: a centennial for notch signaling. *Cancer Cell* 25, 318–334.
 53. Thiel, G., Ekici, M., and Rössler, O.G. (2015). RE-1 silencing transcription factor (REST): a regulator of neuronal development and neuronal/endocrine function. *Cell Tissue Res.* 359, 99–109.
 54. Chalisehar, M.D., Wait, S.J., Huang, F., Ireland, A.S., Mukhopadhyay, A., Lee, Y., Schuman, S.S., Guthrie, M.R., Berrett, K.C., Vahrenkamp, J.M., et al. (2019). MYC-driven small-cell lung cancer is metabolically distinct and vulnerable to arginine depletion. *Clin. Cancer Res.* 25, 5107–5121.
 55. Mollaoglu, G., Guthrie, M.R., Böhm, S., Brägelmann, J., Can, I., Ballieu, P.M., Marx, A., George, J., Heinen, C., Chalisehar, M.D., et al. (2017). MYC drives progression of small cell lung cancer to a variant neuroendocrine subtype with vulnerability to aurora kinase inhibition. *Cancer Cell* 31, 270–285.

56. Massó-Vallés, D., Beaulieu, M.E., and Soucek, L. (2020). MYC, MYCL, and MYCN as therapeutic targets in lung cancer. *Expert Opin. Ther. Targets* 24, 101–114.
57. Lum, C., and Alamgeer, M. (2019). Technological and therapeutic advances in advanced small cell lung cancer. *Cancers (Basel)* 11, 1570.
58. Yan, M., Wang, C., He, B., Yang, M., Tong, M., Long, Z., Liu, B., Peng, F., Xu, L., Zhang, Y., et al. (2016). Aurora-A kinase: a potent oncogene and target for cancer therapy. *Med. Res. Rev.* 36, 1036–1079.
59. Sos, M.L., Dietlein, F., Peifer, M., Schöttle, J., Balke-Want, H., Müller, C., Koker, M., Richters, A., Heynck, S., Malchers, F., et al. (2012). A framework for identification of actionable cancer genome dependencies in small cell lung cancer. *Proc. Natl. Acad. Sci. USA* 109, 17034–17039.
60. Sabari, J.K., Lok, B.H., Laird, J.H., Poirier, J.T., and Rudin, C.M. (2017). Unravelling the biology of SCLC: implications for therapy. *Nat. Rev. Clin. Oncol.* 14, 549–561.
61. Schultheis, A.M., Scheel, A.H., Ozretić, L., George, J., Thomas, R.K., Hagemann, T., Zander, T., Wolf, J., and Buettner, R. (2015). PD-L1 expression in small cell neuroendocrine carcinomas. *Eur. J. Cancer* 51, 421–426.
62. Schultheis, A.M., Bos, M., Schmitz, K., Wilsberg, L., Binot, E., Wolf, J., Büttner, R., and Schildhaus, H.U. (2014). Fibroblast growth factor receptor 1 (FGFR1) amplification is a potential therapeutic target in small-cell lung cancer. *Mod. Pathol.* 27, 214–221.
63. Karim, S.M., and Zekri, J. (2012). Chemotherapy for small cell lung cancer: a comprehensive review. *Oncol. Rev.* 6, e4.
64. Bumber, Y. (2018). Tumor mutational burden (TMB) as a biomarker of response to immunotherapy in small cell lung cancer. *J. Thorac. Dis.* 10, 4689–4693.
65. Stewart, C.A., Gay, C.M., Xi, Y., Sivaji, S., Sivakamasundari, V., Fujimoto, J., Bolisetty, M., Hartsfield, P.M., Balasubramanian, V., Chalisahar, M.D., et al. (2020). Single-cell analyses reveal increased intratumoral heterogeneity after the onset of therapy resistance in small-cell lung cancer. *Nat. Can.* 1, 423–436.
66. Meder, L., Büttner, R., and Odenthal, M. (2017). Notch signaling triggers the tumor heterogeneity of small cell lung cancer. *J. Thorac. Dis.* 9, 4884–4888.
67. Ng, J., and Sutherland, K.D. (2020). NOTCH your usual suspect: MYC charged with controlling neuroendocrine cell-fate in small cell lung cancer. *Cancer Cell* 38, 17–20.
68. Zoppi, G., Regairaz, M., Leo, E., Reinhold, W.C., Varma, S., Ballestrero, A., Doroshov, J.H., and Pommier, Y. (2012). Putative DNA/RNA helicase Schlafen-11 (SLFN11) sensitizes cancer cells to DNA-damaging agents. *Proc. Natl. Acad. Sci. USA* 109, 15030–15035.
69. Allison Stewart, C., Tong, P., Cardnell, R.J., Sen, T., Li, L., Gay, C.M., Masrourpour, F., Fan, Y., Bara, R.O., Feng, Y., et al. (2017). Dynamic variations in epithelial-to-mesenchymal transition (EMT), ATM, and SLFN11 govern response to PARP inhibitors and cisplatin in small cell lung cancer. *Oncotarget* 8, 28575–28587.
70. Nogales, V., Reinhold, W.C., Varma, S., Martinez-Cardus, A., Moutinho, C., Moran, S., Heyn, H., Sebio, A., Barnadas, A., Pommier, Y., and Esteller, M. (2016). Epigenetic inactivation of the putative DNA/RNA helicase SLFN11 in human cancer confers resistance to platinum drugs. *Oncotarget* 7, 3084–3097.
71. Gardner, E.E., Lok, B.H., Schneeberger, V.E., Desmeules, P., Miles, L.A., Arnold, P.K., Ni, A., Khodos, I., de Stanchina, E., Nguyen, T., et al. (2017). Chemosensitive relapse in small cell lung cancer proceeds through an EZH2-SLFN11 axis. *Cancer Cell* 31, 286–299.
72. Gopalakrishnan, D., Koshkin, V.S., Ornstein, M.C., Papatsoris, A., and Grivas, P. (2018). Immune checkpoint inhibitors in urothelial cancer: recent updates and future outlook. *Ther. Clin. Risk Manag.* 14, 1019–1040.
73. Herrscher, H., and Robert, C. (2020). Immune checkpoint inhibitors in melanoma in the metastatic, neoadjuvant, and adjuvant setting. *Curr. Opin. Oncol.* 32, 106–113.
74. Pavan, A., Attili, I., Pasello, G., Guarneri, V., Conte, P.F., and Bonanno, L. (2019). Immunotherapy in small-cell lung cancer: from molecular promises to clinical challenges. *J. Immunother. Cancer* 7, 205.
75. Kim, J.H., Kim, H.S., and Kim, B.J. (2017). Prognostic value of smoking status in non-small-cell lung cancer patients treated with immune checkpoint inhibitors: a meta-analysis. *Oncotarget* 8, 93149–93155.
76. Reck, M., Rodríguez-Abreu, D., Robinson, A.G., Hui, R., Csőszi, T., Fülöp, A., Gottfried, M., Peled, N., Tafreshi, A., Cuffe, S., et al.; KEYNOTE-024 Investigators (2016). Pembrolizumab versus chemotherapy for PD-L1-positive non-small-cell lung cancer. *N. Engl. J. Med.* 375, 1823–1833.
77. Chalmers, Z.R., Connelly, C.F., Fabrizio, D., Gay, L., Ali, S.M., Ennis, R., Schrock, A., Campbell, B., Shlien, A., Chmielecki, J., et al. (2017). Analysis of 100,000 human cancer genomes reveals the landscape of tumor mutational burden. *Genome Med.* 9, 34.
78. Sebastian, M., Koschade, S., and Stratmann, J.A. (2019). SCLC, paraneoplastic syndromes, and the immune system. *J. Thorac. Oncol.* 14, 1878–1880.
79. Mansfield, A.S., Kaźarnowicz, A., Karaseva, N., Sánchez, A., De Boer, R., Andric, Z., Reck, M., Atagi, S., Lee, J.S., Garassino, M., et al. (2020). Safety and patient-reported outcomes of atezolizumab, carboplatin, and etoposide in extensive-stage small-cell lung cancer (IMpower133): a randomized phase I/III trial. *Ann. Oncol.* 31, 310–317.
80. Paz-Ares, L., Dvorkin, M., Chen, Y., Reinmuth, N., Hotta, K., Trukhin, D., Statsenko, G., Hochmair, M.J., Özgüroğlu, M., Ji, J.H., et al.; CASPIAN investigators (2019). Durvalumab plus platinum-etoposide versus platinum-etoposide in first-line treatment of extensive-stage small-cell lung cancer (CASPIAN): a randomised, controlled, open-label, phase 3 trial. *Lancet* 394, 1929–1939.
81. Chung, H.C., Piha-Paul, S.A., Lopez-Martin, J., Schellens, J.H.M., Kao, S., Miller, W.H., Jr., Delord, J.P., Gao, B., Planchard, D., Gottfried, M., et al. (2020). Pembrolizumab after two or more lines of previous therapy in patients with recurrent or metastatic SCLC: results from the KEYNOTE-028 and KEYNOTE-158 studies. *J. Thorac. Oncol.* 15, 618–627.
82. Ready, N.E., Ott, P.A., Hellmann, M.D., Zugazagoitia, J., Hann, C.L., de Braud, F., Antonia, S.J., Ascierto, P.A., Moreno, V., Atmaca, A., et al. (2020). Nivolumab monotherapy and nivolumab plus ipilimumab in recurrent small cell lung cancer: results from the CheckMate 032 randomized cohort. *J. Thorac. Oncol.* 15, 426–435.
83. Hornyák, L., Dobos, N., Koncz, G., Karányi, Z., Páll, D., Szabó, Z., Halmos, G., and Székely, L. (2018). The role of indoleamine-2,3-dioxygenase in cancer development, diagnostics, and therapy. *Front. Immunol.* 9, 151.
84. Katz, J.B., Muller, A.J., and Prendergast, G.C. (2008). Indoleamine 2,3-dioxygenase in T-cell tolerance and tumoral immune escape. *Immunol. Rev.* 222, 206–221.
85. Sloan, K.E., Eustace, B.K., Stewart, J.K., Zehetmeier, C., Torella, C., Simeone, M., Roy, J.E., Unger, C., Louis, D.N., Ilag, L.L., and Jay, D.G. (2004). CD155/PVR plays a key role in cell motility during tumor cell invasion and migration. *BMC Cancer* 4, 73.
86. Friedlaender, A., Addeo, A., and Banna, G. (2019). New emerging targets in cancer immunotherapy: the role of TIM3. *ESMO Open* 4 (Suppl 3), e000497.
87. Leclerc, M., Mezquita, L., Guillebot De Nerville, G., Tihy, I., Malenica, I., Chouaib, S., and Mami-Chouaib, F. (2019). Recent advances in lung cancer immunotherapy: input of T-cell epitopes associated with impaired peptide processing. *Front. Immunol.* 10, 1505.
88. Doyle, A., Martin, W.J., Funa, K., Gazdar, A., Carney, D., Martin, S.E., Linnoila, I., Cuttitta, F., Mulshine, J., Bunn, P., et al. (1985). Markedly decreased expression of class I histocompatibility antigens, protein, and mRNA in human small-cell lung cancer. *J. Exp. Med.* 161, 1135–1151.
89. Doyle, L.A., Cuttitta, F., Mulshine, J.L., Bunn, P.A., and Minna, J.D. (1987). Markedly different antibody responses to immunized small cell and non-small cell lung cancer cells. *Cancer Res.* 47, 5009–5013.
90. Yazawa, T., Kamma, H., Fujiwara, M., Matsui, M., Horiguchi, H., Satoh, H., Fujimoto, M., Yokoyama, K., and Ogata, T. (1999). Lack of class II transactivator causes severe deficiency of HLA-DR expression in small cell lung cancer. *J. Pathol.* 187, 191–199.
91. Arbour, K.C., Mezquita, L., Long, N., Rizvi, H., Auclin, E., Ni, A., Martínez-Bernal, G., Ferrara, R., Lai, W.V., Hendriks, L.E.L., et al. (2018). Impact of baseline steroids on efficacy of programmed cell death-1 and programmed death-ligand 1 blockade in patients with non-small-cell lung cancer. *J. Clin. Oncol.* 36, 2872–2878.
92. König, D., and Läubli, H. (2020). Mechanisms of immune-related complications in cancer patients treated with immune checkpoint inhibitors. *Pharmacology*. Published online July 28, 2020. <https://doi.org/10.1159/000509081>.
93. Mizrahi, M., and Ilan, Y. (2009). The gut mucosa as a site for induction of regulatory T-cells. *Curr. Pharm. Des.* 15, 1191–1202.

94. Anderson, R., Theron, A.J., and Rapoport, B.L. (2019). Immunopathogenesis of immune checkpoint inhibitor-related adverse events: roles of the intestinal microbiome and Th17 cells. *Front. Immunol.* **10**, 2254.
95. Atrati, F., Groen, H.J.M., Byers, L.A., Garralda, E., Lolkema, M.P., Sangha, R.S., Viteri Ramirez, S., Chae, Y.K., Camidge, D.R., Gabrail, N.Y., et al. (2017). Phase 1/2 study of veliparib (V) combined with carboplatin (Cb) and etoposide (E) in patients (pts) with extensive-stage disease (ED) small cell lung cancer (SCLC) and other solid tumors: phase 1 results. *J. Clin. Oncol.* **35** (15, Suppl), 8530.
96. Wainberg, Z.A., de Bono, J.S., Mina, L., Sachdev, J., Byers, L.A., Chugh, R., Zhang, C., Henshaw, J.W., Dorr, A., Glaspy, J., et al. (2013). Update on first-in-man trial of novel oral PARP inhibitor BMN 673 in patients with solid tumors. *Mol. Cancer Ther.* **12** (11, Suppl), C295.
97. Pietanza, M.C., Waqar, S.N., Krug, L.M., Dowlati, A., Hann, C.L., Chiappori, A., Owonikoko, T.K., Woo, K.M., Cardnell, R.J., Fujimoto, J., et al. (2018). Randomized, double-blind, phase II study of temozolomide in combination with either veliparib or placebo in patients with relapsed-sensitive or refractory small-cell lung cancer. *J. Clin. Oncol.* **36**, 2386–2394.
98. Owonikoko, T.K., Dahlberg, S.E., Sica, G.L., Wagner, L.I., Wade, J.L., 3rd, Srkalovic, G., Lash, B.W., Leach, J.W., Leal, T.B., Aggarwal, C., and Ramalingam, S.S. (2019). Randomized phase II trial of cisplatin and etoposide in combination with veliparib or placebo for extensive-stage small-cell lung cancer: ECOG-ACRIN 2511 study. *J. Clin. Oncol.* **37**, 222–229.
99. AbbVie (2018). Phase 3 trial of Rova-T as second-line therapy for advanced small-cell lung cancer (TAHOE study), <https://news.abbvie.com/news/phase-3-trial-rova-t-as-second-line-therapy-for-advanced-small-cell-lung-cancer-tahoe-study-halted.htm>.
100. Morgensztern, D., Besse, B., Greillier, L., Santana-Davila, R., Ready, N., Hann, C.L., Glisson, B.S., Farago, A.F., Dowlati, A., Rudin, C.M., et al. (2019). Efficacy and safety of rovalpituzumab tesirine in third-line and beyond patients with DLL3-expressing, relapsed/refractory small-cell lung cancer: results from the phase II TRINITY study. *Clin. Cancer Res.* **25**, 6958–6966.
101. Cheng, Y., Wang, Q., Li, K., Shi, J., Wu, L., Han, B., Chen, G., He, J., Wang, J., Qin, H., et al. (2018). Anlotinib as third-line or further-line treatment in relapsed SCLC: a multicentre, randomized, double-blind phase 2 trial. *J. Thorac. Oncol.* **13** (10, Suppl), S351.
102. Soria, J.C., Johnson, B.E., and Chevalier, T.L. (2003). Imatinib in small cell lung cancer. *Lung Cancer* **41** (Suppl 1), S49–S53.
103. Chiappori, A.A., Otterson, G.A., Dowlati, A., Traynor, A.M., Horn, L., Owonikoko, T.K., Ross, H.J., Hann, C.L., Abu Hejleh, T., Nieva, J., et al. (2016). A randomized phase II study of linsitinib (OSI-906) versus topotecan in patients with relapsed small-cell lung cancer. *Oncologist* **21**, 1163–1164.
104. Ellis, P.M., Shepherd, F.A., Laurie, S.A., Goss, G.D., Olivo, M., Powers, J., Seymour, L., and Bradbury, P.A. (2014). NCIC CTG IND.190 phase I trial of dalotuzumab (MK-0646) in combination with cisplatin and etoposide in extensive-stage small-cell lung cancer. *J. Thorac. Oncol.* **9**, 410–413.
105. Owonikoko, T.K., Niu, H., Nackaerts, K., Csozsi, T., Ostoros, G., Mark, Z., Baik, C., Joy, A.A., Chouaid, C., Jaime, J.C., et al.; C14018 study investigators (2020). Randomized phase II study of paclitaxel plus alisertib versus paclitaxel plus placebo as second-line therapy for SCLC: primary and correlative biomarker analyses. *J. Thorac. Oncol.* **15**, 274–287.
106. Spigel, D.R., Townley, P.M., Waterhouse, D.M., Fang, L., Adiguzel, I., Huang, J.E., Karlin, D.A., Faoro, L., Scappaticci, F.A., and Socinski, M.A. (2011). Randomized phase II study of bevacizumab in combination with chemotherapy in previously untreated extensive-stage small-cell lung cancer: results from the SALUTE trial. *J. Clin. Oncol.* **29**, 2215–2222.
107. Trigo, J., Subbiah, V., Besse, B., Moreno, V., López, R., Sala, M.A., Peters, S., Ponce, S., Fernández, C., Alfaro, V., et al. (2020). Lurbinectedin as second-line treatment for patients with small-cell lung cancer: a single-arm, open-label, phase 2 basket trial. *Lancet Oncol.* **21**, 645–654.
108. Schulze, A.B., Evers, G., Kerkhoff, A., Mohr, M., Schliemann, C., Berdel, W.E., and Schmidt, L.H. (2019). Future options of molecular-targeted therapy in small cell lung cancer. *Cancers (Basel)* **11**, 690.
109. Byers, L.A., Wang, J., Nilsson, M.B., Fujimoto, J., Saintigny, P., Yordy, J., Giri, U., Peyton, M., Fan, Y.H., Diao, L., et al. (2012). Proteomic profiling identifies dysregulated pathways in small cell lung cancer and novel therapeutic targets including PARP1. *Cancer Discov.* **2**, 798–811.
110. Calabrese, C.R., Almassy, R., Barton, S., Batey, M.A., Calvert, A.H., Canan-Koch, S., Durkacz, B.W., Hostomsky, Z., Kumpf, R.A., Kyle, S., et al. (2004). Anticancer chemosensitization and radiosensitization by the novel poly(ADP-ribose) polymerase-1 inhibitor AG14361. *J. Natl. Cancer Inst.* **96**, 56–67.
111. Tata, P.R., and Rajagopal, J. (2017). Plasticity in the lung: making and breaking cell identity. *Development* **144**, 755–766.
112. Melichar, B., Adenis, A., Lockhart, A.C., Bennouna, J., Dees, E.C., Kayaleh, O., Obermannova, R., DeMichele, A., Zatloukal, P., Zhang, B., et al. (2015). Safety and activity of alisertib, an investigational aurora kinase A inhibitor, in patients with breast cancer, small-cell lung cancer, non-small-cell lung cancer, head and neck squamous-cell carcinoma, and gastro-oesophageal adenocarcinoma: a five-arm phase 2 study. *Lancet Oncol.* **16**, 395–405.
113. Vallières, E., Shepherd, F.A., Crowley, J., Van Houtte, P., Postmus, P.E., Carney, D., Chansky, K., Shaikh, Z., and Goldstraw, P.; International Association for the Study of Lung Cancer International Staging Committee and Participating Institutions (2009). The IASLC Lung Cancer Staging Project: proposals regarding the relevance of TNM in the pathologic staging of small cell lung cancer in the forthcoming (seventh) edition of the TNM classification for lung cancer. *J. Thorac. Oncol.* **4**, 1049–1059.
114. Schreiber, D., Rineer, J., Weedon, J., Vongtama, D., Wortham, A., Kim, A., Han, P., Choi, K., and Rotman, M. (2010). Survival outcomes with the use of surgery in limited-stage small cell lung cancer: should its role be re-evaluated? *Cancer* **116**, 1350–1357.
115. Varlotto, J.M., Recht, A., Flickinger, J.C., Medford-Davis, L.N., Dyer, A.-M., and DeCamp, M.M. (2011). Lobectomy leads to optimal survival in early-stage small cell lung cancer: a retrospective analysis. *J. Thorac. Cardiovasc. Surg.* **142**, 538–546.
116. Takei, H., Kondo, H., Miyaoka, E., Asamura, H., Yoshino, I., Date, H., Okumura, M., Tada, H., Fujii, Y., Nakanishi, Y., et al.; Japanese Joint Committee of Lung Cancer Registry (2014). Surgery for small cell lung cancer: a retrospective analysis of 243 patients from Japanese Lung Cancer Registry in 2004. *J. Thorac. Oncol.* **9**, 1140–1145.
117. Combs, S.E., Hancock, J.G., Boffa, D.J., Decker, R.H., Dettterbeck, F.C., and Kim, A.W. (2015). Bolstering the case for lobectomy in stages I, II, and IIIA small-cell lung cancer using the National Cancer Data Base. *J. Thorac. Oncol.* **10**, 316–323.
118. Hoda, M.A., Klikovits, T., and Klepetko, W. (2018). Controversies in oncology: surgery for small cell lung cancer? It's time to rethink the case. *ESMO Open* **3**, e000366.
119. Dowell, J.E. (2010). Small cell lung cancer: are we making progress? *Am. J. Med. Sci.* **339**, 68–76.
120. Taniguchi, H., Sen, T., and Rudin, C.M. (2020). Targeted therapies and biomarkers in small cell lung cancer. *Front. Oncol.* **10**, 741.
121. Kim, D.-W., Kim, K.-C., Kim, K.-B., Dunn, C.T., and Park, K.-S. (2018). Transcriptional deregulation underlying the pathogenesis of small cell lung cancer. *Transl. Lung Cancer Res.* **7**, 4–20.
122. Furuta, M., Sakakibara-Konishi, J., Kikuchi, H., Yokouchi, H., Nishihara, H., Minemura, H., Harada, M., Yamazaki, S., Akie, K., Fujita, Y., et al.; Hokkaido Lung Cancer Clinical Study Group (2019). Analysis of DLL3 and ASCL1 in surgically resected small cell lung cancer (HOT1702). *Oncologist* **24**, e1172–e1179.
123. Takagi, S., Ishikawa, Y., Mizutani, A., Iwasaki, S., Matsumoto, S., Kamada, Y., Nomura, T., and Nakamura, K. (2017). LSD1 inhibitor T-3775440 inhibits SCLC cell proliferation by disrupting LSD1 interactions with SNAG domain proteins INSM1 and GF11B. *Cancer Res.* **77**, 4652–4662.
124. Augert, A., Eastwood, E., Ibrahim, A.H., Wu, N., Grunblatt, E., Basom, R., Liggitt, D., Eaton, K.D., Martins, R., Poirier, J.T., et al. (2019). Targeting NOTCH activation in small cell lung cancer through LSD1 inhibition. *Sci. Signal.* **12**, eaau2922.
125. Jia, D., Augert, A., Kim, D.W., Eastwood, E., Wu, N., Ibrahim, A.H., Kim, K.B., Dunn, C.T., Pillai, S.P.S., Gazdar, A.F., et al. (2018). *Crebbp* loss drives small cell lung cancer and increases sensitivity to HDAC inhibition. *Cancer Discov.* **8**, 1422–1437.

126. Tsoukalas, N., Aravantinou-Fatorou, E., Baxevanos, P., Tolia, M., Tsapakidis, K., Galanopoulos, M., Liontos, M., and Kyrgias, G. (2018). Advanced small cell lung cancer (SCLC): new challenges and new expectations. *Ann. Transl. Med.* 6, 145.
127. Poirier, J.T., Dobromilskaya, I., Moriarty, W.F., Peacock, C.D., Hann, C.L., and Rudin, C.M. (2013). Selective tropism of Seneca Valley virus for variant subtype small cell lung cancer. *J. Natl. Cancer Inst.* 105, 1059–1065.
128. Cao, L., Zhang, R., Liu, T., Sun, Z., Hu, M., Sun, Y., Cheng, L., Guo, Y., Fu, S., Hu, J., et al. (2018). Seneca Valley virus attachment and uncoating mediated by its receptor anthrax toxin receptor 1. *Proc. Natl. Acad. Sci. USA* 115, 13087–13092.
129. Inno, A., Stagno, A., and Gori, S. (2018). Schlafen-11 (SLFN11): a step forward towards personalized medicine in small-cell lung cancer? *Transl. Lung Cancer Res.* 7 (Suppl 4), S341–S345.
130. Gay, C.M., Diao, L., Stewart, C.A., Xi, Y., Cardnell, R.J., Swisher, S.G., Roth, J.A., Glisson, B.S., Wang, J., Heymach, J.V., et al. (2019). Inter- and intra-tumoral variations in ASCL1, NEUROD1, and POU2F3 transcriptional programs underlie three distinct molecular subtypes of small cell lung cancers. *Cancer Res.* 79 (13, Suppl), 3772. <https://doi.org/10.1158/1538-7445.AM2019-3772>.
131. Shibata, M., Ham, K., and Hoque, M.O. (2018). A time for YAP1: tumorigenesis, immunosuppression and targeted therapy. *Int. J. Cancer* 143, 2133–2144.
132. Lee, B.S., Park, D.I., Lee, D.H., Lee, J.E., Yeo, M.K., Park, Y.H., Lim, D.S., Choi, W., Lee, D.H., Yoo, G., et al. (2017). Hippo effector YAP directly regulates the expression of PD-L1 transcripts in EGFR-TKI-resistant lung adenocarcinoma. *Biochem. Biophys. Res. Commun.* 491, 493–499.
133. Horie, M., Saito, A., Ohshima, M., Suzuki, H.I., and Nagase, T. (2016). YAP and TAZ modulate cell phenotype in a subset of small cell lung cancer. *Cancer Sci.* 107, 1755–1766.
134. Bebb, C.M., Thomas, E.S., Chen, Z., Stroh, J., Androulidaki, A., Schmitt, A., Höhne, M.N., Stüker, L., de Pádua Alves, C., Khonsari, A., et al. (2020). Ferroptosis response segregates small cell lung cancer (SCLC) neuroendocrine subtypes. *bioRxiv*. <https://doi.org/10.1101/2020.07.11.198408>.
135. Yang, D., Denny, S.K., Greenside, P.G., Chaikovskiy, A.C., Brady, J.J., Ouadah, Y., Granja, J.M., Jahchan, N.S., Lim, J.S., Kwok, S., et al. (2018). Intertumoral heterogeneity in SCLC is influenced by the cell type of origin. *cancer Discov.* 8, 1316–1331.
136. Nakazawa, K., Kurishima, K., Tamura, T., Kagohashi, K., Ishikawa, H., Satoh, H., and Hizawa, N. (2012). Specific organ metastases and survival in small cell lung cancer. *Oncol. Lett.* 4, 617–620.
137. Park, K.S., Liang, M.C., Raiser, D.M., Zamponi, R., Roach, R.R., Curtis, S.J., Walton, Z., Schaffer, B.E., Roake, C.M., Zmoos, A.F., et al. (2011). Characterization of the cell of origin for small cell lung cancer. *Cell Cycle* 10, 2806–2815.
138. Rudin, C.M., Awad, M.M., Navarro, A., Gottfried, M., Peters, S., Csösz, T., Cheema, P.K., Rodriguez-Abreu, D., Wollner, M., Yang, J.C., et al.; KEYNOTE-604 Investigators (2020). Pembrolizumab or placebo plus etoposide and platinum as first-line therapy for extensive-stage small-cell lung cancer: randomized, double-blind, phase III KEYNOTE-604 study. *J. Clin. Oncol.* 38, 2369–2379.
139. Besse, B., Menis, J., Bironzo, P., Gervais, R., Greillier, L., Monnet, I., Livi, L., Young, R., Decroisette, C., Cloarec, N., et al. (2020). LBA85—REACTION: A phase II study of etoposide and cis/carboplatin with or without pembrolizumab in untreated extensive small cell lung cancer. *Ann. Oncol.* 31 (Suppl 4), S1211–S1212.
140. Reck, M., Luft, A., Szczesna, A., Havel, L., Kim, S.-W., Akerley, W., Pietanza, M.C., Wu, Y.L., Zielinski, C., Thomas, M., et al. (2016). Phase III randomized trial of ipilimumab plus etoposide and platinum versus placebo plus etoposide and platinum in extensive-stage small-cell lung cancer. *J. Clin. Oncol.* 34, 3740–3748.
141. Owonikoko, T.K., Kim, H.R., Govindan, R., Ready, N., Reck, M., Peters, S., Dakhil, S.R., Navarro, A., Rodriguez-Cid, J., Schenker, M., et al. (2019). Nivolumab (nivo) plus ipilimumab (ipi), nivo, or placebo (pbo) as maintenance therapy in patients (pts) with extensive disease small cell lung cancer (ED-SCLC) after first-line (1L) platinum-based chemotherapy (chemo): results from the double-blind, randomized phase III CheckMate 451 study. *Ann. Oncol.* 30 (Suppl 2), ii77.
142. Gadgeel, S.M., Pennell, N.A., Fidler, M.J., Halmos, B., Bonomi, P., Stevenson, J., Schneider, B., Sukari, A., Ventimiglia, J., Chen, W., et al. (2018). Phase II study of maintenance pembrolizumab in patients with extensive-stage small cell lung cancer (SCLC). *J. Thorac. Oncol.* 13, 1393–1399.
143. Antonia, S.J., López-Martin, J.A., Bendell, J., Ott, P.A., Taylor, M., Eder, J.P., Jäger, D., Pietanza, M.C., Le, D.T., de Braud, F., et al. (2016). Nivolumab alone and nivolumab plus ipilimumab in recurrent small-cell lung cancer (CheckMate 032): a multicentre, open-label, phase 1/2 trial. *Lancet Oncol.* 17, 883–895.
144. Fischer, R.N., George, J., Scheel, A.H., Schloesser, H.A., Vehreschild, M., Brossart, P., Engel-Riedel, W., Griesinger, F., Grohé, C., Kambartel, K.-O., et al. (2019). BIOLUMA: a phase II trial of nivolumab in combination with ipilimumab to evaluate efficacy and safety in lung cancer and to evaluate biomarkers predictive for response—preliminary results from the SCLC cohort. *J. Clin. Oncol.* 37 (15, Suppl), 8563.
145. Reck, M., Vicente, D., Ciuleanu, T., Gettinger, S., Peters, S., Horn, L., Audigier-Valette, C., Pardo, N., Juan-Vidal, O., Cheng, Y., et al. (2018). Efficacy and safety of nivolumab (nivo) monotherapy versus chemotherapy (chemo) in recurrent small cell lung cancer (SCLC): results from CheckMate 331. *Ann. Oncol.* 29 (Suppl 10), x43.
146. Ott, P.A., Elez, E., Hirt, S., Kim, D.-W., Morosky, A., Saraf, S., Piperdi, B., and Mehnert, J.M. (2017). Pembrolizumab in patients with extensive-stage small-cell lung cancer: results from the phase Ib KEYNOTE-028 study. *J. Clin. Oncol.* 35, 3823–3829.
147. Chung, H.C., Lopez-Martin, J.A., Kao, S.C.-H., Miller, W.H., Ros, W., Gao, B., Marabelle, A., Gottfried, M., Zer, A., Delord, J.-P., et al. (2018). Phase 2 study of pembrolizumab in advanced small-cell lung cancer (SCLC): KEYNOTE-158. *J. Clin. Oncol.* 36 (15, Suppl), 8506.
148. Cho, D.C., Mahipal, A., Dowlati, A., Chow, W.A., Segal, N.H., Chung, K.Y., Schneider, B.J., Nemunaitis, J.J., Abdul Razak, A.R., Tsai, F.Y.-C., et al. (2018). Safety and clinical activity of durvalumab in combination with tremelimumab in extensive disease small-cell lung cancer (ED-SCLC). *J. Clin. Oncol.* 36 (15, Suppl), 8517.