

A check-list of the family Phoridae (Diptera) of Hungary\*

L. Ádám and L. Papp

**A check-list of the family Phoridae (Diptera) of Hungary** — A total of 207 species of Phoridae recorded hitherto in Hungary is listed with the names of the other species whose known occurrence data make probable their finding also in Hungary. The genera *Aenigmatias*, *Plectanocnema* and *Woodiphora* are new for the Hungarian fauna.

An application for a grant for studies on insect guilds on “very small-sized feeding sources” (more concretely droppings of forest animals, dead snails, dead mice, decaying fungi, oozing sap of deciduous woods, etc.) was accepted by the Hungarian National Scientific Research Fund in 1995. Early in the preparation for these studies it became obvious that phorid species are important members of all these guilds and during the first sampling in 1995 we obtained an abundant and rather various phorid material on these sources.

In order to be prepared for the handling and identification of the Phoridae, last year a key was constructed (based on Schmitz’s keys and on Disney’s (1983) key, supplemented by Dr Disney’s MS key for the *Manual of Palaearctic/European Diptera*) in Hungarian. In 1995–1996 the dry material in the Diptera collection of the Department of Zoology, Hungarian Natural History Museum, Budapest (HNHM) (some 11 000 individuals, double mounted on minute pins) has been assigned to genera by the authors in order to facilitate the identification of the phorids in insect samples collected in our mountain forests. We did not encounter special difficulties in sorting them to genera but — of course — we did when we tried to identify species, particularly so for the species of *Megaselia*. After all, numerous interesting specimens were found: Representatives of 19 genera were found in the “new” materials (collected after 1957); no specimens of *Dohrniphora*, *Menoziola* and *Tubicera* were identified (which were recorded in Schmitz’s works (1922 to 1953)) but specimens of *Aenigmatias*, *Plectanocnema* and *Woodiphora* were found, which genera had not been reported formerly. It seems probable that some species new to science are also included.

Logically the next step in our preparation was a bibliographic search for the faunistic data on the species of the family Phoridae from our country. This work yielded the present paper.

From the summary of the faunistic data, a total of 207 species belonging to the family Phoridae has been reported to occur in Hungary. Some species are mentioned in Schmitz’s (1928, 1940, 1941*a*, 1941*b*) and Disney’s (1991) works merely as from “Hun-

gary", without closer locality data. Of course, this is not a critique on Schmitz's works: he was really great but records like "U" for "Ungarn" (e.g., Schmitz 1928) cannot serve as a solid base for inclusion of a species in the faunistic list for Hungary. As it is well known, the present Hungary covers ca 1/3 of the area of the country before 1920. In addition, a very significant part of the material in the era of K. Kertész's curatorship was collected outside of the present borders but all the names were written on labels in Hungarian orthography, and as Magyarized names of localities which lie now in Slovakia, Croatia, Romania, etc. Scientists other than Hungarians have always had problems with these circumstances and so labels with "Hungaria" or similar notes have caused much confusion. For the present paper we accepted all the records of this kind but since all the old material identified by Schmitz perished in fire in 1956 in the HNHM (so no voucher specimens of the former records exist today), retaining any species on the Hungarian list, which is not represented in the material collected after 1957, must be done with critical care.

In the list below we use square brackets for the specific names, which have not been recorded from Hungary but which are expected to occur here. We based our decisions on their inclusion mainly on Disney's (1991) catalogue. A further 248 species of this kind are listed. We think the phorid fauna of Hungary must be over 500 species but various special collections need to be done in order to add more species to those represented in previous collections from Hungary. We can regard this paper as a very first step in the study of the phorids of the Carpathian Basin.

#### LIST OF SPECIES

##### **Aenigmatias** Meinert, 1890

- [**Aenigmatias brevifrons** (Schmitz, 1955)  
(*Platyphora*)]  
[**Aenigmatias dorni** (Enderlein, 1908)  
(*Oniscomyia*)]  
[**Aenigmatias franzi** Schmitz, 1950]  
[**Aenigmatias lubbocki** (Verrall, 1877)  
(*Platyphora*)]  
[**Aenigmatias picipes** Schmitz, 1927]  
[**Aenigmatias pyrenaicus** (Becker, 1912)  
(*Psalidesma*)]

##### **Anevrina** Lioy, 1864

- Anevrina curvinervis** (Becker, 1901) (*Phora*)  
— Schmitz 1953: 203 (not in Disney 1991)  
[**Anevrina setigera** (Loew, 1874) (*Phora*)]  
**Anevrina thoracica** (Meigen, 1804)  
(*Trineura*) — Schmitz 1928: 141  
**Anevrina unispinosa** (Zetterstedt, 1860)  
(*Trineura*) — Schmitz 1924: 80 (as  
*Parastenophora*; not in Disney 1991)  
**Anevrina urbana** (Meigen, 1830) (*Phora*) —  
Schmitz 1953: 203 (not in Disney 1991)

##### **Beckerina** Malloch, 1910

- Beckerina umbrimargo** (Becker, 1901)  
(*Phora*) — Schmitz 1941a: 11

##### **Borophaga** Enderlein, 1924

- [**Borophaga agilis** (Meigen, 1830) (*Phora*)]  
[**Borophaga carinifrons** (Zetterstedt, 1848)  
(*Phora*)]  
**Borophaga cephalotes** (Schmitz, 1922)  
(*Hypocera*) — Schmitz 1922: 37; type-  
locality: Gyón  
**Borophaga erythrocerca** (Meigen, 1830)  
(*Phora*) — Schmitz 1924: 80 (as *Hypocera*  
*bernuthi* (Egger, 1862))  
**Borophaga femorata** (Meigen, 1830) (*Phora*)  
— Schmitz 1924: 80 (as *Hypocera*  
*flavimana* (Meigen, 1830))  
[**Borophaga germanica** (Schmitz, 1918)  
(*Hypocera*)]  
**Borophaga incrassata** (Meigen, 1830)  
(*Phora*) — Schmitz 1928: 142  
[**Borophaga irregularis** (Wood, 1912)  
(*Hypocera*)]  
[**Borophaga subsultans** (Linnaeus, 1767)  
(*Musca*)]



**Chaetopleurophora** Schmitz, 1922

- [**Chaetopleurophora bohemanni** (Becker, 1901) (*Phora*)]  
**Chaetopleurophora erythronota** (Strobl, 1892) (*Phora*) — Schmitz 1928: 142  
**Chaetopleurophora pygidialis** Schmitz, 1941 — Schmitz 1953: 203 (not in Disney 1991)  
 [**Chaetopleurophora spinosa** Schmitz, 1935]  
**Chaetopleurophora spinosior** Schmitz, 1938 — Schmitz 1940: 120  
**Chaetopleurophora spinosissima** (Strobl, 1892) (*Phora*) — Thalhammer 1899: 36 (as *Phora*); Schmitz 1924: 80

**Conicera** Meigen, 1830

- Conicera dauci** (Meigen, 1830) (*Phora*) — Schmitz 1928: 141 (as *Conicera atra* Meigen, 1830)  
**Conicera floricola** Schmitz, 1938 — Schmitz 1940: 120  
 [**Conicera schnittmanni** Schmitz, 1926]  
 [**Conicera sensilipes** Schmitz, 1938]  
**Conicera similis** (Haliday, 1833) (*Phora*) — Schmitz 1924: 80 (as *Conicera similis* and *C. pauxilla* Schmitz, 1920; not in Disney 1991)  
 [**Conicera tarsalis** Schmitz, 1920]  
**Conicera tibialis** Schmitz, 1925 — Schmitz 1925: 119; type-locality: Pusztá-Szent-Lőrincz

**Diplonevra** Lioy, 1864

- Diplonevra abbreviata** (von Roser, 1840) (*Phora*) — Schmitz 1928: 142 (as *Diplonevra*)  
**Diplonevra altipetax** (Schmitz, 1935) (*Diplonevra*) — Schmitz 1953: 204 (not in Disney 1991)  
 [**Diplonevra amphichaeta** (Schmitz, 1949) (*Diplonevra*)]  
**Diplonevra concinna** (Meigen, 1830) (*Phora*) — Schmitz 1924: 80 (as *Dohrniphora*)  
**Diplonevra crassicornis** (Meigen, 1830) (*Phora*) — Schmitz 1924: 80 (as *Dohrniphora dudai* Schmitz, 1920)  
**Diplonevra florea** (Fabricius, 1794) (*Bibio*) — Thalhammer 1899: 36 (as *Phora florea* and *Ph. flexuosa* Egger, 1862); Schmitz 1924: 80 (as *Dohrniphora abdominalis* (Fallén, 1823))

**[Diplonevra freyi** (Schmitz, 1927)*(Diplonevra)***Diplonevra funebris** (Meigen, 1830) (*Phora*) — Schmitz 1953: 204 (not in Disney 1991)**Diplonevra glabra** (Schmitz, 1927) (*Diplonevra*) — Schmitz 1928: 142 (as *Diplonevra parcepilosa* Schmitz, 1927)**Diplonevra lophochaeta** (Schmitz, 1927) (*Diplonevra*) — Schmitz 1927: 60; type-locality: Gyón**Diplonevra nitidula** (Meigen, 1830) (*Phora*) — Schmitz 1928: 142 (as *Diplonevra*)**[Diplonevra oldenbergi** (Schmitz, 1920) (*Dohrniphora*)]**[Diplonevra pachycera** (Schmitz, 1918) (*Dohrniphora concinna* var.)]**Diplonevra pilosella** (Schmitz, 1927) (*Diplonevra*) — Schmitz 1953: 204 (not in Disney 1991)**[Diplonevra praealpina** (Schmitz, 1948) (*Diplonevra*)]**[Diplonevra refulgens** (Schmitz, 1949) (*Diplonevra*)]**Diplonevra sesquicornis** (Schmitz, 1927) (*Diplonevra*) — Schmitz 1927: 59; type-locality: Budapest**[Diplonevra unisetalis** (Schmitz, 1927) (*Diplonevra*)]**Diplonevra unispinalis** (Schmitz, 1927) (*Diplonevra*) — Schmitz 1927: 60; type-locality: Visegrád**Dohrniphora** Dahl, 1898**Dohrniphora cornuta** (Bigot, 1856) (*Phora*) — Schmitz 1953: 204 (as *Diplonevra*; not in Disney 1991)**Gymnophora** Macquart, 1835**Gymnophora arcuata** (Meigen, 1830) (*Phora*) — Thalhammer 1899: 36; Schmitz 1928: 142**[Gymnophora healeyae** Disney, 1980]**Gymnophora integralis** Schmitz, 1920 — Schmitz 1928: 142**Gymnophora nigripennis** Schmitz, 1926 — Schmitz 1928: 142**Gymnophora quartomollis** Schmitz, 1920 — Schmitz 1928: 142

A species new to science with extremely enlarged male 1st flagellomere has also been found in the newly collected material.

**Gymnoptera** Lioy, 1864

**Gymnoptera longicostalis** Schmitz, 1933 — Schmitz 1940: 128 — This species was split in 1993 so earlier records require checking specimens.

**Gymnoptera vitripennis** (Meigen, 1830) (*Phora*) — Schmitz, 1928: 142 (not mentioned from Hungary in Schmitz's later works; not in Disney 1991)

**Hypocera** Lioy, 1864

**Hypocera mordellaria** (Fallén, 1823) (*Trineura*) — Schmitz 1928: 142 (as *Hypocera subsultans*: Schmitz, 1928)

**Megaselia** Rondani, 1856

[**Megaseli abdita** Schmitz, 1959]

**Megaselia abludens** Schmitz, 1927 — Schmitz 1928: 144 (not in Disney 1991)

[**Megaselia aculeata** (Schmitz, 1919) (*Aphiochaeta*)]

[**Megaselia acuta** Schmitz, 1935]

**Megaselia aequalis** (Wood, 1909) (*Phora*) — Schmitz 1924: 81 (as *Aphiochaeta*)

[**Megaselia affinis** (Wood, 1909) (*Phora*)]

[**Megaselia albicans** (Wood, 1908) (*Phora*)]

**Megaselia albicaudata** (Wood, 1910) (*Phora*) — Schmitz 1924: 83 (as *Aphiochaeta*)

[**Megaselia albiclava** (Schmitz, 1926) (*Aphiochaeta*)]

**Megaselia albocingulata** (Strobl, 1906) (*Phora*) — Schmitz 1924: 83 (as *Aphiochaeta languescens* Schmitz, 1924; type-locality: Gyón)

[**Megaselia alpina** Schmitz, 1965 — Syntype(s) from "Ungarn, Tatra", i.e., it has not been found in Hungary (cf. Disney 1991: 156)]

**Megaselia alticolella** (Wood, 1909) (*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta*)

**Megaselia altifrons** (Wood, 1909) (*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta*)

**Megaselia analis** (Lundbeck, 1920) (*Aphiochaeta*) — Schmitz 1953: 205 (not in Disney 1991)

**Megaselia angelicae** (Wood, 1910) (*Phora*) — Schmitz 1924: 83 (as *Aphiochaeta*)

[**Megaselia angularis** (Schmitz, 1924) (*Aphiochaeta*)]

[**Megaselia annulipes** (Schmitz, 1921) (*Aphiochaeta*)]

[**Megaselia aristica** (Schmitz, 1920) (*Aphiochaeta*)]

[**Megaselia armata** (Wood, 1909) (*Phora*)]

[**Megaselia arquata** Schmitz, 1935]

[**Megaselia atrosericea** Schmitz, 1927]

[**Megaselia badia** Schmitz, 1938]

[**Megaselia barbulate** (Wood, 1909) (*Phora*)]

[**Megaselia basicrinialis** Schmitz, 1953]

**Megaselia basispinata** (Lundbeck, 1920) (*Aphiochaeta*) — Schmitz 1924: 82 (as *Aphiochaeta*)

[**Megaselia basitumida** Schmitz, 1927]

[**Megaselia beatricis** Colyer, 1962]

**Megaselia beckeri** (Wood, 1909) (*Phora*) — Schmitz 1928: 142

**Megaselia berndseni** (Schmitz, 1919) (*Aphiochaeta*) — Schmitz 1924: 83 (as *Aphiochaeta berndseni* and *A. pygmaeoides* Lundbeck, 1921)

[**Megaselia beyeri** Schmitz, 1965]

{**Phora bicolor** Meigen, 1830 — Thalhammer 1899: 36 (doubtful species in *Megaselia*)}

**Megaselia bovista** (Gimmerthal, 1848) (*Phora*) — Schmitz 1924: 83 (as *Aphiochaeta*)

[**Megaselia brevicornis** Schmitz, 1938]

**Megaselia brevicostalis** (Wood, 1910) (*Phora*) — Schmitz 1924: 83 (as *Aphiochaeta*; "von 8 ungarischen Orten")

[**Megaselia brevifemorata** (Schmitz, 1926) (*Aphiochaeta*)]

[**Megaselia brevior** (Schmitz, 1924) (*Aphiochaeta*)]

[**Megaselia brevipipes** (Lundbeck, 1920) (*Aphiochaeta*)]

**Megaselia breviseta** (Wood, 1912) (*Aphiochaeta*) — Schmitz 1924: 81 (as *Aphiochaeta*)

**Megaselia brevissima** (Schmitz, 1924) (*Aphiochaeta*) — Schmitz 1924: 83 (as *Aphiochaeta*; not in Disney 1991)

**Megaselia breviterga** (Lundbeck, 1920) (*Aphiochaeta*) — Schmitz 1953: 205 (not in Disney 1991)

[**Megaselia brunnea** (Schmitz, 1920) (*Aphiochaeta*)]

**Megaselia brunneicornis** (Schmitz, 1920) (*Aphiochaeta*) — Schmitz 1924: 81 (as *Aphiochaeta*)

[**Megaselia brunneipennis** Costa, 1857]

**Megaselia campestris** (Wood, 1908) (*Phora*) — Schmitz 1924: 81 (as *Aphiochaeta*)

[**Megaselia capillipes** Schmitz, 1929]



- [*Megaselia capronata* Schmitz, 1940]  
 [*Megaselia chaetopyga* (Lundbeck, 1921) (*Aphiochaeta*)]  
*Megaselia ciliata* (Zetterstedt, 1848) (*Trineura*) — Schmitz 1924: 81 (as *Aphiochaeta*)  
 [*Megaselia cinerea* Schmitz, 1938]  
*Megaselia cinereifrons* (Strobl, 1910) (*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta*)  
 [*Megaselia cirratula* Schmitz, 1948]  
 [*Megaselia cirricauda* Colyer, 1962]  
 [*Megaselia cirriventris* Schmitz, 1929]  
 [*Megaselia clara* (Schmitz, 1921) (*Aphiochaeta*)]  
 [*Megaselia claricornis* Colyer, 1962]  
 [*Megaselia coacta* (Lundbeck, 1920) (*Aphiochaeta*)]  
 [*Megaselia coetanea* Schmitz, 1929]  
 [*Megaselia coccyx* Schmitz, 1965]  
 [*Megaselia coei* Schmitz, 1938]  
*Megaselia collini* (Wood, 1909) (*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta*)  
*Megaselia communiformis* (Schmitz, 1918) (*Aphiochaeta*) — Schmitz 1928: 142  
*Megaselia conformis* (Wood, 1909) (*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta*)  
*Megaselia consetigera* (Schmitz, 1925) (*Aphiochaeta*) — Schmitz 1928: 143  
 [*Megaselia constricta* Colyer, 1962]  
 [*Megaselia constrictior* Schmitz, 1929]  
 [*Megaselia correlata* (Schmitz, 1918) (*Aphiochaeta*)]  
*Megaselia costalis* (von Roser, 1840) (*Phora*) — Schmitz 1928: 144  
 [*Megaselia cothurnata* (Schmitz, 1919) (*Aphiochaeta*)]  
*Megaselia crassicosta* (Strobl, 1892) (*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta*)  
 [*Megaselia crassilla* Schmitz, 1926]  
 [*Megaselia crassipes* (Wood, 1909) (*Phora*)]  
 [*Megaselia crassivenia* Schmitz, 1927]  
 [*Megaselia crinita* Schmitz, 1939]  
 [*Megaselia criniticauda* Colyer, 1962]  
*Megaselia curvicapilla* Schmitz, 1947 — Schmitz 1953: 204 (not in Disney 1991)  
 [*Megaselia curtivenia* Schmitz, 1928]  
*Megaselia dahli* (Becker, 1901) (*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta*)  
*Megaselia deltomera* (Schmitz, 1924) (*Aphiochaeta*) — Schmitz 1953: 205  
 [*Megaselia densior* Schmitz, 1927]  
 [*Megaselia devia* Schmitz, 1936]  
 [*Megaselia differens* Schmitz, 1948]
- [*Megaselia dimidia* Schmitz, 1926]  
 [*Megaselia dipsacosa* Smith, 1977]  
*Megaselia discreta* (Wood, 1909) (*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta*; not in Disney 1991)  
*Megaselia diversa* (Wood, 1909) (*Phora*) — Schmitz 1953: 205 — Syn.: *Megaselia sordescens* Schmitz, 1927 (Disney 1993)  
 [*Megaselia dubitalis* (Wood, 1908) (*Phora*)]  
 [*Megaselia eisfelderiae* Schmitz, 1948]  
*Megaselia elongata* (Wood, 1914) (*Phora*) — Schmitz 1928: 143 (as *Megaselia cuspidata* (Schmitz, 1919))  
*Megaselia emarginata* (Wood, 1908) (*Phora*) — Schmitz 1924: 81 (as *Aphiochaeta*)  
 [*Megaselia erecta* (Wood, 1910) (*Phora*)]  
*Megaselia errata* (Wood, 1912) (*Phora*) — Schmitz 1924: 81 (as *Aphiochaeta*)  
 [*Megaselia exangulata* Schmitz, 1947]  
*Megaselia fenestralis* (Schmitz, 1919) (*Aphiochaeta*) — Schmitz 1924: 82 (as *Aphiochaeta*)  
 [*Megaselia feronia* Schmitz, 1934]  
 [*Megaselia filamentosa* Schmitz, 1958]  
*Megaselia flava* (Fallén, 1823) (*Trineura*) — Schmitz 1924: 82 (as *Aphiochaeta*)  
*Megaselia flavescens* (Wood, 1909) (*Phora*) — Schmitz 1928: 143  
*Megaselia flavicans* Schmitz, 1935 — Schmitz 1941a: 17  
*Megaselia flavicoxa* (Zetterstedt, 1848) (*Trineura*) — Schmitz 1924: 81 (as *Aphiochaeta*)  
 [*Megaselia frameata* Schmitz, 1927]  
*Megaselia frontalis* (Wood, 1909) (*Phora*) — Schmitz 1928: 144  
 [*Megaselia fumata* (Malloch, 1909) (*Phora*)]  
*Megaselia fungivora* (Wood, 1909) (*Phora*) — Schmitz 1928: 143  
 [*Megaselia furcatipennis* Schmitz, 1934]  
 [*Megaselia furcilla* (Schmitz, 1956) (*Plastophora*)]  
 [*Megaselia furva* Schmitz, 1929]  
*Megaselia fusca* (Wood, 1909) (*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta*; not in Disney 1991)  
 [*Megaselia fuscinervis* (Wood, 1908) (*Phora*)]  
 [*Megaselia fuscinaula* (Schmitz, 1926) (*Aphiochaeta*)]  
 [*Megaselia fuscipalpis* (Lundbeck, 1920) (*Aphiochaeta*)]  
 [*Megaselia fuscopleuralis* Schmitz, 1929]

- [**Megaselia fuscovariana** Schmitz, 1933]  
**Megaselia giraudii** (Egger, 1862) (*Phora*) — Schmitz 1924: 81 (as *Aphiochaeta rata* (Wood, 1908))
- [**Megaselia glabrifrons** (Wood, 1909) (*Phora*)]
- [**Megaselia gregaria** (Wood, 1910) (*Phora*)]  
**Megaselia groenlandica** (Lundbeck, 1901) (*Phora*) — Disney 1991: 167
- Megaselia halterata** (Wood, 1910) (*Phora*) — Schmitz 1924: 83 (as *Aphiochaeta*)
- [**Megaselia hendersoni** Disney, 1979]  
**Megaselia hibernans** Schmitz, 1934]  
**Megaselia hilaris** Schmitz, 1927 — Schmitz 1928: 144
- Megaselia hirsuta** (Wood, 1910) (*Phora*) — Schmitz 1928: 143 (not in Disney 1991)
- Megaselia hirticrus** (Schmitz, 1918) (*Aphiochaeta*) — Schmitz 1953: 205 (not in Disney 1991)
- Megaselia hirtiventris** (Wood, 1909) (*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta*)
- [**Megaselia hortensis** (Wood, 1909) (*Phora*)]  
**Megaselia humeralis** (Zetterstedt, 1838) (*Trineura*) — Schmitz 1924: 81 (as *Aphiochaeta*)
- Megaselia hyalipennis** (Wood, 1912) (*Aphiochaeta*) — Schmitz 1924: 82 (as *Aphiochaeta*)
- Megaselia hypopygialis** (Lundbeck, 1920) (*Aphiochaeta*) — Schmitz 1924: 82 (as *Aphiochaeta*)
- [**Megaselia ignobilis** (Schmitz, 1919) (*Aphiochaeta*)]  
**Megaselia impinguata** Schmitz, 1935]  
**Megaselia incongruens** Schmitz, 1940]  
**Megaselia incontaminata** (Schmitz, 1926) (*Aphiochaeta*)
- Megaselia incrassata** (Schmitz, 1920) (*Aphiochaeta*) — Schmitz 1928: 144
- Megaselia indifferens** (Lundbeck, 1920) (*Aphiochaeta*) — Schmitz 1924: 82 (as *Aphiochaeta*)
- [**Megaselia indigesta** (Schmitz, 1920) (*Aphiochaeta*)]  
**Megaselia infrapospita** (Wood, 1909) (*Phora*)]
- [**Megaselia inquinata** Schmitz, 1953]  
**Megaselia insecta** Schmitz, 1953 — Schmitz 1953: 207 (not in Disney 1991)
- [**Megaselia insons** (Lundbeck, 1920) (*Aphiochaeta*)]
- [**Megaselia intercostata** (Lundbeck, 1921) (*Aphiochaeta*)]  
**Megaselia intersecta** Schmitz, 1935]  
**Megaselia intonsa** Schmitz, 1948]  
**Megaselia involuta** (Wood, 1910) (*Phora*) — Schmitz 1924: 83 (as *Aphiochaeta*)
- [**Megaselia lactipennis** (Lundbeck, 1920) (*Aphiochaeta*)]  
**Megaselia laeviceps** Schmitz, 1948]  
**Megaselia largifrontalis** Schmitz, 1939]  
**Megaselia lata** (Wood, 1910) (*Phora*) — Schmitz 1924: 83 (as *Aphiochaeta*)
- Megaselia latericia** Schmitz, 1935 — Schmitz 1928: 145 (not in Disney 1991)
- Megaselia latifemorata** (Becker, 1901) (*Phora*) — Schmitz 1924: 81 (as *Aphiochaeta*)
- Megaselia latifrons** (Wood, 1910) (*Phora*) — Schmitz 1928: 143
- [**Megaselia latior** Schmitz, 1936]  
**Megaselia latipalpis** (Schmitz, 1921) (*Aphiochaeta*) — Schmitz 1953: 205 (not in Disney 1991)
- [**Megaselia limburgensis** (Schmitz, 1918) (*Aphiochaeta*)]  
**Megaselia longicostalis** (Wood, 1912) (*Aphiochaeta*) — Schmitz 1924b: 131 (“bedarf der Nachprüfung”, Schmitz, 1953: 205; not in Disney 1991)
- Megaselia longipalpis** (Wood, 1910) (*Phora*) — Schmitz 1924: 83 (as *Aphiochaeta*)
- Megaselia longiseta** (Wood, 1909) (*Phora*) — Schmitz 1928: 145
- Megaselia lucifrons** (Schmitz, 1918) (*Aphiochaeta*) — Schmitz 1924: 82 (as *Aphiochaeta subnitida* Lundbeck, 1920; not in Disney 1991)
- Megaselia luminifrons** (Schmitz, 1926) (*Aphiochaeta*) — Disney 1991: 171
- Megaselia lutea** (Meigen, 1930) (*Phora*) — Schmitz 1928: 145
- Megaselia luteipes** (Schmitz, 1918) (*Aphiochaeta*) — Schmitz 1924: 81 (as *Aphiochaeta*; not in Disney 1991)
- Megaselia lutescens** (Wood, 1910) (*Phora*) — Schmitz 1928: 145
- [**Megaselia magnifica** (Lundbeck, 1920) (*Aphiochaeta*)]  
**Megaselia major** (Wood, 1912) (*Aphiochaeta*) — Schmitz 1953: 205
- Megaselia mallochi** (Wood, 1909) (*Phora*) — Schmitz 1924: 81 (as *Aphiochaeta*)



- Megaselia manicata** (Wood, 1910) (*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta*)
- [**Megaselia manualis** (Schmitz, 1919) (*Aphiochaeta*)]
- Megaselia maura** (Wood, 1910) (*Phora*) — Disney 1991: 172
- Megaselia meconicera** (Speiser, 1925) (*Aphiochaeta*) — Schmitz 1953: 205 (not in Disney 1991)
- Megaselia meigeni** (Becker, 1901) (*Phora*) — Schmitz 1924: 81 (as *Aphiochaeta*)
- Megaselia melanocephala** (von Roser, 1840) (*Phora*) — Schmitz 1953: 205 (not in Disney 1991)
- [**Megaselia melanostola** Schmitz, 1942]
- [**Megaselia miki** Schmitz, 1929]
- Megaselia minor** (Zetterstedt, 1848) (*Trineura*) — Thalhammer 1899: 36 (as *Phora*); Schmitz 1924: 81 (as *Aphiochaeta minor* and *A. angustifrons* Wood, 1912)
- Megaselia minuta** (Aldrich, 1892) (*Phora*) — Schmitz 1953: 205 (as *Megaselia luminosa* Schmitz, 1952)
- Megaselia mixta** (Schmitz, 1918) (*Aphiochaeta*) — Schmitz 1928: 145
- [**Megaselia montana** Schmitz, 1935]
- [**Megaselia mortenseni** (Lundbeck, 1920) (*Aphiochaeta*)]
- Megaselia nasoni** (Malloch, 1912) (*Aphiochaeta*) — Schmitz 1924: 81 (as *Aphiochaeta coaequalis* Schmitz, 1919)
- Megaselia nigra** (Meigen, 1830) (*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta albidohalteris*; Schmitz, 1924)
- Megaselia nigriceps** (Loew, 1866) (*Phora*) — Schmitz 1928: 143 (as *Megaselia projecta* (Becker, 1901))
- [**Megaselia nigripalpis** (Lundbeck, 1920) (*Aphiochaeta*)]
- {**Phora nitidifrons** Strobl, 1892 — Thalhammer 1899: 36 (doubtful species in *Megaselia*)}
- [**Megaselia norica** Schmitz, 1929]
- Megaselia oblongifrons** Schmitz, 1939 — Schmitz 1953: 205 (not in Disney 1991)
- Megaselia obscuripennis** (Wood, 1909) (*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta*)
- Megaselia ochripes** Schmitz, 1953 — Schmitz 1953: 206; type-locality: Kőszeg
- [**Megaselia offuscata** (Schmitz, 1921) (*Aphiochaeta*)]
- [**Megaselia oldenbergi** (Schmitz, 1919) (*Aphiochaeta*)]
- [**Megaselia oxybelorum** Schmitz, 1928]
- [**Megaselia pachydactyla** Schmitz, 1953]
- [**Megaselia pallidizona** (Lundbeck, 1920) (*Aphiochaeta*)]
- Megaselia palmeni** (Becker, 1901) (*Phora*) — Schmitz 1928: 143
- Megaselia paludosa** (Wood, 1908) (*Phora*) — Schmitz 1924: 81 (as *Aphiochaeta*)
- Megaselia parva** (Wood, 1909) (*Phora*) — Schmitz 1924: 81 (as *Aphiochaeta*)
- [**Megaselia paupera** (Lundbeck, 1920) (*Aphiochaeta*)]
- [**Megaselia pectinifera** Schmitz, 1926]
- [**Megaselia pectoraliformis** Colyer, 1962]
- Megaselia pectoralis** (Wood, 1910) (*Phora*) — Schmitz 1928: 143
- [**Megaselia pectorella** Schmitz, 1929]
- Megaselia pedatella** (Schmitz, 1926) (*Aphiochaeta*) — Schmitz 1928: 143
- Megaselia perdistans** (Schmitz, 1924) (*Aphiochaeta*) — Schmitz 1928: 145
- [**Megaselia perfusca** Schmitz, 1935]
- [**Megaselia petraea** Schmitz, 1934]
- [**Megaselia phoenicura** (Schmitz, 1926) (*Aphiochaeta*)]
- Megaselia picta** (Lehmann, 1822) (*Phora*) — Schmitz 1924: 81 (as *Aphiochaeta*)
- [**Megaselia picotorufa** (Colyer, 1957) (*Plastophora*)]
- [**Megaselia pilifemur** (Lundbeck, 1921) (*Aphiochaeta*)]
- Megaselia pleuralis** (Wood, 1909) (*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta*)
- Megaselia plurispinulosa** (Zetterstedt, 1860) (*Trineura*) — Schmitz 1924: 81 (as *Aphiochaeta giraudii*; Schmitz, 1924)
- Megaselia posticata** (Strobl, 1898) (*Phora*) — Disney 1991: 177
- Megaselia praeacuta** (Schmitz, 1919) (*Aphiochaeta*) — Schmitz 1953: 205 (not in Disney 1991)
- [**Megaselia prodroma** (Lundbeck, 1921) (*Aphiochaeta*)]
- [**Megaselia producta** (Schmitz, 1921) (*Aphiochaeta*) — Syn.: *Megaselia sordescens* auctt., nec Schmitz, 1927 (Disney 1993)]
- [**Megaselia prolongata** Schmitz, 1954]
- Megaselia propinqua** (Wood, 1909) (*Phora*) — Schmitz 1924: 81 (as *Aphiochaeta*)
- [**Megaselia propior** Colyer, 1956]

- [*Megaselia protarsalis* Schmitz, 1927]  
 [*Megaselia pseudociliata* (Strobl, 1910) (*Aphiochaeta*)]  
*Megaselia pseudogiraudii* (Schmitz, 1920) (*Aphiochaeta*) — Schmitz 1924: 81 (as *Aphiochaeta*)  
 [*Megaselia pubecula* Schmitz, 1927]  
*Megaselia pulicaria* (Fallén, 1823) (*Trineura*) — Thalhammer 1899: 36 (as *Phora*); Schmitz 1924: 82 (as *Aphiochaeta pulicaria* and *A. angusta* (Wood, 1909))  
 [*Megaselia pullipalpis* Colyer, 1962]  
*Megaselia pumila* (Meigen, 1830) (*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta atripes* Brues, 1915)  
*Megaselia pusilla* (Meigen, 1830) (*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta*)  
*Megaselia pygmaea* (Zetterstedt, 1848) (*Trineura*) — Schmitz 1924: 83 (as *Aphiochaeta*)  
*Megaselia quadriseta* (Schmitz, 1918) (*Aphiochaeta*) — Schmitz 1924: 81 (as *Aphiochaeta*)  
*Megaselia rivalis* (Wood, 1909) (*Phora*) — Schmitz 1928: 144 (as *Megaselia ebenina* (Schmitz, 1920); not mentioned from Hungary in Schmitz's later works; not in Disney 1991)  
 [*Megaselia robusta* Schmitz, 1928]  
*Megaselia rubella* (Schmitz, 1920) (*Aphiochaeta*) — Schmitz 1928: 145  
*Megaselia rubescens* (Wood, 1912) (*Aphiochaeta*) — Schmitz 1928: 145  
*Megaselia rubida* (Schmitz, 1918) (*Aphiochaeta*) — Schmitz 1924: 81 (as *Aphiochaeta*)  
 [*Megaselia rubicornis* (Schmitz, 1919) (*Aphiochaeta*)]  
*Megaselia rudis* (Wood, 1909) (*Phora*) — Schmitz 1924: 81 (as *Aphiochaeta*)  
*Megaselia rufa* (Wood, 1908) (*Phora*) — Schmitz 1924: 81 (as *Aphiochaeta*)  
*Megaselia ruficornis* (Meigen, 1830) (*Phora*) — Schmitz 1924: 81 (as *Aphiochaeta*)  
 [*Megaselia rufifrons* (Wood, 1910) (*Phora*)]  
*Megaselia rufipes* (Meigen, 1804) (*Trineura*) — Thalhammer 1899: 36 (as *Phora pulicaria* v. *rufipes* and *Ph. pulicaria* v. *heracleellae* Bouché, 1834); Schmitz 1924: 82 (as *Aphiochaeta*)  
 [*Megaselia rupestris* Schmitz, 1934]  
 [*Megaselia sanguinea* (Schmitz, 1922) (*Aphiochaeta*)]  
 [*Megaselia scalaris* (Loew, 1866) (*Phora*)]  
 [*Megaselia schutti* Schmitz, 1936]  
*Megaselia scutellaris* (Wood, 1909) (*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta*)  
 [*Megaselia sepulchralis* (Lundbeck, 1920) (*Aphiochaeta*)]  
 [*Megaselia sericata* Schmitz, 1935]  
 [*Megaselia serrata* (Wood, 1910) (*Phora*)]  
 [*Megaselia setifer* (Lundbeck, 1920) (*Aphiochaeta*)]  
 [*Megaselia setulipalpis* Schmitz, 1938]  
 [*Megaselia sicaria* (Colyer, 1962) (*Plastophora*)]  
*Megaselia similifrons* Schmitz, 1934 — Schmitz 1941a: 34  
 [*Megaselia simplex* (Wood, 1910) (*Phora*)]  
*Megaselia simulans* (Wood, 1912) (*Aphiochaeta*) — Schmitz 1953: 205  
 [*Megaselia sinuata* Schmitz, 1926]  
 [*Megaselia solitaria* Schmitz, 1934]  
*Megaselia sordida* (Zetterstedt, 1838) (*Trineura*) — Thalhammer 1899: 36 (as *Phora*); Schmitz 1924: 82 (as *Aphiochaeta*; not in Disney 1991)  
 [*Megaselia specularis* Schmitz, 1935]  
 [*Megaselia speiseri* Schmitz, 1929]  
 [*Megaselia spinata* (Wood, 1910) (*Phora*)]  
*Megaselia spinicincta* (Wood, 1910) (*Phora*) — Schmitz 1928: 145  
 [*Megaselia spinigera* (Wood, 1908) (*Phora*)]  
 [*Megaselia stichata* (Lundbeck, 1920) (*Aphiochaeta*)]  
 [*Megaselia stigmatica* (Schmitz, 1920) (*Aphiochaeta*)]  
 [*Megaselia striolata* Schmitz, 1940]  
*Megaselia styloprocta* (Schmitz, 1921) (*Aphiochaeta*) — Schmitz 1924: 81 (as *Aphiochaeta*)  
 [*Megaselia subcarpalis* (Lundbeck, 1920) (*Aphiochaeta*)]  
 [*Megaselia subconvexa* (Lundbeck, 1920) (*Aphiochaeta*)]  
*Megaselia subfraudulenta* Schmitz, 1933 — Schmitz 1953: 205 (not in Disney, 1991)  
 [*Megaselia subfuscipes* Schmitz, 1935]  
*Megaselia subnudipennis* (Schmitz, 1919) (*Aphiochaeta*) — Schmitz 1928: 145  
 [*Megaselia subpalpalis* (Lundbeck, 1920) (*Aphiochaeta*)]  
*Megaselia subpleuralis* (Wood, 1909) (*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta*)



- Megaselia subtumida** (Wood, 1909) (*Phora*)  
— Schmitz 1924: 82 (as *Aphiochaeta*; not in Disney 1991)
- Megaselia sulphuripes** (Meigen, 1830)  
(*Phora*) — Schmitz 1924: 82 (as *Aphiochaeta*; "von 8 ungarischen Orten")
- Megaselia superciliata** (Wood, 1910) (*Phora*)  
— Schmitz 1928: 144
- Megaselia surdifrons** (Wood, 1909) (*Phora*)  
— Schmitz 1941a: 34
- Megaselia sylvatica** (Wood, 1910) (*Phora*) —  
Schmitz 1924: 82 (as *Aphiochaeta*  
*impolluta* Schmitz, 1920)
- [**Megaselia tabida** Colyer, 1956]
- Megaselia tama** (Schmitz, 1926)  
(*Aphiochaeta*) — Schmitz 1928: 144
- Megaselia tarsalis** (Wood, 1910) (*Phora*) —  
Schmitz 1924: 83 (as *Aphiochaeta*)
- [**Megaselia tarsella** (Lundbeck, 1921)  
(*Aphiochaeta*)]
- [**Megaselia tarsicia** Schmitz, 1926]
- [**Megaselia tenebricola** Schmitz, 1934]
- [**Megaselia tergata** (Lundbeck, 1920)  
(*Aphiochaeta*)]
- [**Megaselia testacea** Schmitz, 1938]
- [**Megaselia thalhammeri** Schmitz, 1935]
- [**Megaselia trichorrhoea** (Schmitz, 1921)  
(*Aphiochaeta*)]
- [**Megaselia triquetra** Schmitz, 1927]
- Megaselia tumida** (Wood, 1909) (*Phora*) —  
Schmitz 1924: 82 (as *Aphiochaeta*)
- [**Megaselia uliginosa** (Wood, 1909) (*Phora*)]
- Megaselia unguicularis** (Wood, 1909)  
(*Phora*) — Schmitz 1928: 144 (not  
mentioned from Hungary in Schmitz's later  
works)
- [**Megaselia unichaeta** Colyer, 1962]
- [**Megaselia unicolor** (Schmitz, 1919)  
(*Aphiochaeta*)]
- [**Megaselia ustulata** (Schmitz, 1920)  
(*Aphiochaeta*)]
- [**Megaselia valvata** Schmitz, 1935]
- Megaselia variana** Schmitz, 1926 — Schmitz  
1924: 82 (as *Aphiochaeta variabilis*:  
Schmitz, 1924)
- [**Megaselia verna** Schmitz, 1932]
- [**Megaselia vernalis** (Wood, 1909) (*Phora*)]
- Megaselia verralli** (Wood, 1910) (*Phora*) —  
Schmitz 1953: 205 (not in Disney 1991)
- Megaselia vestita** (Wood, 1914)  
(*Aphiochaeta*) — Schmitz 1924: 83 (as  
*Aphiochaeta*)
- [**Megaselia villicauda** Schmitz, 1927]
- [**Megaselia virilis** (Schmitz, 1919)  
(*Aphiochaeta*)]
- [**Megaselia waagei** Schmitz, 1935]
- Megaselia woodi** (Lundbeck, 1922)  
(*Aphiochaeta*) — Schmitz 1924: 82 (as  
*Aphiochaeta*; not in Disney 1991)
- Megaselia xanthozona** (Strobl, 1892) (*Phora*)  
— Thalhammer 1899: 36 (as *Phora*);  
Schmitz 1924: 82 (as *Aphiochaeta*)
- Megaselia zonata** (Zetterstedt, 1838)  
(*Trineura*) — Schmitz 1924: 82 (as  
*Aphiochaeta*)
- Menoziola** Schmitz, 1927
- Menoziola schmitzi** (Menozi, 1921)  
(*Apocephalus*) — Schmitz 1928: 141
- Metopina** Macquart, 1835
- Metopina braueri** (Strobl, 1880)  
(*Drepanophora*) — Schmitz 1953: 206 (not  
in Disney 1991)
- [**Metopina crassinervis** Schmitz, 1920]
- [**Metopina formicomendicola** Schmitz, 1927]
- Metopina galeata** (Haliday, 1833) (*Phora*) —  
Schmitz 1924: 83 (not in Disney 1991)
- [**Metopina heselhausi** Schmitz, 1914]
- [**Metopina oligoneura** (Mik, 1867) (*Phora*)]
- Metopina perpusilla** (Six, 1878) (*Leptophora*)  
— Schmitz 1941a: 35 (not in Disney 1991)
- [**Metopina pilieta** Schmitz, 1936]
- [**Metopina trochanteralis** Schmitz, 1953]
- [**Metopina ulrichi** Disney, 1979]
- Phalacrotophora** Schmitz, 1934
- Phalacrotophora berolinensis** Schmitz, 1920  
— Schmitz 1924: 83
- Phalacrotophora fasciata** (Fallén, 1823)  
(*Trineura*) — Schmitz 1953: 206 (as  
*Phalacrotophora*; not in Disney 1991)
- [**Phalacrotophora paradoxa** Schmitz, 1949]
- [**Phalacrotophora pictofasciata** Schmitz,  
1919]
- Phalacrotophora spectabilis** Schmitz, 1925  
— Schmitz 1928: 146
- Phora** Latreille, 1796
- [**Phora adducta** Schmitz, 1955]
- [**Phora artifrons** Schmitz, 1920]
- Phora atra** (Meigen, 1804) (*Trineura*) —  
Thalhammer 1899: 36 (as *Trineura aterrima*)

- (Fabricius, 1794); Schmitz 1924: 81 (as *Phora aterrime* (Fabricius, 1794))
- [*Phora bullata* Schmitz, 1927]
- [*Phora convallium* Schmitz, 1928]
- [*Phora convergens* Schmitz, 1920]
- Phora dubia* (Zetterstedt, 1848) (*Trineura*) — Schmitz 1928: 146 (as *Phora schineri* (Becker, 1901))
- Phora edentata* Schmitz, 1920 — Schmitz 1924: 81
- Phora hamata* Schmitz, 1927 — Schmitz 1953: 203
- Phora holosericea* Schmitz, 1920 — Schmitz 1928: 146
- Phora horrida* Schmitz, 1920 — Schmitz 1928: 146
- [*Phora hyperborea* Schmitz, 1927]
- [*Phora indivisa* Schmitz, 1948]
- [*Phora obscura* (Zetterstedt, 1848) (*Trineura*)]
- [*Phora penicillata* Schmitz, 1920]
- [*Phora pubipes* Schmitz, 1920]
- Phora stictica* Meigen, 1830 — Schmitz 1928: 146
- Phora tinctoria* Schmidt, 1920 — Schmitz 1928: 146
- Phora velutina* Meigen, 1830 — Schmitz 1953: 203 (not in Disney 1991)
- Plectanocnema** Schmitz, 1926
- Plectanocnema nudipes* (Becker, 1901) (*Phora*) — A genus new to Hungary. One male: Budapest, Pestszentlőrinc, Péterhalmi-erdő, tölgyes [= oak forest], 19. VIII. 1995, L. Papp; 1 male: Bükki N. P.: Szarvaskő, 30. VII. 1984, L. Papp
- Pseudacteon** Coquillett, 1907
- [*Pseudacteon brevicauda* Schmitz, 1925]
- [*Pseudacteon fennicus* Schmitz, 1927]
- Pseudacteon formicarum* (Verrall, 1877) (*Phora*) — Schmitz 1924: 83
- Pseudacteon lundbecki* Schmitz, 1924 — Schmitz 1953: 206 (not in Disney 1991)
- Spiniphora** Malloch, 1909
- Spiniphora bergenstammi* (Mik, 1864) (*Phora*) — Schmitz 1953: 203 (not in Disney 1991). Found again in Hungary (4 individuals)
- [*Spiniphora dorsalis* (Becker, 1901) (*Phora*)]
- [*Spiniphora excisa* (Becker, 1901) (*Phora*)]
- [*Spiniphora jugorum* (Schmitz, 1924) (*Paraspiniphora*)]
- Spiniphora maculata* (Meigen, 1830) (*Phora*) — Schmitz 1928: 146. Found again: Budapest, Pestszentlőrinc, Péterhalmi-erdő
- Spiniphora strobli* (Becker, 1901) (*Phora*) — Schmitz 1928: 146
- [*Stichillus* Enderlein, 1924]
- [*Stichillus coronatus* (Becker, 1901) (*Phora*)]
- Triphleba** Rondani, 1856
- [*Triphleba aequalis* (Schmitz, 1919) (*Trupheoneura*)]
- Triphleba antricola* (Schmitz, 1918) (*Pseudostenophora*) — Schmitz 1928: 146
- Triphleba aprilina* (Schmitz, 1918) (*Trupheoneura*) — Schmitz 1924: 80 (as *Trupheoneura radiosetosa* Schmitz, 1918)
- [*Triphleba aptina* (Schiner, 1853) (*Phora*)]
- [*Triphleba autumnalis* (Becker, 1901) (*Phora*)]
- [*Triphleba bicornuta* (Strobl, 1910) (*Phora*)]
- [*Triphleba bifida* Schmitz, 1949]
- [*Triphleba citreiformis* (Becker, 1901) (*Phora*)]
- [*Triphleba crassinervis* (Strobl, 1910) (*Phora*)]
- [*Triphleba dentata* Schmitz, 1943]
- Triphleba distinguenda* (Strobl, 1892) (*Phora*) — Schmitz 1928: 147 (as *Triphleba unicalcarata* (Becker, 1901))
- Triphleba dudai* (Schmitz, 1918) (*Trupheoneura*) — Schmitz 1924: 80 (as *Trupheoneura pauciseta* Schmitz, 1918)
- [*Triphleba excisa* (Lundbeck, 1921) (*Trupheoneura*)]
- [*Triphleba flexipalpis* Schmitz, 1927]
- [*Triphleba forfex* Schmitz, 1943]
- [*Triphleba gracilis* (Wood, 1907) (*Phora*)]
- [*Triphleba hentrichi* Schmitz, 1934]
- Triphleba hyalinata* (Meigen, 1830) (*Phora*) — Schmitz 1928: 146
- [*Triphleba hypopygialis* (Schmitz, 1918) (*Trupheoneura*)]
- [*Triphleba inaequalis* Schmitz, 1943]
- [*Triphleba intempesta* (Schmitz, 1918) (*Trupheoneura*)]
- Triphleba intermedia* (Malloch, 1908) (*Phora*) — Schmitz 1924: 80 (as *Trupheoneura*)



- [*Triphleba latipalpis* (Schmitz, 1924)  
(*Trupheoneura*)]
- Triphleba longifurcata*** (Schmitz, 1922)  
(*Parastenophora*) — Schmitz 1922: 34;  
type-locality: Gyón
- Triphleba lugubris*** (Meigen, 1830) (*Phora*)  
— Schmitz 1924: 80 (as *Trupheoneura*; not  
in Disney 1991)
- [*Triphleba luteifemorata* (Wood, 1906)  
(*Phora*)]
- [*Triphleba lyria* Schmitz, 1935]
- [*Triphleba minuta* (Fabricius, 1787) (*Empis*)]
- [*Triphleba novembrina* Schmitz, 1943]
- Triphleba nudipalpis*** (Becker, 1901)  
(*Trupheoneura*) — Schmitz 1953: 204 (not  
in Disney 1991)
- Triphleba opaca*** (Meigen, 1830) (*Phora*) —  
Thalhammer 1899: 36 (as *Phora nigricornis*  
Egger, 1862); Schmitz 1924: 80 (as  
*Trupheoneura*)
- [*Triphleba pachyneurella* (Schmitz, 1919)  
(*Trupheoneura*)]
- [*Triphleba palposa* (Zetterstedt, 1848)  
(*Trineura*)]
- Triphleba papillata*** (Wingate, 1906) (*Phora*)  
— Schmitz 1928: 146
- [*Triphleba smithi* Disney, 1982]
- [*Triphleba subcompleta* Schmitz, 1927]
- Triphleba transparens*** (Schmitz, 1922)  
(*Parastenophora*) — Schmitz 1928: 146
- Triphleba trinervis*** (Becker, 1901) (*Phora*)  
— Schmitz 1940: 128
- [*Triphleba tumidula* (Schmitz, 1918)  
(*Trupheoneura*)]
- [*Triphleba vitrea* (Wood, 1906) (*Phora*)]
- [*Triphleba zernyi* Schmitz, 1927]
- Tubicera** Schmitz, 1920
- Tubicera lichtwardti*** Schmitz, 1920 —  
Schmitz 1924: 83
- Woodiphora** Schmitz, 1926
- [***Woodiphora retroversa*** (Wood, 1908)  
(*Phora*)]
- A species was found in our material from  
Hungary but it has not been identified.

## LITERATURE

- Disney, R. H. L. (1983): Scuttle Flies. Diptera Phoridae (except Megaselia). — In: *Handbk Ident. Br. Insects* Vol. 10, Part 6. Royal Entomological Society of London, pp. 81.
- Disney, R. H. L. (1989): Scuttle Flies. Diptera Phoridae. Genus Megaselia. — In: *Handbk Ident. Br. Insects* Vol. 10, Part 8. Royal Entomological Society of London, pp. 155.
- Disney, R. H. L. (1991): Family Phoridae. — In: Soós, Á. and Papp, L. (eds): *Catalogue of Palaearctic Diptera*. Volume 7. Dolichopodidae–Platypzeidae. Akadémiai Kiadó, Budapest, pp. 143–204.
- Disney, R. H. L. (1993): Notes on European Phoridae (Diptera). — *Br. J. Ent. Nat. Hist.* 6: 107–118.
- Schmitz, H. (1922): Neue europäische Phoriden des Ungarischen Nationalmuseums. — *Ent. Ber., Amst.* 6: 33–39.
- Schmitz, H. (1924a): Europäische Phoriden des Ungarischen National-Museums. — *Annls hist.-nat. Mus. natn hung.* 21: 79–86.
- Schmitz, H. (1924b): Nachtrag zu den Verzeichnis der Phoriden des Ungarischen Nationalmuseums. — *Natuurh. Maandbl.* 13: 131.
- Schmitz, H. (1925): Drei neue europäische Phoriden des Ungarischen National-Museums. — *Annls hist.-nat. Mus. natn hung.* 22: 119–123.
- Schmitz, H. (1928): Die geographische verbreitung der europaeischen Phoridenarten. — *Natuurh. Maandbl.* 17: 139–147.
- Schmitz, H. (1938): 33. Phoridae. — In: Lindner, E. (ed.): *Die Fliegen der palaearktischen Region*. 4 (7), Lieferung 138, pp. 1–64.
- Schmitz, H. (1940): Kritisches Verzeichnis der paläarktischen Phoriden. Mit Angabe ihrer Verbreitung. — *Natuurh. Maandbl.* 29: 96, 101–104, 118–120, 127–129.

- Schmitz, H. (1941a): Kritisches Verzeichnis der paläarktischen Phoriden. Mit Angabe ihrer Verbreitung. — *Natuurh. Maandbl.* 30: 11–12, 15–17, 34–35.
- Schmitz, H. (1941b): 33. Phoridae. — In: Lindner, E. (ed.): *Die Fliegen der palaearktischen Region.* 4 (7), Lieferung 141, pp. 65–128.
- Schmitz, H. (1943a): 33. Phoridae. — In: Lindner, E. (ed.): *Die Fliegen der palaearktischen Region.* 4 (7), Lieferung 147, pp. 129–160.
- Schmitz, H. (1943b): 33. Phoridae. — In: Lindner, E. (ed.): *Die Fliegen der palaearktischen Region.* 4 (7), Lieferung 149, pp. 161–192.
- Schmitz, H. (1949): 33. Phoridae. — In: Lindner, E. (ed.): *Die Fliegen der palaearktischen Region.* 4 (7), Lieferung 160, pp. 193–240.
- Schmitz, H. (1951): 33. Phoridae. — In: Lindner, E. (ed.): *Die Fliegen der palaearktischen Region.* 4 (7), Lieferung 165, pp. 241–272.
- Schmitz, H. (1952): 33. Phoridae. — In: Lindner, E. (ed.): *Die Fliegen der palaearktischen Region.* 4 (7), Lieferung 171, pp. 273–320.
- Schmitz, H. (1953): Ungarische und andere paläarktische Phoriden des Ungarischen Nationalmuseums (Diptera). — *Annl. hist.-nat. Mus. natn. hung.* 3: 203–211.
- Schmitz, H. (1955): 33. Phoridae. — In: Lindner, E. (ed.): *Die Fliegen der palaearktischen Region.* 4 (7), Lieferung 180, pp. 321–386.
- Schmitz, H. (1956): 33. Phoridae. — In: Lindner, E. (ed.): *Die Fliegen der palaearktischen Region.* 4 (7), Lieferung 187, pp. 387–416.
- Schmitz, H. (1957): 33. Phoridae. — In: Lindner, E. (ed.): *Die Fliegen der palaearktischen Region.* 4 (7), Lieferung 196, pp. 417–464.
- Schmitz, H. (1958): 33. Phoridae. — In: Lindner, E. (ed.): *Die Fliegen der palaearktischen Region.* 4 (7), Lieferung 202, pp. 465–512.
- Schmitz, H. und Beyer, E. M. (1965a): 33. Phoridae. — In: Lindner, E. (ed.): *Die Fliegen der palaearktischen Region.* 4 (7), Lieferung 258, pp. 513–560.
- Schmitz, H. und Beyer, E. M. (1965b): 33. Phoridae. — In: Lindner, E. (ed.): *Die Fliegen der palaearktischen Region.* 4 (7), Lieferung 260, pp. 561–608.
- Schmitz, H. und Delage, A. (1974): 33. Phoridae. — In: Lindner, E. (ed.): *Die Fliegen der palaearktischen Region.* 4 (7), Lieferung 301, pp. 609–664.
- Schmitz, H. und Delage, A. (1981): 33. Phoridae. — In: Lindner, E. (ed.): *Die Fliegen der palaearktischen Region.* 4 (7), Lieferung 325, pp. 665–712.
- Thalhammer, J. (1899): Ordo. Diptera. — In: *A Magyar Birodalom Állatvilága. A Magyar Birodalomból eddig ismert állatok rendszeres lejtroma.* A K. M. Természettudományi Társulat, Budapest, 1–76 pp.

(Received: May 3, 1996)

Authors' address: L. ÁDÁM and L. PAPP  
Department of Zoology  
Hungarian Natural History Museum  
Baross u. 13  
H-1088 Budapest  
HUNGARY