

AKADÉMIAI KIADÓ

Journal of Behavioral Addictions

13 (2024) 4, 1003–1013

DOI:  
10.1556/2006.2024.00053  
© 2024 The Author(s)

FULL-LENGTH REPORT



# Rethinking cutoff values for the South Oaks Gambling Screen: Sex-specific insights and DSM-5 severity adjustments in Gambling Disorder assessment

IGNACIO LUCAS<sup>1,2</sup> , ROSER GRANERO<sup>2,3,4</sup> ,  
BERNAT MORA-MALTAS<sup>2,5</sup> ,  
FERNANDO FERNÁNDEZ-ARANDA<sup>1,2,3,6</sup> and  
SUSANA JIMÉNEZ-MURCIA<sup>1,2,3,6\*</sup>

<sup>1</sup> Clinical Psychology Department, Bellvitge University Hospital, Barcelona, Spain

<sup>2</sup> Psychoneurobiology of Eating and Addictive Behaviours Group, Bellvitge Biomedical Research Institute (IDIBELL), Barcelona, Spain

<sup>3</sup> CIBER de Fisiopatología de la Obesidad y Nutrición (CIBERObn), Instituto de Salud Carlos III, Barcelona, Spain

<sup>4</sup> Departament de Psicobiologia i Metodologia, Universitat Autònoma de Barcelona, Barcelona Spain

<sup>5</sup> Doctoral Program in Medicine and Translational Research, University of Barcelona, Barcelona, Spain

<sup>6</sup> Department of Clinical Sciences, School of Medicine and Health Sciences, University of Barcelona, Barcelona, Spain

Received: March 14, 2024 • Revised manuscript received: August 26, 2024 • Accepted: August 31, 2024  
Published online: October 4, 2024

## ABSTRACT

**Background and aims:** The South Oaks Gambling Screen (SOGS) is a commonly used tool for screening potential gambling problems. The SOGS score has been found to be correlated with the DSM-5 criteria for Gambling Disorder (GD). However, one of its main limitations is the high rate of false positives. The objective of this study was to establish more accurate cutoff points for the SOGS based on the DSM-5 criteria and severity levels, taking into account sex-specific samples. Additionally, we aimed to validate these cutoff points using external measures. **Methods:** The study sample comprised 4,516 patients (398 women and 4,118 men) who sought treatment for GD. Of these patients, 4,203 met the DSM-5 criteria for GD, while 313 did not meet the threshold for GD diagnosis. **Results:** The recommended cutoff value for the SOGS is eight for men (Sensitivity (Se): 82.9%, Specificity (Sp): 86.2%) and seven for women (Se: 85.6%, Sp: 77.4%). For moderate severity of GD, the cutoff points are nine for men (Se: 82.1%, Sp: 82.0%) and eight for women (Se: 86.3%, Sp: 73.3%), while for severe cases of GD, the cutoff point is ten for both sexes (Se: 83.0%, Sp: 56.7% for men; Se: 80.0%, Sp: 77.4% for women). These cutoff values are validated by evidence of worse psychopathological states, more dysfunctional personality traits, and risky behaviours commonly associated with GD. **Discussion and Conclusions:** These findings support adjusting the reference values for the SOGS to eight in males and seven in females to enhance the classification of individuals potentially experiencing GD. The use of higher cutoff values has significant implications for clinical and research purposes, enabling a more precise assessment of the diagnosis and severity of GD.

## KEYWORDS

Receiver Operating Characteristics (ROC), gambling disorder, South Oaks Gambling Screen (SOGS), DSM-5, severity

\*Corresponding author.  
E-mail: [sjimenez@bellvitgehospital.cat](mailto:sjimenez@bellvitgehospital.cat)

## INTRODUCTION

The South Oaks Gambling Screen (SOGS) (Lesieur & Blume, 1987) is a widely used instrument designed for screening the possible presence of gambling problems in clinical,



epidemiological and research contexts (Petry, 2005). Its structure is based on the third edition of the Diagnostic and Statistical Manual of Mental Disorders III (DSM-III) criteria for pathological gambling, the former version of current gambling disorder (GD) diagnosis of the DSM-5 and DSM-5-TR (American Psychiatric Association, 2013, 2022). Despite being designed based on DSM-III criteria, the SOGS has a similar etiological structure of later versions of the manual (Slutske, Zhu, Meier, & Martin, 2011) and demonstrated to be a rapid and reliable instrument to evaluate the presence of gambling related problems (Esparza-Reig, Guillén-Riquelme, Martí-Vilar, & González-Sala, 2021; Stinchfield, 2002). The scale is composed of 20 items with multiple choice, assessing a single dimension. The score ranges from 0 to 20, and its original cutoff point is 5, therefore a score from 5 to 20 is considered as an indicator of the possible presence of gambling related problems. The SOGS allows its administration by professionals, non-professionals not specialized in psychological assessment, administration by computer or even self-administration, and has been adapted and validated to several languages and cultures, showing good psychometric properties (de Oliveira, Silva, & da Silveira, 2002; Duvarci, Varan, Coşkunol, & Ersoy, 1997; Echeburúa, Báez, Fernández-Montalvo, & Páez, 1994; Gyollai et al., 2011; Haddad, Roukoz, Akel, & Hallit, 2022; Kido & Shimazaki, 2007; Petry, Armentano, Kuoch, Norinth, & Smith, 2003; Tang, Wu, Tang, & Yan, 2010).

In clinical settings, screening instruments are very useful, as they provide reliable, valid information about the possible presence of a disorder and are frequently the first step towards the solution of a serious problem. In this line, easy to administer screening instruments are crucial in order to optimize public health resources providing the adequate assistance to each patient considering its individual characterization (precision medicine). Nevertheless, like almost all brief assessment instruments, the SOGS has some limitations and, hence, has not been exempt of criticism. For instance, half of the SOGS items pertain to the criterion of borrowing money to gamble. As a result, the mere presence of this particular criterion could lead to a positive screening score (Stinchfield, 2002). Additionally, other studies suggest that the SOGS may be assessing two distinctive dimensions related to gambling issues (Orford, Sproston, & Erens, 2003; Salonen et al., 2017). Furthermore, one of the primary critiques of the SOGS is the high presence of false positives, i.e. many patients surpassing the threshold but not presenting the condition (Battersby, Thomas, Tolchard, & Esterman, 2002; Otto et al., 2020; Stinchfield, 2002; Tang et al., 2010).

Goodie et al. (2013) examined the utility of the SOGS to predict the presence of gambling disorder in a sample of community individuals who presented frequent gambling behaviours. Their results yielded a high number of false positives of GD for the original cutoff value of five. However, the authors proposed that if the cutoff was elevated to eight, the number of false positives was reduced drastically while it has almost no impact in the false negative rates. In the same line, previous validity studies also reported that the SOGS cutoff value should be elevated to eight (de Oliveira et al.,

2009; Duvarci & Varan, 2001; Tang et al., 2010). Moreover, the SOGS scores range up to 20, and these cutoff points consider the same risk for all patients who scored above them. Hence, these proposals aim solely at providing a cutoff point to assess the possible presence of a gambling problem. The fifth version of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5) introduced three different severity levels to the diagnosis of GD. These three severity levels are determined by the number of fulfilled criteria: mild (4–5 criteria), moderate (6–7 criteria) and severe (8–9 criteria) and were established mirroring those proposed for substance use disorders (SUD). To specify the severity of the disorder is a useful feature to predict the prognosis of the disorder and, therefore, to adapt the treatment to the necessities of each patient. The SOGS scores correlate with the number of DSM criteria for GD (Cox, Enns, & Michaud, 2004; de Oliveira et al., 2009; Goodie et al., 2013; Stinchfield, 2002; Tang et al., 2010), with other measures of gambling frequency and severity (Barbaranelli, Vecchione, Fida, & Podio-Guidugli, 2013; Petry, 2003), and with more impulsive choices in a delay discounting paradigm (Alessi & Petry, 2003). Therefore, the SOGS have been widely used as a measure of severity of GD (Aloi et al., 2022; Chase & Clark, 2010; Ciccarelli, Cosenza, Nigro, & D'Olimpio, 2022; Jiménez-Murcia et al., 2019; Mathieu, Barrault, Brunault, & Varescon, 2020; Muela, Ventura-Lucena, Navas, & Perales, 2023; Nigro et al., 2022; Panayiotou, Artemi, Theodorou, Theodorou, & Neophytou, 2023; Rogier, Picci, & Velotti, 2019), even though it was not initially designed for this purpose (Lesieur & Blume, 1987). Other widely used screening tools for addiction problems, like the Alcohol Use Disorders Identification Test (AUDIT) (Saunders, Aasland, Babor, de la Fuente, & Grant, 1993) and the Drug Use Disorders Identification Test (DUDIT) (Berman, Bergman, Palmstierna, & Schlyter, 2005) have different cutoff points for different risk levels.

Moreover, the AUDIT and the DUDIT also have different cutoff scores depending on sex, being slightly lower for women than for men (Babor, Higgins-Biddle, Saunders, Monteiro, & World Health Organization (WHO), 2001; Berman et al., 2005; Hildebrand, 2015; Reinert & Allen, 2007). Nevertheless, the SOGS do not have different scores by sex although men and women tend to exhibit different profiles of GD. Women are more prone to engage in non-strategic forms of gambling (e.g. bingo, lotteries), while men are more likely to conduct strategic gambling (e.g. poker, sports betting) (Gartner, Bickl, Härtl, Loy, & Häffner, 2022; Grant, Chamberlain, Schreiber, & Odlaug, 2012; Jiménez-Murcia, Granero, Fernández-Aranda, & Menchón, 2020; Zakiniaiez & Potenza, 2018). Additionally, women present higher comorbidity with depression and anxiety symptomatology (Gartner et al., 2022; Grant, Chamberlain, et al., 2012). In contrast, men have less previous comorbidity with depression and anxiety problems but higher comorbid alcohol and substance use (Gartner et al., 2022; Grant, Chamberlain, et al., 2012). In addition, women usually have a later onset, but a more rapid evolution of the disorder (i.e. telescoping phenomenon) (Grant, Odlaug, &



Mooney, 2012; Lucas et al., 2023; Zakiniaieiz, Cosgrove, Mazure, & Potenza, 2017), as well as higher rates in autolytic ideation and behaviour (Valenciano-Mendoza et al., 2021). Unlike physiological conditions where gender differences are clear-cut, the behavioural context of gambling presents a more complex scenario. However, the sex-specific differences in gambling behaviour, preferences, and clinical outcomes suggest that separate cutoffs may improve the sensitivity and specificity of the SOGS.

Therefore, the objectives of this study are, first, to establish cutoff values for the SOGS that align with the diagnostic criteria and the severity levels for GD outlined in the DSM-5, while accounting for variations in these cutoff values based on the sex of the assessed individuals. Second, we aim to assess the discriminant validity of the proposed cutoff values in relation to psychopathological variables and gambling behaviour.

## METHODS

### Participants

The sample consisted of 4,516 patients (398 women and 4,118 men) who sought treatment in the Behavioural Addictions Unit of the University Hospital of Bellvitge, a public hospital in Spain certified as a tertiary care centre for the treatment of GD. All of them were assessed by experienced clinical psychologists in two sessions before the treatment. During the first session, the clinical psychologist conducted a semi-structured interview to assess the presence of GD and to explore various aspects of gambling behaviour and socio-demographic data. During this first session, they also signed the informed consent to participate in the study. During the second assessment session, participants completed a battery of validated psychometric instruments, including the Diagnostic Questionnaire for Pathological Gambling (Jiménez-Murcia et al., 2009; Stinchfield, 2003). Both the semi-structured interview and the Diagnostic Questionnaire for Pathological Gambling were used to ensure diagnostic accuracy. 4,203 patients met the threshold for GD ( $\geq 4$  criteria of the DSM-5), while 313 of them did not reach the threshold for GD diagnosis and, accordingly, their condition was considered as problematic gambling. The recruitment took place between January-2005 and March-2023.

### Instruments

**Semi-structured clinical interview.** The aim of this interview was to assess clinical and sociodemographic variables (Jiménez-Murcia, Aymamí-Sanromà, Gómez-Pena, Álvarez-Moya, & Vallejo, 2006), including the age at which problematic gambling behaviour began, the presence of accumulated debts related to gambling, the occurrence of illegal acts related to gambling disorder, gambling preferences (non-strategic, strategic, or mixed), type of gambling (offline, online, or mixed) (Jiménez-Murcia et al., 2020), substance use, autolytic behaviour, suicidal ideation, age, marital status, highest level of education attained,

employment status, personal income, and family income. Social position was calculated using Hollingshead's index (Hollingshead, 2011).

**South Oaks Gambling Screen.** As described in the introduction, the SOGS is a widely used screening instrument for assessing the possibility of a gambling problem (Lesieur & Blume, 1987). It consists of 20 items that assess gambling symptoms and possible negative consequences. SOGS scores have showed correlation with DSM criteria for GD (Cox et al., 2004; de Oliveira et al., 2009; Goodie et al., 2013; Stinchfield, 2002; Tang et al., 2010) and other measures of gambling severity (Barbaranelli et al., 2013; Petry, 2003). The Spanish version of the SOGS was used in this study (Echeburúa et al., 1994) and the measured internal consistency was in the adequate range, with a Chronbach's alpha of 0.703.

**Symptom Checklist-Revised.** The Symptom Checklist-Revised (SCL-90-R) (Derogatis, 1994) is a self-report instrument designed to assess psychopathological symptoms. The instrument uses a 5-point Likert scale and consists of 90 items. It provides dimensions of somatization, obsessive-compulsive symptoms, interpersonal sensitivity, depression, anxiety, hostility, phobic anxiety, paranoid ideation and psychotic symptomatology. In addition, the instrument provides three global measures: Global Severity Index (GSI), Positive Symptom Total (PST) and Positive Symptom Distress Index (PSDI). The Spanish version used in this study (Derogatis, 2002) showed good internal consistency, with Chronbach's alphas ranging from 0.783 to 0.980 for all dimensions.

**Temperament and Character Inventory-Revised.** The Temperament and Character Inventory-Revised (TCI-R) (Cloninger, 1999) is a self-report instrument designed for the evaluation of temperament and character traits. The instrument uses a 5-point Likert scale and consists of 240 items. The temperament dimensions include harm avoidance, novelty seeking, reward dependence and persistence, while the character dimensions include self-directedness, cooperativeness and self-transcendence. The Spanish version used in this study (Gutiérrez-Zotes et al., 2004) had a Cronbach's alpha between 0.703 and 0.859 for all dimensions.

**DSM-5 criteria for gambling disorder.** Diagnostic criteria for GD were assessed using the Spanish adaptation of the Diagnostic Questionnaire for Pathological Gambling (Jiménez-Murcia et al., 2009; Stinchfield, 2003). This self-report measure consists of 19 items coded on a binary scale (Yes/No). This instrument is a valid and reliable instrument to evaluate the presence of DSM-IV criteria for pathological gambling. The results were re-evaluated and recoded post hoc according to the DSM-5 criteria for GD. The internal consistency for this study was adequate, with a Chronbach's alpha of 0.761.

### Statistical analysis

The statistical analysis was done with Stata18 for Windows (Stata-Corp, 2023). The analysis of the global accuracy of the



SOGS as a screening measure to discriminate the GD severity level (as defined by DSM-5 criteria, defined as the external reference gold standard in this work) was done with Receiver Operating Characteristics (ROC) methodology. The area under the ROC curve (AUC) measured the global accuracy of the SOGS across all the potential cutoff points. The optimal cutoff point for the SOGS was selected considering the usual indexes: sensitivity (Se), specificity (Sp), false alarm rate (FAR, the complement of Sp), predictive negative value (PNV), predictive positive value (PPV), and false discovery rate (FDR, the complement of PPV). Since the primary aim of the screening tools is to maximize the accuracy to identify at-high-risk individuals (with the assumption that the costs of false negative classifications are higher than false positive ones), the selection of the optimal cutoff point was based on Se indexes no lower than 0.80 (Zhou, Obuchowski, & McClish, 2002).

Next, two groups based on the SOGS optimal cutoff point were compared (low screening score *versus* high screening score), using chi-square tests ( $\chi^2$ ) for categorical variables and analysis of variance (ANOVA) for quantitative variables. These procedures included the calculation of the effect size for the differences, with the standardized indexes Cramer's- $V$  and partial eta-square ( $\eta_p^2$ ) (mild-moderate effect size was considered for  $V > 0.15$  and  $\eta_p^2 > 0.10$ , and high-large effect for  $V > 0.30$  and  $\eta_p^2 > 0.25$ ) (Levine & Hullett, 2002). In addition, Finner's procedure was used for controlling the increase in Type-I error due to multiple significance tests (Finner & Gontscharuk, 2009).

## Ethics

This study was conducted in accordance with the Declaration of Helsinki. The study was approved by the Ethics Committee for Clinical Research of the University Hospital of Bellvitge (ref. 34/05, 307/06).

## RESULTS

### Description of the sample

Mean age was 42.4 years-old ( $SD = 13.8$ ). Most participants in the study were married (44.2%), with low education levels (58.2% primary or less), employed (56.2%), and pertained to low social position index (51.6%). A detailed description of the sociodemographic variables of the sample is displayed in Table 1, which includes the frequency distribution by sex.

### Severity distribution

Based on the DSM-5 criteria,  $n = 4,203$  patients (93.1%) met clinical criteria for GD and  $n = 313$  were within the problematic gambling condition ( $<4$  criteria). No difference by sex was observed for this classification ( $\chi^2 = 0.50$ ,  $p = 0.480$ ,  $V = 0.01$ ).

Figure 1 shows the bar-chart with the distribution of the gambling severity based on the DSM-5 criteria. The highest severity level achieved the prevalence in the study, with no differences between sexes (GD:  $\chi^2 = 2.99$ ,  $p = 0.225$ ,  $V = 0.03$ ; problematic gambling:  $\chi^2 = 0.40$ ,  $p = 0.821$ ,  $V = 0.04$ ).

### Cutoff points

Figure 2 displays the results of the ROC analysis aimed at the selection of the optimal cutoff points of the SOGS for discriminating between the GD severity levels (each plot shows the impact of the different SOGS cutoff points on the Se, Sp, PPV, PNV, FDR and FAR). Among the women subsample, the best SOGS divisions were 7 (for mild GD), 8 (for moderate GD) and 10 (for severe GD). Among men subsamples, the best divisions were 8, 9 and 10 (for mild, moderate and severe GD). Table 2 displays the complete

Table 1. Sociodemographic descriptive of the sample

|                                     |                      | Total            |           | Women          |           | Men              |           | <i>p</i>      |
|-------------------------------------|----------------------|------------------|-----------|----------------|-----------|------------------|-----------|---------------|
|                                     |                      | <i>N</i> = 4,516 |           | <i>N</i> = 398 |           | <i>N</i> = 4,118 |           |               |
| <sup>1</sup> Quantitative variables |                      | <i>Mean</i>      | <i>SD</i> | <i>Mean</i>    | <i>SD</i> | <i>Mean</i>      | <i>SD</i> |               |
| Age (years-old)                     |                      | 42.37            | 13.82     | 49.98          | 12.81     | 41.64            | 13.70     | <b>0.001*</b> |
| <sup>1</sup> Categorical variables  |                      | <i>n</i>         | %         | <i>n</i>       | %         | <i>n</i>         | %         |               |
| Marital status                      | Single               | 1,904            | 42.2%     | 170            | 42.7%     | 1,734            | 42.1%     | <b>0.001*</b> |
|                                     | Married - in couple  | 1,997            | 44.2%     | 149            | 37.4%     | 1,848            | 44.9%     |               |
|                                     | Divorced - separated | 615              | 13.6%     | 79             | 19.8%     | 536              | 13.0%     |               |
| Education                           | Primary/less         | 2,629            | 58.2%     | 245            | 61.6%     | 2,384            | 57.9%     | 0.188         |
|                                     | Secondary            | 1,593            | 35.3%     | 124            | 31.2%     | 1,469            | 35.7%     |               |
|                                     | University           | 294              | 6.5%      | 29             | 7.3%      | 265              | 6.4%      |               |
| Employment                          | Unemployed           | 1,978            | 43.8%     | 215            | 54.0%     | 1,763            | 42.8%     | <b>0.001*</b> |
|                                     | Employed             | 2,538            | 56.2%     | 183            | 46.0%     | 2,355            | 57.2%     |               |
| Social position index               | High                 | 67               | 1.5%      | 3              | 0.8%      | 64               | 1.6%      | <b>0.001*</b> |
|                                     | Mean-high            | 216              | 4.8%      | 15             | 3.8%      | 201              | 4.9%      |               |
|                                     | Mean                 | 467              | 10.3%     | 44             | 11.1%     | 423              | 10.3%     |               |
|                                     | Mean-low             | 1,436            | 31.8%     | 77             | 19.3%     | 1,359            | 33.0%     |               |
|                                     | Low                  | 2,330            | 51.6%     | 259            | 65.1%     | 2,071            | 50.3%     |               |

Note. <sup>1</sup>Comparisons of means with analysis of variance (ANOVA). <sup>2</sup>Comparison of proportions with chi-square tests ( $\chi^2$ ). SD: standard deviation. Bold:  $p < 0.05$ .








Fig. 1. Severity distribution based on DSM-5 criteria

Note. GD present: mild (4–5 criteria), moderate (6–7 criteria), severe (8–9 criteria).  
Problematic gambling: mild (1 criteria), moderate (2 criteria), severe (3 criteria).




Fig. 2. Identification of the optimal cutoff for the SOGS: ROC analysis for women (left) and men (right)

Note. Se: sensitivity. Sp: specificity. PPV: predictive value for a positive. PNV: predictive value for a negative. FDR: False discovery rate. FAR: false alarm rate.




Table 2. Identification of the optimal cutoff for the SOGS: ROC analysis

| Optimal cutoff for the SOGS: ROC analysis |             |        |        |       |       |        |       |       |       |        |       |
|-------------------------------------------|-------------|--------|--------|-------|-------|--------|-------|-------|-------|--------|-------|
| Subsample                                 | DSM-5       | Cutoff | Se     | Sp    | PPV   | PNV    | FDR   | FAR   | AUC   | 95% CI | AUC   |
| Women ( <i>n</i> = 398)                   | GD mild     | 7      | 85.6%  | 77.4% | 97.8% | 31.8%  | 22.6% | 2.2%  | 0.752 | 0.652  | 0.851 |
|                                           | GD moderate | 8      | 86.3%  | 73.3% | 90.2% | 65.4%  | 26.7% | 9.8%  | 0.810 | 0.760  | 0.859 |
|                                           | GD Severe   | 10     | 80.0%  | 77.4% | 75.7% | 81.0%  | 22.6% | 24.3% | 0.782 | 0.736  | 0.827 |
| Men ( <i>n</i> = 4,118)                   | GD mild     | 8      | 82.9%  | 86.2% | 98.8% | 92.2%  | 13.8% | 1.2%  | 0.828 | 0.804  | 0.851 |
|                                           | GD moderate | 9      | 82.1%  | 82.0% | 93.2% | 60.4%  | 18.0% | 6.8%  | 0.840 | 0.826  | 0.854 |
|                                           | GD Severe   | 10     | 83.0%  | 56.7% | 60.1% | 81.0%  | 43.3% | 39.9% | 0.799 | 0.786  | 0.813 |
| Validity for the cutoff = 5               |             |        |        |       |       |        |       |       |       |        |       |
| Subsample                                 | DSM-5       | Cutoff | Se     | Sp    | PPV   | PNV    | FDR   | FAR   | AUC   | 95% CI | AUC   |
| Women ( <i>n</i> = 398)                   | GD mild     | 5      | 97.0%  | 29.0% | 94.0% | 45.7%  | 71.0% | 6.0%  | 0.752 | 0.652  | 0.851 |
|                                           | GD moderate | 5      | 99.0%  | 23.8% | 78.7% | 89.1%  | 76.2% | 21.3% | 0.810 | 0.760  | 0.859 |
|                                           | GD Severe   | 5      | 100.0% | 14.2% | 50.8% | 100.0% | 85.8% | 49.2% | 0.782 | 0.736  | 0.827 |
| Men ( <i>n</i> = 4,118)                   | GD mild     | 5      | 98.1%  | 36.5% | 95.4% | 97.9%  | 63.5% | 4.6%  | 0.828 | 0.804  | 0.851 |
|                                           | GD moderate | 5      | 99.6%  | 24.0% | 79.7% | 95.0%  | 76.0% | 20.3% | 0.840 | 0.826  | 0.854 |
|                                           | GD Severe   | 5      | 99.9%  | 12.5% | 47.3% | 99.7%  | 87.5% | 52.7% | 0.799 | 0.786  | 0.813 |

Note. Se: sensitivity. Sp: specificity. PPV: predictive value for a positive screening score. PNV: predictive value for a negative screening score. FDR: false discovery rate. FAR: false alarm rate. AUC: area under the ROC curve. 95% CI: 95% confidence interval.

assessment of these selected thresholds for the SOGS. This table also includes the validity-accuracy indexes for a cutoff equal to 5, which provide unacceptable results in specificity, regardless of the severity of the gambling disorder and the sex of the patient.

### Discriminant validity of the optimal cutoff points

The patients were classified in two groups based on the cutoff point which discriminated at least mild GD severity level (SOGS = 7 for women and SOGS = 8 for men). The results of the comparisons between the individuals with low versus high score is displayed in Table 3. Among women, those with high SOGS screening score reported a younger onset of the GD, higher number of DSM-5 criteria for GD, worse psychopathology state (higher mean scores in the SCL90-R), higher scores in the novelty seeking and harm avoidance traits, lower score in the self-directedness, higher likelihood of alcohol consumption and higher risk of illegal behaviour related with the gambling activity. Among men, those with a high SOGS screening score reported a younger onset of the GD, longer duration of the GD, higher number of DSM-5 criteria for GD, worse psychopathology state, more dysfunctional personality traits, higher likelihood of impulsivity behaviours (autolytic and suicidal behaviour, as well as substance use), and a higher risk of debts due to the gambling activity.

## DISCUSSION

The findings of this research agree with raising the cutoff score of the SOGS from five to eight in order to determine the probable presence of gambling disorder in men who show gambling activity. The precise threshold of eight has previously been suggested as the optimal cutoff value for the

SOGS (de Oliveira et al., 2009; Duvarci & Varan, 2001; Goodie et al., 2013; Tang et al., 2010). However, this cutoff value had still not been validated with a large sample of treatment-seeking patients in any prior study. Additionally, this study assessed if different cutoff values could be recommended for men and women, as proposed in other screening instruments for addictive disorders (Babor et al., 2001; Berman et al., 2005; Hildebrand, 2015; Reinert & Allen, 2007). According to these results, the ideal cutoff value for women should be seven, one point less than for men. The literature that explored sex differences in GD has identified different profiles of GD for men and women (Gartner et al., 2022; Grant, Chamberlain, et al., 2012; Jiménez-Murcia et al., 2020; Zakiniaiez & Potenza, 2018). The lower cutoff value for women is consistent with these differential profiles, as women exhibit a more rapid evolution of the disorder and, thus, may be prompt to present GD with fewer symptoms as assessed by the SOGS (Grant, Odlaug, & Mooney, 2012; Lucas et al., 2023; Zakiniaiez et al., 2017).

Moreover, these results propose different scores depending on the level of severity of GD. For men, the cutoff value for assessing moderate severity would be nine, while for women this cutoff value would be eight, in line with the different cutoff scores for possible GD. However, according to these results, no different cutoff values either for men or women should be considered for severe GD. A SOGS score equal or greater than ten may represent a high risk of presenting severe GD for both men and women. A cutoff of ten in the SOGS had already been proposed to precisely identify severe gambling problems (Battersby et al., 2002; Dickerson, Baron, Hong, & Cottrell, 1996). These results support the use of the SOGS to assess GD severity from a dimensional perspective (Barbaranelli et al., 2013; Goodie et al., 2013; Petry, 2003; Stinchfield, 2002; Tang et al., 2010).



Table 3. Discriminant validity of the proposed cutoff points

| SOGS →                 | <sup>1</sup> Women |       |                |       |               |                             | <sup>2</sup> Men |       |                  |       |               |                             |
|------------------------|--------------------|-------|----------------|-------|---------------|-----------------------------|------------------|-------|------------------|-------|---------------|-----------------------------|
|                        | Low; n = 63        |       | High.; n = 335 |       |               |                             | Low; n = 799     |       | High.; n = 3,319 |       |               |                             |
|                        | Mean               | SD    | Mean           | SD    | p             | η <sub>p</sub> <sup>2</sup> | Mean             | SD    | Mean             | SD    | p             | η <sub>p</sub> <sup>2</sup> |
| Onset GD (yrs-old)     | 41.62              | 14.31 | 36.98          | 11.89 | <b>0.006*</b> | 0.019                       | 33.50            | 14.26 | 28.50            | 11.08 | <b>0.001*</b> | 0.027                       |
| Duration GD (yrs)      | 5.62               | 5.45  | 6.06           | 6.07  | 0.592         | 0.001                       | 5.45             | 6.14  | 6.27             | 6.09  | <b>0.001*</b> | 0.003                       |
| DSM-5 total            | 5.03               | 2.02  | 7.12           | 1.90  | <b>0.001*</b> | <b>0.137</b>                | 4.84             | 1.81  | 7.23             | 1.75  | <b>0.001*</b> | <b>0.223<sup>†</sup></b>    |
| SCL: Somatization      | 1.31               | 0.90  | 1.71           | 0.91  | <b>0.002*</b> | 0.025                       | 0.60             | 0.59  | 0.97             | 0.78  | <b>0.001*</b> | 0.038                       |
| SCL: Obs/compulsive    | 1.13               | 0.84  | 1.62           | 0.87  | <b>0.001*</b> | 0.042                       | 0.74             | 0.66  | 1.21             | 0.80  | <b>0.001*</b> | 0.056                       |
| SCL: Interp. sensitive | 0.96               | 0.82  | 1.53           | 0.87  | <b>0.001*</b> | 0.054                       | 0.65             | 0.65  | 1.09             | 0.82  | <b>0.001*</b> | 0.045                       |
| SCL: Depressive        | 1.61               | 0.98  | 2.21           | 0.89  | <b>0.001*</b> | 0.056                       | 0.98             | 0.74  | 1.56             | 0.88  | <b>0.001*</b> | 0.069                       |
| SCL: Anxiety           | 1.03               | 0.86  | 1.62           | 0.89  | <b>0.001*</b> | 0.055                       | 0.59             | 0.59  | 1.07             | 0.79  | <b>0.001*</b> | 0.058                       |
| SCL: Hostility         | 0.65               | 0.66  | 1.17           | 0.86  | <b>0.001*</b> | 0.050                       | 0.58             | 0.63  | 0.99             | 0.84  | <b>0.001*</b> | 0.039                       |
| SCL: Phobic anxiety    | 0.63               | 0.77  | 0.91           | 0.88  | <b>0.020*</b> | 0.014                       | 0.27             | 0.45  | 0.50             | 0.66  | <b>0.001*</b> | 0.021                       |
| SCL: Paranoid id.      | 0.94               | 0.83  | 1.33           | 0.83  | <b>0.001*</b> | 0.029                       | 0.61             | 0.63  | 0.98             | 0.78  | <b>0.001*</b> | 0.036                       |
| SCL: Psychotic         | 0.77               | 0.70  | 1.25           | 0.79  | <b>0.001*</b> | 0.051                       | 0.53             | 0.57  | 0.97             | 0.76  | <b>0.001*</b> | 0.054                       |
| SCL: PST score         | 1.09               | 0.75  | 1.59           | 0.74  | <b>0.001*</b> | 0.058                       | 0.67             | 0.54  | 1.11             | 0.69  | <b>0.001*</b> | 0.066                       |
| SCL: GSI score         | 42.73              | 21.58 | 58.11          | 17.52 | <b>0.001*</b> | 0.087                       | 32.97            | 19.65 | 48.48            | 20.47 | <b>0.001*</b> | 0.084                       |
| SCL: PSDI score        | 2.03               | 0.57  | 2.32           | 0.62  | <b>0.001*</b> | 0.029                       | 1.65             | 0.50  | 1.92             | 0.57  | <b>0.001*</b> | 0.035                       |
| TCI: Novelty seeking   | 102.51             | 12.97 | 111.15         | 11.68 | <b>0.001*</b> | 0.066                       | 102.50           | 13.13 | 110.70           | 12.77 | <b>0.001*</b> | 0.060                       |
| TCI: Harm avoidance    | 104.86             | 16.90 | 110.09         | 15.40 | <b>0.015*</b> | 0.015                       | 96.97            | 14.35 | 100.58           | 16.12 | <b>0.001*</b> | 0.008                       |
| TCI: Reward-Depend.    | 101.84             | 14.14 | 102.15         | 12.38 | 0.858         | 0.000                       | 99.01            | 14.02 | 97.39            | 13.55 | <b>0.003*</b> | 0.002                       |
| TCI: Persistence       | 103.46             | 16.99 | 104.01         | 16.80 | 0.811         | 0.000                       | 107.58           | 18.25 | 108.85           | 18.62 | 0.081         | 0.001                       |
| TCI: Self-directedness | 129.90             | 20.14 | 118.80         | 17.59 | <b>0.001*</b> | 0.048                       | 138.16           | 19.61 | 126.01           | 19.50 | <b>0.001*</b> | 0.057                       |
| TCI: Cooperative.      | 135.75             | 13.17 | 134.12         | 13.45 | 0.378         | 0.002                       | 133.51           | 14.66 | 128.87           | 15.40 | <b>0.001*</b> | 0.014                       |
| TCI: Self-Transcend.   | 68.03              | 14.23 | 68.31          | 14.15 | 0.887         | 0.000                       | 60.65            | 13.85 | 63.41            | 13.94 | <b>0.001*</b> | 0.006                       |
|                        | n                  | %     | n              | %     | p             | V                           | n                | %     | n                | %     | p             | V                           |
| Autolytic behaviour    | 9                  | 14.3% | 50             | 14.9% | 0.896         | 0.007                       | 31               | 3.9%  | 211              | 6.4%  | <b>0.008*</b> | 0.042                       |
| Suicidal ideation      | 11                 | 17.5% | 84             | 25.1% | 0.193         | 0.065                       | 86               | 10.8% | 677              | 20.4% | <b>0.001*</b> | 0.098                       |
| Tobacco                | 28                 | 44.4% | 184            | 54.9% | 0.126         | 0.077                       | 432              | 54.1% | 2,078            | 62.6% | <b>0.001*</b> | 0.069                       |
| Alcohol                | 1                  | 1.6%  | 32             | 9.6%  | <b>0.035</b>  | 0.105                       | 90               | 11.3% | 584              | 17.6% | <b>0.001*</b> | 0.068                       |
| Drugs                  | 1                  | 1.6%  | 17             | 5.1%  | 0.222         | 0.061                       | 50               | 6.3%  | 406              | 12.2% | <b>0.001*</b> | 0.075                       |
| Debts due to gambling  | 22                 | 34.9% | 155            | 46.3% | 0.096         | 0.083                       | 243              | 30.4% | 1,821            | 54.9% | <b>0.001*</b> | <b>0.193<sup>†</sup></b>    |
| Illegal acts           | 6                  | 9.5%  | 71             | 21.2% | <b>0.031*</b> | 0.108                       | 87               | 10.9% | 937              | 28.2% | <b>0.001*</b> | <b>0.159<sup>†</sup></b>    |

Note. <sup>1</sup>High score: SOGS≥7. <sup>2</sup>High score: High score: SOGS≥8. SD: standard deviation. Effect size (η<sub>p</sub><sup>2</sup> for ANOVA and Cramer-V for χ<sup>2</sup> test). \*Bold: significant comparison. <sup>†</sup>Effect size: mild/moderate to high/large (η<sub>p</sub><sup>2</sup> > 0.10 or V > 0.15).

The second objective of this study was to validate the heightened cutoff points by using external measures related to the GD. The discriminant validity of these cutoff values is proven by the higher general psychopathological symptomatology, dysfunctional personality traits, sooner gambling behaviour onset, more alcohol use and more illegal acts in those individuals with a SOGS value higher than eight (for men) and seven (for women). Moreover, in men there are also differences in gambling duration, autolytic behaviours and suicidal ideations, tobacco and drug use and debts. For women, the lack of differences in the GD duration may be associated to the telescoping effect (Grant, Odlaug, & Mooney, 2012; Lucas et al., 2023; Zakiniaieiz et al., 2017). Furthermore, the absence of differences in the rates of suicidal ideation and self-harming behaviour in women could be explained as being independent of the level of severity (Valenciano-Mendoza et al., 2021). Moreover, while substance-related disorders correlate with GD in men, this association is not observed in women (Gartner et al., 2022). Also, the presence of higher debt rates in males with a SOGS

score higher than eight may be associated with the preference for online based strategic forms of gambling (Baenas et al., 2024), associated with a rapid increase in debt levels (Jiménez-Murcia et al., 2020). These differences support the investigation of sex-specific cutoffs to enhance diagnostic accuracy. However, it is essential to exercise caution when interpreting these cutoffs, particularly to prevent the potential stigmatisation of women. Overall, these results provide discriminant validity for these cutoff points, as more severe gambling profile is present in those individuals who score above the proposed thresholds.

Objective identification of potential issues of the instrument is crucial for both clinical and research purposes. In clinical settings, false positives may be considered less severe errors than false negatives (Goodie et al., 2013; Stinchfield, 2002, 2003). This is because untreated medical conditions may have greater adverse effects than the side effects, expenses or other consequences that may result from a false positive diagnosis. Therefore, in clinical settings aiming to identify potential gambling problems, a score of five may



be convenient, but the elevated rates of false positives have to be considered. On the other hand, for epidemiological and research purposes where SOGS is employed as a tool to assess possible gambling severity, it would be recommended to raise the cutoff points to decrease the occurrence of false positives and provide a more accurate depiction of gambling disorder. In line with this and prior research (de Oliveira et al., 2009; Duvarci & Varan, 2001; Goodie et al., 2013; Tang et al., 2010), a fitting SOGS score to differentiate possible positive cases of gambling disorder would be eight for males and, according to these results, seven for females. The findings of this research can assist not only to endorse the increase of the cutoff values in upcoming studies but also to accurately interpret prior research that applied the SOGS to measure the severity of GD in samples with gambling behaviours. Those studies with individuals with gambling behaviour but low SOGS scores ought to contemplate that their sample may exhibit noteworthy differences with clinically diagnosed patients (Otto et al., 2020; Stinchfield, 2002; Tang et al., 2010). Moreover, when categorizing the severity of the GD using the SOGS, the proposed cutoff values for the different severity levels may serve as a reference of the scores of a large clinical sample of patients diagnosed with GD with different severity gradients (Barbaranelli et al., 2013; Goodie et al., 2013; Petry, 2003; Stinchfield, 2002; Tang et al., 2010). In addition, for clinical purposes, a screening score corresponding to a higher severity level could prove beneficial in determining treatment urgency, serving as a form of triage, providing a more personalized intervention and, thus, optimizing the time and resources applied to evaluate possible cases of GD.

## Limitations

The findings of this study have to be interpreted in light of its limitations. Firstly, sex differences have been considered in this study, however, other individual features could have an impact on the optimal cutoff values, such as age (Granero et al., 2020) or cultural context (Battersby et al., 2002; Hayano, Dong, Miyata, & Kasuga, 2021). Secondly, obtained cutoff values for women may be tentative due to sample size limitations. Future research should continue to refine these cutoffs with larger and more diverse samples in the context of gambling behaviour. Thirdly, the severity levels are useful to determine which patients would require a more rapid or a specific intervention, but DSM-5 severity levels for GD are not exempt from criticism, as they have not been proven to be as effective for GD as they have for SUD and not all symptoms may be equivalent or interchangeable (Grant, Odlaug, & Chamberlain, 2017; Hasin et al., 2013; Mestre-Bach et al., 2019). While this study adheres to the current standard, future work should explore alternative approaches to severity assessment that may offer a more refined understanding of GD. Lastly, the entire sample included in this study exhibited at least one DSM-5 symptom, thereby suggesting that investigating this phenomenon including non-problem gamblers could be an interesting possibility for future research, providing a more comprehensive validation of the SOGS cutoffs.

## CONCLUSIONS

This study is the first to validate the cutoff values for the SOGS both for males and females, also adding cutoff values for the different severity gradients of GD proposed in the DSM-5. Moreover, consistent with previous studies, these results support the dimensional use of the SOGS as a measure to assess the severity of GD, but, at the same time, suggest that the reference values of the SOGS should be increased to eight in males and seven in females in order to optimize the categorization of individuals with a potential GD. The rise in cutoff values has significant implications for both clinical practice and research purposes, as it allows for a more precise assessment of GD diagnosis and severity levels.

*Funding sources:* We thank CERCA Programme/Generalitat de Catalunya for guarantee institutional support. This manuscript and research were supported by grants from the Delegación del Gobierno para el Plan Nacional sobre Drogas (2021I031), Plan Nacional sobre Drogas Convocatoria de subvenciones para proyectos de investigación financiados con fondos europeos 2022 (EXP2022/008847), Ministerio de Ciencia e Innovación (PDI2021-124887OB-I00 supported by MCIN/ AEI/10.13039/501100011033 and FEDER “Una manera de hacer Europa”), Instituto de Salud Carlos III (ISCIII) (FIS PI20/00132 and cofounded by FEDER funds/European Regional Development Fund (ERDF), a way to build Europe). CIBERObn is an initiative of ISCIII. Additional funding was received by AGAUR-Generalitat de Catalunya (2021-SGR-00824) and European Union’s Horizon 2020 research and innovation programme under Grant agreement no. 847879 (PRIME/H2020, Prevention and Remediation of Insulin Multimorbidity in Europe) and no. 101080219 (eprObes). IL was supported by the Ministerio de Ciencia e Innovación (Juan de la Cierva-Formación program, FJC2021-046494-I). RG is supported by the Catalan Institution for Research and Advanced Studies (ICREA-Academia, 2021-Programme). The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

*Authors’ contribution:* IL: Conceptualization, Writing-Original draft; RG: Data analysis, Writing-Original draft; BM-M: Conceptualization, Writing-Review and editing; FF-A: Writing—review and editing, Funding acquisition; SJ-M: Conceptualization, Writing—review and editing, Funding acquisition. All authors approved the final version of the manuscript. All authors had full access to all data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

*Conflict of interest:* FF-A and SJ-M received consultancy honoraria from Novo Nordisk. None of these funding sources are related to this study, and the funding institution had no role in the study design or the collection, analysis, and interpretation of the data, the writing of the manuscript, or the decision to submit the paper for publication.





*Data availability:* The datasets generated during and/or analysed during the current study are not publicly available due to ethical restrictions in order to protect the confidentiality of the participants, but are available from the corresponding author on reasonable request.

## REFERENCES

- Alessi, S. M., & Petry, N. M. (2003). Pathological gambling severity is associated with impulsivity in a delay discounting procedure. *Behavioural Processes*, 64(3), 345–354. [https://doi.org/10.1016/S0376-6357\(03\)00150-5](https://doi.org/10.1016/S0376-6357(03)00150-5).
- Aloi, M., Riccelli, C., Piterà, F., Notaro, M., Curcio, V., Pullia, L., ... De Fazio, P. (2022). Impaired metacognitive differentiation, high difficulty in controlling impulses and non-acceptance of emotions are associated with the severity of gambling disorder. *Journal of Gambling Studies*, 39(1), 1–11. <https://doi.org/10.1007/s10899-021-10099-y>.
- American Psychiatric Association (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). American Psychiatric Association.
- American Psychiatric Association (2022). *Diagnostic and statistical manual of mental disorders* (5th ed.). Text Revision: American Psychiatric Association.
- Babor, T. F., Higgins-Biddle, J. C., Saunders, J. B., & Monteiro, M. G., & World Health Organization (WHO) (2001). *AUDIT: The alcohol use disorders identification test: Guidelines for use in primary health care*. World Health Organization.
- Baenas, I., Mora-Maltas, B., Etxandi, M., Lucas, I., Granero, R., Fernández-Aranda, F., ... Jiménez-Murcia, S. (2024). Cluster analysis in gambling disorder based on sociodemographic, neuropsychological, and neuroendocrine features regulating energy homeostasis. *Comprehensive Psychiatry*, 128(October 2023), 152435. <https://doi.org/10.1016/j.comppsy.2023.152435>.
- Barbaranelli, C., Vecchione, M., Fida, R., & Podio-Guidugli, S. (2013). Estimating the prevalence of adult problem gambling in Italy with SOGS and PGSI. *Journal of Gambling Issues*, 28, 1–24. <https://doi.org/10.4309/jgi.2013.28.3>.
- Battersby, M. W., Thomas, L. J., Tolchard, B., & Esterman, A. (2002). The South Oaks Gambling Screen: A review with reference to Australian use. *Journal of Gambling Studies*, 18(3), 257–271. <https://doi.org/10.1023/a:1016895221871>.
- Berman, A. H., Bergman, H., Palmstierna, T., & Schlyter, F. (2005). Evaluation of the Drug Use Disorders Identification Test (DUDIT) in criminal justice and detoxification settings and in a Swedish population sample. *European Addiction Research*, 11(1), 22–31. <https://doi.org/10.1159/000081413>.
- Chase, H. W., & Clark, L. (2010). Gambling severity predicts midbrain response to near-miss outcomes. *Journal of Neuroscience*, 30(18), 6180–6187. <https://doi.org/10.1523/JNEUROSCI.5758-09.2010>.
- Ciccarelli, M., Cosenza, M., Nigro, G., & D'Olimpio, F. (2022). Does craving increase gambling severity? The role of attentional bias. *Journal of Affective Disorders*, 317, 403–408. <https://doi.org/10.1016/j.jad.2022.08.068>.
- Cloninger, C. R. (1999). *The Temperament and character inventory—revised*. Center for Psychobiology of Personality, Washington University.
- Cox, B. J., Enns, M. W., & Michaud, V. (2004). Comparisons between The South Oaks Gambling Screen and a DSM-IV—based interview in a community survey of problem gambling. *The Canadian Journal of Psychiatry*, 49(4), 258–264. <https://doi.org/10.1177/070674370404900406>.
- de Oliveira, M. P. M. T., da Silveira, D. X., de Carvalho, S. V. B., Collakis, S. T., Bizeto, J., & Silva, M. T. A. (2009). Reliability, validity and classification accuracy of The South Oaks Gambling Screen in a Brazilian sample. *Journal of Gambling Studies*, 25(4), 557–568. <https://doi.org/10.1007/s10899-009-9152-x>.
- de Oliveira, M. P. M. T., Silva, M. T. A., & da Silveira, D. X. da. (2002). Validity study of The South Oaks Gambling Screen (SOGS) among distinct groups of Brazilian gamblers. *Revista Brasileira de Psiquiatria*, 24(4), 170–176. <https://doi.org/10.1590/S1516-44462002000400005>.
- Derogatis, L. R. (1994). *Symptom checklist 90-R. Administration, Scoring and procedures manual-II for the revised version*. Clinical Psychometric Research.
- Derogatis, L. R. (2002). *SCL-90-R. Cuestionario de 90 Síntomas-Manual* (TEA (ed.)).
- Dickerson, M. G., Baron, E., Hong, S. M., & Cottrell, D. (1996). Estimating the extent and degree of gambling related problems in the Australian population: A national survey. *Journal of Gambling Studies*, 12(2), 161–178. <https://doi.org/10.1007/BF01539172>.
- Duvarci, I., & Varan, A. (2001). Reliability and validity study of the Turkish form of The South Oaks Gambling Screen. *Turk Psikiyatri Dergisi*, 12(1), 34–45.
- Duvarci, I., Varan, A., Coşkunol, H., & Ersoy, M. A. (1997). DSM-IV and The South Oaks Gambling Screen: Diagnosing and assessing pathological gambling in Turkey. *Journal of Gambling Studies*, 13(3), 193–206. <https://doi.org/10.1023/A:1024927115449>.
- Echeburúa, E., Báez, C., Fernández-Montalvo, J., & Páez, D. (1994). Cuestionario de juego patológico de South Oaks (SOGS): Validación Española. [The South Oaks Gambling Screen Questionnaire: The Spanish Validation]. *Análisis y Modificación de Conducta*, 20, 769–791.
- Esparza-Reig, J., Guillén-Riquelme, A., Martí-Vilar, M., & González-Sala, F. (2021). A reliability generalization meta-analysis of the South Oaks Gambling Screen (Sogs). *Psicothema*, 33(3), 490–499. <https://doi.org/10.7334/psicothema2020.449>.
- Finner, H., & Gontscharuk, V. (2009). Controlling the familywise error rate with plug-in estimator for the proportion of true null hypotheses. *Journal of the Royal Statistical Society. Series B: Statistical Methodology*, 71(5), 1031–1048. <https://doi.org/10.1111/j.1467-9868.2009.00719.x>.
- Gartner, C., Bickl, A., Härtl, S., Loy, J. K., & Häffner, L. (2022). Differences in problem and pathological gambling: A narrative review considering sex and gender. *Journal of Behavioral Addictions*, 11(2), 267–289. <https://doi.org/10.1556/2006.2022.00019>.
- Goodie, A. S., MacKillop, J., Miller, J. D., Fortune, E. E., Maples, J., Lance, C. E., & Campbell, W. K. (2013). Evaluating The South



- Oaks Gambling Screen with DSM-IV and DSM-5 criteria. *Assessment*, 20(5), 523–531. <https://doi.org/10.1177/1073191113500522>.
- Granero, R., Jiménez-Murcia, S., Fernández-Aranda, F., Del Pino-Gutiérrez, A., Mena-Moreno, T., Mestre-Bach, G., ... Menchón, J. M. (2020). Presence of problematic and disordered gambling in older age and validation of the South Oaks Gambling Scale. *Plos One*, 15(5), 1–22. <https://doi.org/10.1371/journal.pone.0233222>.
- Grant, J. E., Chamberlain, S. R., Schreiber, L. R. N., & Odlaug, B. L. (2012). Gender-related clinical and neurocognitive differences in individuals seeking treatment for pathological gambling. *Journal of Psychiatric Research*, 46(9), 1206–1211. <https://doi.org/10.1016/j.jpsychires.2012.05.013>.
- Grant, J. E., Odlaug, B. L., & Chamberlain, S. R. (2017). Gambling disorder, DSM-5 criteria and symptom severity. *Comprehensive Psychiatry*, 75, 1–5. <https://doi.org/10.1016/j.comppsy.2017.02.006>.
- Grant, J. E., Odlaug, B. L., & Mooney, M. E. (2012). Telescoping phenomenon in pathological gambling. *Journal of Nervous & Mental Disease*, 200(11), 996–998. <https://doi.org/10.1097/NMD.0b013e3182718a4d>.
- Gutiérrez-Zotes, J. A., Bayón, C., Montserrat, C., Valero, J., Labad, A., Cloninger, C. R., & Fernández-Aranda, F. (2004). Temperament and Character Inventory Revised (TCI-R). Standardization and normative data in a general population sample. *Actas Espanolas de Psiquiatria*, 32(1), 8–15. <http://www.ncbi.nlm.nih.gov/pubmed/14963776>.
- Gyollai, A., Urbán, R., Kun, B., Paksi, B., Arnold, P., Balázs, H., ... Demetrovics, Z. (2011). Problematic and pathologic gambling in Hungary: The Hungarian version and use of The South Oaks Gambling Screen (SOGS-HU). *Psychiatria Hungarica : A Magyar Pszichiatriai Tarsaság Tudományos Folyóirata*, 26(4), 230–240. <http://www.ncbi.nlm.nih.gov/pubmed/22058255>.
- Haddad, P., Roukoz, R., Akel, M., & Hallit, S. (2022). Gambling problems among Lebanese adults: Arabic-language version of The South Oaks Gambling Screen (SOGS) scale validation and correlates. *BMC Psychology*, 10(1), 1–10. <https://doi.org/10.1186/s40359-022-00727-6>.
- Hasin, D. S., O'Brien, C. P., Auriacombe, M., Borges, G., Bucholz, K., Budney, A., ... Grant, B. F. (2013). DSM-5 criteria for substance use disorders: Recommendations and rationale. *American Journal of Psychiatry*, 170(8), 834–851. <https://doi.org/10.1176/appi.ajp.2013.12060782>.
- Hayano, S., Dong, R., Miyata, Y., & Kasuga, S. (2021). The study of differences by region and type of gambling on the degree of gambling addiction in Japan. *Scientific Reports*, 11(1), 1–9. <https://doi.org/10.1038/s41598-021-92137-8>.
- Hildebrand, M. (2015). The psychometric properties of the drug use disorders identification test (DUDIT): A review of recent research. *Journal of Substance Abuse Treatment*, 53(January 2005), 52–59. <https://doi.org/10.1016/j.jsat.2015.01.008>.
- Hollingshead, A. B. (2011). Four factor index of social status. *Yale Journal of Sociology*, 8, 21–51.
- Jiménez-Murcia, S., Aymamí-Sanromà, N., Gómez-Pena, M., Álvarez-Moya, E., & Vallejo, J. (2006). *Protocols de tractament cognitivoconductual pel joc patològic i d'altres addiccions no tòxiques. [Protocols of cognitive-behaviour therapy for pathological gambling and other behavioural addictions]*. Hospital Universitari de Bellvitge, Departament de Salut, Generalitat de Catalunya.
- Jiménez-Murcia, S., Granero, R., Fernández-Aranda, F., Aymamí, N., Gómez-Peña, M., Mestre-Bach, G., ... Menchón, J. M. (2019). Developmental trajectories of gambling severity after cognitive-behavioral therapy. *European Psychiatry*, 60, 28–40. <https://doi.org/10.1016/j.eurpsy.2019.04.001>.
- Jiménez-Murcia, S., Granero, R., Fernández-Aranda, F., & Menchón, J. M. (2020). Comparison of gambling profiles based on strategic versus non-strategic preferences. *Current Opinion in Behavioral Sciences*, 31(November 2019), 13–20. <https://doi.org/10.1016/j.cobeha.2019.09.001>.
- Jiménez-Murcia, S., Stinchfield, R., Álvarez-Moya, E., Jaurrieta, N., Bueno, B., Granero, R., ... Vallejo, J. (2009). Reliability, validity, and classification accuracy of a Spanish translation of a measure of DSM-IV diagnostic criteria for pathological gambling. *Journal of Gambling Studies*, 25(1), 93–104. <https://doi.org/10.1007/s10899-008-9104-x>.
- Kido, M., & Shimazaki, T. (2007). Reliability and validity of the modified Japanese version of The South Oaks Gambling Screen (SOGS). *The Japanese Journal of Psychology*, 77(6), 547–552. <https://doi.org/10.4992/jjpsy.77.547>.
- Lesieur, H. R., & Blume, S. B. (1987). The South Oaks Gambling Screen (SOGS): A new instrument for the identification of pathological gamblers. *American Journal of Psychiatry*, 144(9), 1184–1188. <https://doi.org/10.1176/ajp.144.9.1184>.
- Levine, T. R., & Hullett, C. R. (2002). Eta squared, partial eta squared, and misreporting of effect size in communication research. *Human Communication Research*, 28(4), 612–625. <https://doi.org/10.1111/j.1468-2958.2002.tb00828.x>.
- Lucas, I., Granero, R., Fernández-Aranda, F., Solé-Morata, N., Demetrovics, Z., Baenas, I., ... Jiménez-Murcia, S. (2023). Gambling disorder duration and cognitive behavioural therapy outcome considering gambling preference and sex. *Journal of Psychiatric Research*, 158, 341–349. <https://doi.org/10.1016/j.jpsychires.2022.12.031>.
- Mathieu, S., Barrault, S., Brunault, P., & Varescon, I. (2020). The role of gambling type on gambling motives, cognitive distortions, and gambling severity in gamblers recruited online. *Plos One*, 15(10 October), 1–14. <https://doi.org/10.1371/journal.pone.0238978>.
- Mestre-Bach, G., Steward, T., Granero, R., Fernández-Aranda, F., del Pino-Gutiérrez, A., Mallorquí-Bagué, N., ... Jiménez-Murcia, S. (2019). The predictive capacity of DSM-5 symptom severity and impulsivity on response to cognitive-behavioral therapy for gambling disorder: A 2-year longitudinal study. *European Psychiatry*, 55, 67–73. <https://doi.org/10.1016/j.eurpsy.2018.09.002>.
- Muela, I., Ventura-Lucena, J. M., Navas, J. F., & Perales, J. C. (2023). The associative learning roots of affect-driven impulsivity and its role in problem gambling: A replication attempt and extension of Quintero et al. (2020). *Journal of Behavioral Addictions*, 12(1), 201–218. <https://doi.org/10.1556/2006.2023.00009>.
- Nigro, G., Matarazzo, O., Ciccarelli, M., Pizzini, B., Sacco, M., & Cosenza, M. (2022). Positive illusions: The role of cognitive distortions related to gambling and temporal perspective in



- chasing behavior. *Journal of Gambling Studies*, 38(3), 889–904. <https://doi.org/10.1007/s10899-021-10068-5>.
- Orford, J., Sproston, K., & Erens, B. (2003). SOGS and DSM-IV in the British gambling prevalence survey: Reliability and factor structure. *International Gambling Studies*, 3(1), 53–65. <https://doi.org/10.1111/j.1360-0443.1993.tb02093.x>.
- Otto, J. L., Smolenski, D. J., Garvey Wilson, A. L., Evatt, D. P., Campbell, M. S., Beech, E. H., ... Belsher, B. E. (2020). A systematic review evaluating screening instruments for gambling disorder finds lack of adequate evidence. *Journal of Clinical Epidemiology*, 120, 86–93. <https://doi.org/10.1016/j.jclinepi.2019.12.022>.
- Panayiotou, G., Artemi, T. F., Theodorou, M., Theodorou, C., & Neophytou, K. (2023). Not “just for fun”: Gambling, substance use, and the transdiagnostic role of emotion regulation. *Cogent Psychology*, 10(1). <https://doi.org/10.1080/23311908.2023.2183677>.
- Petry, N. M. (2003). Validity of a gambling scale for the addiction severity index. *The Journal of Nervous and Mental Disease*, 191(6), 399–407. <https://doi.org/10.1097/01.nmd.0000071589.20829.db>.
- Petry, N. M. (2005). Assessment. In *Pathological gambling: Etiology, comorbidity, and treatment* (Issue 21, pp. 35–54). American Psychological Association. <https://doi.org/10.1037/10894-003>.
- Petry, N. M., Armentano, C., Kuoch, T., Norinth, T., & Smith, L. (2003). Gambling participation and problems among South East Asian refugees to the United States. *Psychiatric Services*, 54(8), 1142–1148. <https://doi.org/10.1176/appi.ps.54.8.1142>.
- Reinert, D. F., & Allen, J. P. (2007). The alcohol use disorders identification test: An update of research findings. *Alcoholism: Clinical and Experimental Research*, 31(2), 185–199. <https://doi.org/10.1111/j.1530-0277.2006.00295.x>.
- Rogier, G., Picci, G., & Velotti, P. (2019). Struggling with happiness: A pathway leading depression to gambling disorder. *Journal of Gambling Studies*, 35(1), 293–305. <https://doi.org/10.1007/s10899-018-09825-w>.
- Salonen, A. H., Rosenström, T., Edgren, R., Volberg, R., Alho, H., & Castrén, S. (2017). Dimensions of The South Oaks Gambling Screen in Finland: A cross-sectional population study. *Scandinavian Journal of Psychology*, 58(3), 228–237. <https://doi.org/10.1111/sjop.12357>.
- Saunders, J. B., Aasland, O. G., Babor, T. F., de la Fuente, J. R., & Grant, M. (1993). Development of the alcohol use disorders identification test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption-II. *Addiction*, 88(6), 791–804. <https://doi.org/10.1111/j.1360-0443.1993.tb02093.x>.
- Slutske, W. S., Zhu, G., Meier, M. H., & Martin, N. G. (2011). Disordered gambling as defined by the diagnostic and statistical manual of mental disorders and The South Oaks Gambling Screen: Evidence for a common etiologic structure. *Journal of Abnormal Psychology*, 120(3), 743–751. <https://doi.org/10.1037/a0022879>.
- Stata-Corp (2023). *Stata statistical software: Release 18*. StataCorp LLC.
- Stinchfield, R. (2002). Reliability, validity, and classification accuracy of The South Oaks Gambling Screen (SOGS). *Addictive Behaviors*, 27(1), 1–19. [https://doi.org/10.1016/S0306-4603\(00\)00158-1](https://doi.org/10.1016/S0306-4603(00)00158-1).
- Stinchfield, R. (2003). Reliability, validity, and classification accuracy of a measure of DSM-IV diagnostic criteria for pathological gambling. *American Journal of Psychiatry*, 160(1), 180–182. <https://doi.org/10.1176/appi.ajp.160.1.180>.
- Tang, C. S. kum, Wu, A. M. S., Tang, J. Y. C., & Yan, E. C. W. (2010). Reliability, validity, and cut scores of the South Oaks Gambling Screen (SOGS) for Chinese. *Journal of Gambling Studies*, 26(1), 145–158. <https://doi.org/10.1007/s10899-009-9147-7>.
- Valenciano-Mendoza, E., Fernández-Aranda, F., Granero, R., Gómez-Peña, M., Moragas, L., Pino-Gutierrez, A. del, ... Jiménez-Murcia, S. (2021). Suicidal behavior in patients with gambling disorder and their response to psychological treatment: The roles of gender and gambling preference. *Journal of Psychiatric Research*, 143, 317–326. <https://doi.org/10.1016/j.jpsychires.2021.09.027>.
- Zakiniæiz, Y., Cosgrove, K. P., Mazure, C. M., & Potenza, M. N. (2017). Does telescoping exist in male and female gamblers? Does it matter? *Frontiers in Psychology*, 8(SEP), 1–5. <https://doi.org/10.3389/fpsyg.2017.01510>.
- Zakiniæiz, Y., & Potenza, M. N. (2018). Gender-related differences in addiction: A review of human studies. *Current Opinion in Behavioral Sciences*, 23, 171–175. <https://doi.org/10.1016/j.cobeha.2018.08.004>.
- Zhou, X., Obuchowski, N. A., & McClish, D. K. (2002). *Statistical Methods in diagnostic medicine*. Wiley. <https://doi.org/10.1002/9780470317082>.

