

Almost sure growth of integrated supOU processes

Danijel Grahovac^{*1} and Péter Kevei^{†2}

¹*School of Applied Mathematics and Informatics, J. J. Strossmayer University of Osijek, Trg Ljudevita Gaja 6, 31000 Osijek, Croatia*

²*Bolyai Institute, University of Szeged, Aradi vértanúk tere 1, 6720 Szeged, Hungary*

Abstract

Superpositions of Ornstein-Uhlenbeck processes allow a flexible dependence structure, including long range dependence for OU-type processes. Their complex asymptotics are governed by three effects: the behavior of the Lévy measure both at infinity and at zero, and the behavior at zero of the measure governing the dependence. We establish almost sure rates of growth depending on the characteristics of the process and prove a Marcinkiewicz–Zygmund type SLLN for the integrated process.

AMS 2010 Subject Classifications: 60G17, 60F15, 60G55, 60G57

Keywords: supOU processes, infinitely divisible random measure, Marcinkiewicz–Zygmund type strong law of large numbers, almost sure properties, rate of growth

1 Introduction

A superposition of Ornstein-Uhlenbeck type processes (*supOU process*) [1] is a strictly stationary process $(X(t))_{t \in \mathbb{R}}$ given by

$$X(t) = \iint_{(0, \infty) \times (-\infty, t]} e^{-x(t-s)} \Lambda(dx, ds), \quad (1)$$

where Λ is a homogeneous infinitely divisible independently scattered random measure on $(0, \infty) \times \mathbb{R}$, meaning that for every sequence $A_1, A_2, \dots$

^{*}dgrahova@mathos.hr

[†]kevei@math.u-szeged.hu

of disjoint bounded sets from $\mathcal{B}((0, \infty) \times \mathbb{R})$, the random variables $\Lambda(A_1)$, $\Lambda(A_2)$, $\dots$ are independent. Moreover, for bounded $A \in \mathcal{B}((0, \infty) \times \mathbb{R})$ we have

$$\begin{aligned} & \log \mathbf{E} e^{i\theta \Lambda(A)} \\ &= (\pi \times \text{Leb})(A) \left(i\theta a - \frac{\theta^2}{2} b + \int_{\mathbb{R}} \left(e^{i\theta z} - 1 - i\theta z \mathbf{1}(|z| \leq 1) \right) \lambda(dz) \right), \end{aligned} \quad (2)$$

where π is a measure on $(0, \infty)$ such that $m_{-1}(\pi) := \int_{(0, \infty)} x^{-1} \pi(dx) < \infty$, and (a, b, λ) is a Lévy-Khintchine triplet of some infinitely divisible distribution such that $\int_{|z| \geq 1} \log |z| \lambda(dz) < \infty$. The *generating quadruple* (a, b, λ, π) completely determines the distribution of the supOU process.

The one-dimensional marginal distribution of the supOU process corresponds to that of Ornstein–Uhlenbeck type process driven by the Lévy process L with characteristic triplet (a, b, λ) . In particular,

$$X(t) \stackrel{d}{=} m_{-1}(\pi) \int_0^\infty e^{-s} L(ds),$$

hence the marginal distribution is controlled by the triplet (a, b, λ) . Lévy driven Ornstein–Uhlenbeck processes always have exponentially decaying correlations, whenever their second moment exists. However, supOU processes have more complex dependence structure since the correlation function of X is the Laplace transform of $x^{-1} \pi(dx)$, provided that $\mathbf{E} X(t)^2 < \infty$. See [1, 3, 4] for more details.

In this paper we investigate the almost sure behavior of the *integrated supOU process*

$$X^*(t) = \int_0^t X(u) du. \quad (3)$$

In our main results, Theorems 1 and 2 below, we prove a Marcinkiewicz–Zygmund type strong law of large numbers for the integrated process. The classical Marcinkiewicz–Zygmund theorem, see e.g. [25, Theorem 6.7.1], states that if iid random variables $Y_1, Y_2, \dots$ have finite moment of order $\gamma \in (1, 2)$, then the centered partial sum $S_n = \sum_{i=1}^n (Y_i - \mathbf{E} Y_i)$ is $o(n^{1/\gamma})$ almost surely. For the same to hold for X^* , the main condition is that

$$\iint_{(0, \infty) \times \mathbb{R}} \frac{|z|^\gamma}{x^\gamma} \mathbf{1}(|z| > x) \pi(dx) \lambda(dz) < \infty. \quad (4)$$

This implies that $\mathbf{E} |X^*(1)|^\gamma < \infty$, however, (4) is stronger than the moment assumption alone, as it is also affected by the behavior of π and λ near 0.

On the other hand, in the purely Gaussian case, a law of iterated logarithm holds (Theorem 3).

The motivation for studying the almost sure rate of growth comes from limiting properties established recently in a series of papers [18, 20, 22]. The integrated supOU processes satisfy distributional limit theorems of the form

$$\left\{ \frac{X^*(Tt) - \mathbf{E}X^*(Tt)}{T^H \widehat{\ell}(T)} \right\} \xrightarrow{\text{fdd}} \{Z(t)\}, \quad \text{as } T \rightarrow \infty, \quad (5)$$

with some slowly varying function $\widehat{\ell}$, $H \in (0, 1)$ and limiting process Z depending on the generating quadruple (a, b, λ, π) . Despite satisfying limit theorems, integrated supOU processes exhibit *moment intermittency* (see e.g. [8, 9, 16, 28, 31, 49]). Namely, their higher order moments may grow faster than those of the limiting process, i.e.

$$\lim_{t \rightarrow \infty} \frac{\log \mathbf{E}|X^*(t) - \mathbf{E}X^*(t)|^q}{\log t} = \begin{cases} Hq, & 0 \leq q \leq \frac{\alpha}{1-H}, \\ q - \alpha, & q \geq \frac{\alpha}{1-H}, \end{cases} \quad (6)$$

where H is the self-similarity parameter of the limiting process Z in (5) and α is parameter related to behavior of π near zero. Intermittency implies that for any $\varepsilon > 0$ and $\delta > 0$ we eventually have [23]

$$\mathbf{P}(|X^*(t) - \mathbf{E}X^*(t)| > ct^{1-\varepsilon}) \geq t^{-\alpha-\delta}. \quad (7)$$

Therefore, the probability that the normalized integrated supOU process exhibits increasingly large values decreases as a power function of time. This small fraction of probability space where $X^*(t)$ is unduly large provides the main contribution to higher order moments resulting in moment intermittency (6). In particular, classical large deviation results with exponentially decaying probabilities do not typically hold in limit theorems for supOU processes [23].

These limiting results suggest that X^* has a similar behavior of moments as the sequence of independent random variables $(Y(n))_{n \in \mathbb{N}}$ given by

$$Y(n) = \begin{cases} n^H, & \text{with probability } 1 - n^{-\alpha}, \\ n, & \text{with probability } n^{-\alpha}. \end{cases}$$

Moreover, for $\alpha < 1$ probabilities $\mathbf{P}(Y(n) \geq n)$ (as those in (7)) are not summable. By the second Borel–Cantelli lemma $Y(n) \geq n$ happens infinitely often.

The main motivation behind this paper is to investigate whether the same is true for paths of integrated supOU processes. Namely, we wonder whether the normalized integrated supOU process exhibits *physical intermittency*, meaning that at large times it exhibits increasingly tall peaks (see e.g. [8, 10, 29, 30, 49]). We show that this not so. In particular, for any $\varepsilon > 0$, we have that

$$\lim_{t \rightarrow \infty} \frac{X^*(t) - \mathbf{E}X^*(t)}{t^{H+\varepsilon}} = 0 \quad \text{a.s.}$$

We conclude that the paths of normalized integrated supOU process do not exhibit increasingly large values infinitely often, even though (6) and (7) would suggest differently at first. This can be compared with the recently observed phenomenon of *dissipation* in SPDEs [29, 30].

Our results provide a rare example of a law of the iterated logarithm type and limsup type results for processes with strong dependence. Such results are typically known only for Gaussian [38, 48] or self-similar processes, like, for example, the law of iterated logarithm for fractional Brownian motion, limsup results for α -stable Lévy processes [44], linear fractional stable process [45], or for general self-similar processes (see [32, 45]). Beyond the class of self-similar processes, let us mention that for Lévy processes the rate of growth depends on the regular variation index of the Lévy measure at infinity (see [44, Section 48] and the references therein).

One may view supOU processes as aggregation of Lévy driven Ornstein-Uhlenbeck processes with random mean reverting parameter. Such models have been extensively studied ever since the seminal paper [24] when it was realized that by aggregating simpler processes one may obtain complex dependence properties like long range dependence (see e.g. [34, 37, 41] and the references therein). However, law of iterated logarithm type results for such processes have been unknown so far. Moreover, distributional limit theorems for supOU processes are similar to that of trawl processes [3, 19, 39, 46]. Hence, our results may pave the way for establishing similar pathwise results for these classes of processes.

From a modeling perspective, supOU processes offer the appealing feature of allowing the marginal distribution and the dependence structure to be modeled independently. They have been used in various applications, mainly in finance where they provide a model for stochastic volatility (see e.g. [5, 11, 15]). In this context, the integrated process (3) is the integrated volatility. SupOU processes have also been applied in other areas, like e.g. in astrophysics [26].

The paper is organized as follows. Section 2 contains the main results.

In Section 3 we give some preliminary facts needed for the proofs and also discuss moments and tail behavior of the integrated process. The proofs of the main results are given in Sections 4 and 5. Section 6 contains the proof of the Gaussian case. In Appendix A we prove a stochastic Fubini theorem and in Appendix B we provide an exotic example of measures π and λ .

Throughout the paper c will denote a constant that may change from row to row. For any measure Q on $(0, \infty)$ we denote

$$m_p(Q) = \int_{(0, \infty)} x^p Q(dx),$$

and for its tail we write

$$\overline{Q}(r) = Q((r, \infty)), \quad r > 0.$$

2 The almost sure growth

The random measure Λ in (1) has a modification with the Lévy–Itô decomposition (see [40, Theorem 4.5] and [4, Theorem 2.2]). We will work with this modification so that

$$\begin{aligned} \Lambda(dx, ds) = & a\pi(dx)ds + \Lambda^G(dx, ds) + \int_{|z| \leq 1} z(\mu - \nu)(dx, ds, dz) \\ & + \int_{|z| > 1} z\mu(dx, ds, dz), \end{aligned} \tag{8}$$

where Λ^G is Gaussian random measure defined as in (2) with generating quadruple $(0, b, 0, \pi)$ and μ is a Poisson random measure on $(0, \infty) \times \mathbb{R} \times \mathbb{R}$ with intensity measure $\nu(dx, ds, dz) = \pi(dx)ds\lambda(dz)$. Furthermore, the existence of a measurable modification of the supOU process $(X(t))_{t \in \mathbb{R}}$ in (1) follows from [2, p. 926]. In the following, we will work with such a modification. However, we note that càdlàg modification does not necessarily exist, see [17, Proposition 1].

Remark 1. The supOU processes were introduced in [1] with a slightly different parametrization ($-xt + s$ in the exponent in (1)). Representation (1) is a moving average representation, which was first used in [13] (see also [14]). The conditions $\int_{(0, \infty)} x^{-1}\pi(dx) < \infty$ and $\int_{|z| > 1} \log |z|\lambda(dz) < \infty$ are necessary and sufficient for the existence of X by Theorem 3.1 in [1]. See also [14, Proposition 2.1] and [4] with the unnecessary assumption that π is a probability measure.

The integrated process (3) may be written in the form

$$X^*(t) = \int_0^t X(u)du = \iint_{(0,\infty) \times \mathbb{R}} f_t(x, s) \Lambda(dx, ds),$$

where

$$f_t(x, s) = \begin{cases} x^{-1}(1 - e^{-xt})e^{xs}, & s \leq 0, \\ x^{-1}(1 - e^{-x(t-s)})\mathbf{1}(s \leq t), & s > 0. \end{cases}$$

The use of the stochastic Fubini theorem is justified in [2, Theorem 3.1] and [22, Lemma 4.1] (see the Appendix for more details). By (8), we have

$$\begin{aligned} X^*(t) &= a \int_{(0,\infty)} \int_{\mathbb{R}} f_t(x, s) \pi(dx) ds + \int_{(0,\infty)} \int_{\mathbb{R}} f_t(x, s) \Lambda^G(dx, ds) \\ &\quad + \int_{(0,\infty)} \int_{\mathbb{R}} \int_{|z| \leq 1} z f_t(x, s) (\mu - \nu)(dx, ds, dz) \\ &\quad + \int_{(0,\infty)} \int_{\mathbb{R}} \int_{|z| > 1} z f_t(x, s) \mu(dx, ds, dz). \end{aligned}$$

Since $\int_{(0,\infty)} \int_{\mathbb{R}} f_t(x, s) \pi(dx) ds = tm_{-1}(\pi)$, the behavior of the deterministic part is clear. Hence, in the following we assume that $a = 0$ whenever $\int_{|z| \leq 1} |z| \lambda(dz) = \infty$, and when $\int_{|z| \leq 1} |z| \lambda(dz) < \infty$, then we assume $a = \int_{|z| \leq 1} z \lambda(dz)$, so that there is no centering. In particular, if $\int_{|z| > 1} |z| \lambda(dz) < \infty$, then

$$\mathbf{E}X^*(t) = \begin{cases} tm_{-1}(\pi) \int_{\mathbb{R}} z \lambda(dz), & \text{if } \int_{|z| \leq 1} |z| \lambda(dz) < \infty, \\ tm_{-1}(\pi) \int_{|z| > 1} z \lambda(dz), & \text{if } \int_{|z| \leq 1} |z| \lambda(dz) = \infty. \end{cases} \quad (9)$$

Moreover, we assume for the moment that there is no Gaussian component in the generating quadruple.

The technique of the proof depends on whether the Poisson random measure needs compensation or not. Hence, we split the results in these two cases.

Theorem 1. *Assume that $a = \int_{|z| \leq 1} |z| \lambda(dz) < \infty$ and $b = 0$.*

1. *If for some $\gamma \in (0, 2)$*

$$\iint_{(0,\infty) \times \mathbb{R}} \frac{|z|^\gamma}{x^\gamma} \mathbf{1}(|z| > x) \pi(dx) \lambda(dz) < \infty, \quad (10)$$

and if $\gamma = 1$ additionally $m_{-1-\delta}(\pi) < \infty$ for some $\delta > 0$, then

$$\lim_{t \rightarrow \infty} \frac{X^*(t) - \mathbf{1}(\gamma \geq 1) \mathbf{E}X^*(t)}{t^{1/\gamma}} = 0 \quad a.s.$$

2. If (10) holds for $\gamma = 2$, then

$$\limsup_{t \rightarrow \infty} \frac{|X^*(t) - \mathbf{E}X^*(t)|}{\sqrt{2t \log \log t}} = \sqrt{\mathbf{Var}X^*(1)} \quad a.s.$$

From Corollary 2 below it follows that (10) implies $\mathbf{E}|X^*(1)|^\gamma < \infty$, which is the moment assumption in the classical Marcinkiewicz–Zygmund theorem (see e.g. [25, Theorem 6.7.1]).

For the $\int_{|z| \leq 1} |z| \lambda(dz) = \infty$ case, we introduce the following indices

$$\begin{aligned} \beta_0 &= \inf \left\{ \beta \geq 0 : \int_{|z| \leq 1} |z|^\beta \lambda(dz) < \infty \right\}, \\ \eta_\infty &= \sup \left\{ \eta \geq 0 : \int_{|z| > 1} |z|^\eta \lambda(dz) < \infty \right\}, \\ \alpha_0 &= \sup \left\{ \alpha \geq 0 : \int_{(0,1]} x^{-\alpha-1} \pi(dx) < \infty \right\}. \end{aligned} \quad (11)$$

The constant β_0 is the Blumenthal–Gettoor index of the Lévy measure λ and η_∞ is the tail index of the Lévy measure λ , which also corresponds to the tail index of the marginal distribution [14, Proposition 3.1]. Assumption $\int_{|z| \leq 1} |z| \lambda(dz) = \infty$ implies that $\beta_0 \geq 1$, while $\beta_0 \leq 2$, as λ is a Lévy measure.

In what follows, we put $\beta = \beta_0$ if $\int_{|z| \leq 1} |z|^{\beta_0} \lambda(dz) < \infty$, otherwise $\beta > \beta_0$ arbitrarily close. Similarly, $\eta \leq \eta_\infty$, $\alpha \leq \alpha_0$ with equality if the corresponding quantity is finite.

Theorem 2. Assume that $\int_{|z| \leq 1} |z| \lambda(dz) = \infty$, $a = 0$ and $b = 0$.

1. If $\beta \leq 1 + \alpha$, assume that (10) holds for some $\gamma \in (0, 2]$ and if $\gamma = 1$ additionally $m_{-1-\delta}(\pi) < \infty$ for some $\delta > 0$. Then

$$\limsup_{t \rightarrow \infty} \frac{|X^*(t) - \mathbf{1}(\gamma \geq 1) \mathbf{E}X^*(t)|}{t^{1/\gamma} \log t} \leq 1 \quad a.s.$$

2. If $\beta \geq 1 + \alpha$, assume that

$$\iint_{(0,\infty) \times \mathbb{R}} \frac{|z|^\gamma}{x^\gamma} \mathbf{1}(|z| > x, |z| > 1) \pi(dx) \lambda(dz) < \infty \quad (12)$$

holds with $\gamma < 1/(1 - \alpha/\beta)$. Then

$$\lim_{t \rightarrow \infty} \frac{X^*(t) - \mathbf{1}(\gamma \geq 1) \mathbf{E}X^*(t)}{t^{1/\gamma}} = 0 \quad a.s.$$

To understand the assumptions, it is worth discussing how (10) is related to indices α , β and η . If (10) holds, then

$$\int_{(0,1]} x^{-\gamma} \int_{|z|>1} |z|^\gamma \lambda(dz) \pi(dx) < \infty,$$

which implies that $\gamma \leq 1 + \alpha$ and $\gamma \leq \eta$. Furthermore, (10) with any $\gamma > 0$ implies that (10) holds with $\gamma = 0$, that is

$$\int_{(0,\infty)} \lambda(\{z : |z| > x\}) \pi(dx) = \int_{\mathbb{R}} \pi((0, |z|)) \lambda(dz) < \infty. \quad (13)$$

Therefore, if $\lambda(\{z : |z| > x\}) = h(x)x^{-\beta_0}$ or $\pi((0, z)) = h(z)z^{1+\alpha_0}$ around 0, where h is a function such that $\lim_{x \downarrow 0} h(x)x^\varepsilon = 0$ and $\lim_{x \downarrow 0} h(x)x^{-\varepsilon} = \infty$ for any $\varepsilon > 0$, then (13) implies $\beta \leq 1 + \alpha$. In particular, this holds if h is slowly varying at 0, i.e. $\lim_{x \downarrow 0} \frac{h(xy)}{h(x)} = 1$ for each $y > 0$ (see e.g. [7]). On the other hand, it is possible to construct rather exotic measures π and λ such that (10) holds but $\beta > 1 + \alpha$ (see the example in the Appendix). Further connections between (10) and indices α , β and η can be seen from the proof of Corollary 1 below.

The following corollary summarizes the main results in terms of the bounds for the *rate of growth* of the (centered) integrated process. For simplicity we assume that π is a finite measure. The generalization to infinite measures resulting in a few more cases, is straightforward.

Corollary 1. *Assume that π is a finite measure. For the rate of growth of the integrated process*

$$R(t) = \frac{\log |X^*(t) - \mathbf{1}(\mathbf{E}X^*(t) < \infty)\mathbf{E}X^*(t)|}{\log t},$$

it holds a.s. that

$$\limsup_{t \rightarrow \infty} R(t) \leq \begin{cases} \frac{1}{2}, & \text{if } \alpha \geq 1 \text{ and } \eta \geq 2, \\ \frac{1}{\eta}, & \text{if } \alpha \geq 1, \eta < 2, \text{ or } \alpha < 1, \eta \leq 1 + \alpha, \beta \leq 1 + \alpha \\ \frac{1}{1+\alpha}, & \text{if } \alpha < 1, \eta > 1 + \alpha \text{ and } \beta \leq 1 + \alpha, \\ 1 - \frac{\alpha}{\beta}, & \text{if } \alpha < 1, \eta > 1 + \alpha, \text{ and } \beta > 1 + \alpha. \end{cases}$$

Proof. Write

$$\begin{aligned}
& \iint_{(0,\infty) \times \mathbb{R}} \frac{|z|^\gamma}{x^\gamma} \mathbf{1}(|z| > x) \pi(dx) \lambda(dz) \\
&= \int_{(0,1]} x^{-\gamma} \pi(dx) \left(\int_{|z| \in (x,1]} |z|^\gamma \lambda(dz) + \int_{|z| > 1} |z|^\gamma \lambda(dz) \right) \\
&\quad + \int_{(1,\infty)} x^{-\gamma} \pi(dx) \int_{|z| > x} |z|^\gamma \lambda(dz) \\
&=: J_1 + J_2 + J_3.
\end{aligned}$$

Then $J_1 < \infty$ if $\gamma \leq 1 + \alpha$ and $\gamma \geq \beta$, $J_2 < \infty$ if $\gamma \leq \eta \wedge (1 + \alpha)$, and $J_3 < \infty$ for $\gamma \leq \eta$ as π is assumed to be a finite measure. We now consider each case separately.

($\alpha \geq 1$ and $\eta \geq 2$): Then $\beta \leq 1 + \alpha$ necessarily holds, and (10) holds with any $\gamma < 2$. Thus the claim follows from Theorem 1 or 2 part 1.

($\alpha \geq 1$, $\eta < 2$), or ($\alpha < 1$, $\eta \leq 1 + \alpha$, $\beta \leq 1 + \alpha$): Then $J_2 < \infty$ for any $\gamma \leq \eta$, and $J_1 < \infty$ for $\gamma \leq 1 + \alpha$ since $1 + \alpha \geq \beta$. Finally, $J_3 < \infty$ for any $\gamma \leq \eta$. Thus (10) holds with $\gamma \leq \eta$, and the claim follows again as above.

($\alpha < 1$, $\eta \geq 1 + \alpha$, $\beta \leq 1 + \alpha$): Then $J_1 < \infty$ since $\gamma \leq 1 + \alpha$ and $\beta \leq 1 + \alpha$, while $J_2 < \infty$ and $J_3 < \infty$ since $\gamma \leq 1 + \alpha \leq \eta$. Thus (10) holds with $\gamma \leq 1 + \alpha$, and the claim follows.

($\alpha < 1$, $\eta \geq 1 + \alpha$, $\beta \geq 1 + \alpha$): Now we use Theorem 2 part 2, and we have to check (12). In this case $J_1 = 0$, $J_2 < \infty$ for $\gamma \leq (1 - \frac{\alpha}{\beta})^{-1} \leq 1 + \alpha \leq \eta$, while $J_3 < \infty$ for $\gamma \leq \eta$. $\square$

To show that the bounds obtained are optimal, we review details of the limit theorems in (5). Suppose that π has a density p which is regularly varying at zero, that is

$$p(x) \sim \alpha_0 \ell(x^{-1}) x^{\alpha_0}, \quad \text{as } x \rightarrow 0,$$

for some $\alpha_0 > 0$ and ℓ slowly varying at infinity. If the integrated process has finite variance, then we have (5) if one of the following holds:

1. $\alpha_0 > 1$, in which case $H = 1/2$ and Z is Brownian motion,
2. $b = 0$, $\alpha_0 \in (0, 1)$ and $\int_{|z| \leq 1} |z|^{1+\alpha_0} \lambda(dz) < \infty$, in which case $H = 1/(1 + \alpha_0)$ and Z is $(1 + \alpha_0)$ -stable Lévy process,
3. $b = 0$, $\alpha_0 \in (0, 1)$ and $\lambda(\{x \in \mathbb{R} : |x| > z\}) \sim cz^{-\beta_0}$ as $z \rightarrow 0$ with $1 + \alpha_0 < \beta_0 < 2$, in which case $H = 1 - \alpha_0/\beta_0$ and Z is β_0 -stable process with dependent increments,

4. $b > 0$ and $\alpha_0 \in (0, 1)$, in which case $H = 1 - \alpha_0/2$ and Z is fractional Brownian motion.

For the infinite variance case, the type of the limit depends additionally on the regular variation index of the marginal distribution [21].

By comparing with the bounds from Theorem 1 and 2, one can see that the bounds are optimal, up to a slowly varying terms. We can conclude that the almost sure growth is of the same order as in the weak limits even though from (7) one would expect differently.

Finally, for completeness we consider the purely Gaussian case, that is when $a = 0$, $b > 0$, $\lambda \equiv 0$. In this case, the following law of iterated logarithm follows from general results on Gaussian processes combined with the asymptotic analysis of variance. The proof is given in Section 6.

Theorem 3. *Assume that $a = 0$, $b > 0$ and $\mu \equiv 0$,*

1. *If $m_{-2}(\pi) < \infty$, then*

$$\limsup_{t \rightarrow \infty} \frac{|X^*(t)|}{\sqrt{2bm_{-2}(\pi)t \log \log t}} = 1 \quad a.s.$$

2. *If π has a density p such that $p(x) \sim \alpha \ell(x^{-1})x^\alpha$ for some $\alpha \in (0, 1)$ and ℓ slowly varying at infinity, then*

$$\limsup_{t \rightarrow \infty} \frac{|X^*(t)|}{\tilde{\sigma} \ell(t)^{\frac{1}{2}} t^{1-\frac{\alpha}{2}} \sqrt{2 \log \log t}} = 1 \quad a.s.$$

$$\text{where } \tilde{\sigma}^2 = b \frac{\Gamma(1+\alpha)}{(2-\alpha)(1-\alpha)}.$$

3 Preliminaries

As the positive and negative jumps of the Poisson random measure μ from (8) are independent, we will show in the proofs of Theorems 1 and 2 that it is enough to consider supOU processes for which this random measure has only positive jumps, i.e. $\lambda((-\infty, 0)) = 0$. Since we consider the Gaussian case separately, we assume that there is no Gaussian component.

The following statement is Theorem 4.1 in [1]. We prove it again here because of the explicit form of the Lévy measure η_t^* .

Lemma 1 ([1, Theorem 4.1]). *The random variable $X^*(t)$ is infinitely divisible with characteristic function*

$$\mathbf{E}e^{i\theta X^*(t)} = \exp \left\{ iA\theta + \int_{(0,\infty)} (e^{i\theta y} - 1 - i\theta y \mathbf{1}(y \leq 1)) \eta_t^*(dy) \right\},$$

where $A \in \mathbb{R}$ is an explicit constant and

$$\eta_t^*(B) = \nu(\{(x, s, z) : f_t(x, s)z \in B\}).$$

Furthermore,

$$\bar{\eta}_t^*(r) = \eta_t^*((r, \infty)) = \int_{(0,\infty)} x^{-1} \pi(dx) \int_{\frac{xr}{1-e^{-xt}}}^{\infty} \bar{\lambda}(z) \frac{1}{z-xr} dz.$$

Proof. The first part follows from [42, Theorem 2.7]. Note that for $r > 0$

$$\begin{aligned} & \{(x, s, z) : f_t(x, s)z > r\} \\ &= \left\{ (x, s, z) : z > \frac{xr}{1-e^{-xt}}, x^{-1} \log \frac{xr}{z(1-e^{-xt})} < s \leq 0 \right\} \\ & \cup \left\{ (x, s, z) : z > \frac{xr}{1-e^{-xt}}, 0 < s < t + x^{-1} \log \left(1 - \frac{xr}{z} \right) \right\}, \end{aligned}$$

thus

$$\begin{aligned} & \bar{\eta}_t^*(r) \\ &= \int_{(0,\infty)} \pi(dx) \int_{(\frac{xr}{1-e^{-xt}}, \infty)} \lambda(dz) \frac{1}{x} \left(\log \frac{z(1-e^{-xt})}{xr} + tx + \log \left(1 - \frac{xr}{z} \right) \right) \\ &= \int_{(0,\infty)} x^{-1} \pi(dx) \int_{(\frac{xr}{1-e^{-xt}}, \infty)} \lambda(dz) \log \left((e^{xt} - 1) \left(\frac{z}{xr} - 1 \right) \right) \\ &= \int_{(0,\infty)} x^{-1} \pi(dx) \int_{\frac{xr}{1-e^{-xt}}}^{\infty} \bar{\lambda}(z) \frac{1}{z-xr} dz, \end{aligned}$$

where the last equality follows after integration by parts combined with the fact that

$$\lim_{u \rightarrow \infty} \bar{\lambda}(u) \log(e^{xt} - 1) \left(\frac{u}{xr} - 1 \right) = 0.$$

The latter limit holds since $\lim_{u \rightarrow \infty} \bar{\lambda}(u) \log u = 0$ by the integrability assumption $\int_{(1,\infty)} \log z \lambda(dz) < \infty$. $\square$

Note that if $m_0(\pi) < \infty$, then $m_{-\beta}(\pi) < \infty$ for all $\beta \in [0, 1]$.

Corollary 2. *Let $t > 0$ and $\beta > 0$. If $\mathbf{E}X^*(t)^\beta < \infty$, then $\int_{(1,\infty)} z^\beta \lambda(dz) < \infty$. On the other hand, if $\int_{(1,\infty)} z^\beta \lambda(dz) < \infty$, then $\mathbf{E}X^*(t)^\beta < \infty$ provided one of the following holds*

1. $\beta \geq 1$,
2. $\beta \in (0, 1)$ and $m_{-\beta}(\pi) < \infty$,
3. $\beta \in (0, 1)$, $m_{-\beta}(\pi) = \infty$ and

$$\int_{(1,\infty)} x^{-\beta} \pi(dx) \int_x^\infty \bar{\lambda}(z) z^{\beta-1} dz < \infty.$$

Proof. By [44, Theorem 25.3], $\mathbf{E}X^*(t)^\beta < \infty$ if and only if $\int_1^\infty r^{\beta-1} \bar{\eta}_t^*(r) dr < \infty$. By Lemma 1, and changing variables $u = \frac{xr}{z}$

$$\begin{aligned} & \int_1^\infty r^{\beta-1} \bar{\eta}_t^*(r) dr \\ &= \int_1^\infty dr \int_{(0,\infty)} \pi(dx) \int_0^\infty dz \frac{r^{\beta-1} \bar{\lambda}(z)}{x(z-xr)} \mathbb{1}\left(z > \frac{xr}{1-e^{-xt}}\right) \\ &= \int_{(0,\infty)} x^{-1} \pi(dx) \int_{x/(1-e^{-xt})}^\infty dz \bar{\lambda}(z) z^{\beta-1} x^{-\beta} \int_{x/z}^{1-e^{-xt}} \frac{u^{\beta-1}}{1-u} du. \end{aligned} \tag{14}$$

The integral with respect to u is an increasing function in z , therefore in each case $\mathbf{E}X^*(t)^\beta < \infty$ implies $\int_1^\infty z^{\beta-1} \bar{\lambda}(z) dz < \infty$, which further implies $\int_{(1,\infty)} z^\beta \lambda(dz) < \infty$.

For the converse, note that

$$\sup_{0 \leq x \leq 1} x^{-\beta} \int_0^{1-e^{-xt}} u^{\beta-1} (1-u)^{-1} du < \infty,$$

while for $x > 1$

$$\int_0^{1-e^{-xt}} u^{\beta-1} (1-u)^{-1} du = t \int_0^x (1-e^{-ty})^{\beta-1} dy \leq cx.$$

Indeed, for $\beta \geq 1$ this follows from $1 - e^{-ty} \leq 1$. For $\beta < 1$ we have

$$t \int_0^x (1 - e^{-ty})^{\beta-1} dy \leq t \int_0^1 (1 - e^{-ty})^{\beta-1} dy + t \int_1^x (1 - e^{-t})^{\beta-1} dy \leq cx.$$

Substituting back into (14) and using also that $u \geq 1 - e^{-u}$, we obtain

$$\begin{aligned} \int_1^\infty \bar{\eta}_t^*(r) r^{\beta-1} dr &\leq c \int_{(0,1]} x^{-1} \pi(dx) \int_{1/t}^\infty \bar{\lambda}(z) z^{\beta-1} dz \\ &\quad + t \int_{(1,\infty)} x^{-\beta} \pi(dx) \int_x^\infty \bar{\lambda}(z) z^{\beta-1} dz, \end{aligned}$$

and the finiteness of the integral follows. $\square$

Thus, in most cases the behavior of π does not have effect on the existence of the moments. However, it does have effect on the existence of exponential moments.

Lemma 2. *Let $\varepsilon_0 = \sup\{s \geq 0 : \pi((0, s)) = 0\}$, $K_0(t) = \frac{1-e^{-\varepsilon_0 t}}{\varepsilon_0}$, with $K_0(t) = t$ for $\varepsilon_0 = 0$. If $\mathbf{E}e^{sX^*(t)} < \infty$, for $s > 0$ then $\int_{(1,\infty)} e^{zy} \lambda(dz) < \infty$ for any $y < K_0(t)s$. On the other hand, if $\int_{(1,\infty)} e^{zK_0(t)s} \lambda(dz) < \infty$, then $\mathbf{E}e^{sX^*(t)} < \infty$.*

Proof. From [44, Theorem 25.3] we see that $\mathbf{E}e^{sX^*(t)} < \infty$ if and only if $\int_{(1,\infty)} e^{sr} \bar{\eta}_t^*(dr) < \infty$, which holds if and only if $\int_1^\infty s e^{sr} \bar{\eta}_t^*(r) dr < \infty$. Using Lemma 1 and changing variables $u = \frac{xr}{z}$ we have

$$\begin{aligned} \int_1^\infty e^{sr} \bar{\eta}_t^*(r) dr &= \int_1^\infty dr \int_{(0,\infty)} \pi(dx) \int_{\frac{xr}{1-e^{-xt}}}^\infty e^{sr} x^{-1} \bar{\lambda}(z) (z-xr)^{-1} dz \\ &= \int_{(0,\infty)} x^{-2} \pi(dx) \int_0^\infty dz \int_{x/z}^{1-e^{-xt}} e^{szu/x} \bar{\lambda}(z) (1-u)^{-1} du \mathbf{1}\left(\frac{x}{z} < 1 - e^{-xt}\right) \\ &= \int_{(0,\infty)} x^{-2} \pi(dx) \int_0^1 (1-u)^{-1} du \int_0^\infty \bar{\lambda}(z) e^{zsu/x} dz \mathbf{1}\left(\frac{x}{z} < u < 1 - e^{-xt}\right). \end{aligned}$$

From the last formula we see that if the integral on the left-hand side is finite, then $\int_1^\infty \bar{\lambda}(z) e^{zy} dz < \infty$ for $y < s(1 - e^{-xt})/x$, where $x \in \text{supp}(\pi)$. Since $(1 - e^{-xt})/x$ is decreasing in x , the first result follows.

Assuming $\int_{(1,\infty)} e^{zK_0(t)s} \lambda(dz) < \infty$, we have

$$\int_1^\infty e^{sr} \bar{\eta}_t^*(r) dr \leq c \int_{(0,\infty)} x^{-2} \pi(dx) \int_0^{1-e^{-xt}} (1-u)^{-1} du = ct. \quad \square$$

Lemma 4.2 in [22] states that if $\bar{\lambda}$ is $(-\gamma)$ -regularly varying at infinity, which we denote by $\bar{\lambda} \in \mathcal{RV}_{-\gamma}$, then $\mathbf{E}X^*(t)^q = \infty$ for all $q > \gamma$, which suggests power-law tail. The following statement makes this observation

precise, by obtaining the exact asymptotics of the tail. We note that the multivariate regular variation result for the integrated process has already been established in [36, Corollary 5.2].

Lemma 3. *Assume that $\bar{\lambda} \in \mathcal{RV}_{-\gamma}$ for some $\gamma > 0$. Further assume that there is $\varepsilon > 0$ such that*

$$\int_{(0,1)} x^{-1-\varepsilon} \pi(\mathrm{d}x) + \int_{(1,\infty)} x^{\varepsilon-\gamma} \pi(\mathrm{d}x) < \infty. \quad (15)$$

Then as $r \rightarrow \infty$

$$\bar{\eta}_t^*(r) \sim \bar{\lambda}(r) \int_{(0,\infty)} x^{-\gamma-1} \int_0^{1-e^{-xt}} y^{\gamma-1} (1-y)^{-1} \mathrm{d}y \pi(\mathrm{d}x).$$

Note that the second integral in (15) is automatically finite (i) for any $\gamma > 0$ and $\varepsilon < \gamma$ if $m_0(\pi) < \infty$, or (ii) for any $\gamma > 1$ and $\varepsilon < \gamma - 1$.

Proof. Assuming $\bar{\lambda}(z) = \ell(z)/z^\gamma$ for some ℓ slowly varying at infinity, we have

$$\int_{\frac{xr}{1-e^{-xt}}}^{\infty} \frac{\bar{\lambda}(z)}{z - xr} \mathrm{d}z = (xr)^{-\gamma} \int_0^{1-e^{-xt}} y^{\gamma-1} (1-y)^{-1} \ell(xr/y) \mathrm{d}y.$$

Note that

$$\lim_{x \downarrow 0} x^{-\gamma} \int_0^{1-e^{-xt}} y^{\gamma-1} (1-y)^{-1} \mathrm{d}y = t^\gamma \gamma^{-1}.$$

Thus, by Lemma 1

$$\begin{aligned} \bar{\eta}_t^*(r) &= \frac{\ell(r)}{r^\gamma} \int_{(0,\infty)} x^{-\gamma-1} \int_0^{1-e^{-xt}} y^{\gamma-1} (1-y)^{-1} \frac{\ell(xr/y)}{\ell(r)} \mathrm{d}y \pi(\mathrm{d}x) \\ &\sim \bar{\lambda}(r) \int_{(0,\infty)} x^{-\gamma-1} \int_0^{1-e^{-xt}} y^{\gamma-1} (1-y)^{-1} \mathrm{d}y \pi(\mathrm{d}x), \quad \text{as } r \rightarrow \infty, \end{aligned}$$

where the last asymptotic equality follows from Potter's bounds [7, Theorem 1.5.6], the integrability assumption, and Lebesgue's dominated convergence theorem. $\square$

4 Proof of Theorem 1

We will first consider the case when $\lambda((-\infty, 0)) = 0$. Note that through this section we assume that $\int_{(0,1]} z\lambda(dz) < \infty$ and that $a = \int_{(0,1]} z\lambda(dz)$, hence there is no centering. The integrated process then has the form

$$X^*(t) = \int_{(0,\infty)} \int_{\mathbb{R}} \int_{(0,\infty)} z f_t(x, s) \mu(dx, ds, dz),$$

and we can decompose it as

$$\begin{aligned} X^*(t) &= \int_{(0,\infty)} \int_{(-\infty,0]} \int_{(0,\infty)} zx^{-1}(1 - e^{-xt})e^{xs} \mu(dx, ds, dz) \\ &\quad + \int_{(0,\infty)} \int_{(0,t]} \int_{(0,\infty)} zx^{-1} \mu(dx, ds, dz) \\ &\quad - \int_{(0,\infty)} \int_{(0,t]} \int_{(0,\infty)} zx^{-1}e^{-x(t-s)} \mu(dx, ds, dz) \\ &=: X_-^*(t) + X_{+,1}^*(t) - X_{+,2}^*(t). \end{aligned} \tag{16}$$

We now consider each of the terms separately. The term X_-^* corresponding to negative jump times is easy to handle, and it is treated in Lemma 4. The main term $X_{+,1}^*$ is a subordinator, a nonnegative increasing Lévy process. We apply the classical Marcinkiewicz–Zygmund theorem, and its process version, to determine the almost sure growth rate of $X_{+,1}^*$. This part is done in Lemma 5. The difficult part is to handle the other negligible term $X_{+,2}^*$ for $\gamma \geq 1$. We need a fine analysis of the Poisson random measure μ : we decompose the space $(0, \infty) \times (0, t] \times (0, \infty)$ to countably many pieces according to the jump sizes and obtain sharp bounds on the number of points in each piece. This is done in Lemma 7, with the necessary bounds in Lemma 6 and Corollary 3. This part of the proof is similar to the techniques used in [10].

Let $(\xi_k, \tau_k, \zeta_k)_{k \geq 0}$ denote the points of the Poisson random measure μ . Then

$$X_-^*(t) = \sum_{\tau_k \leq 0} \frac{\zeta_k}{\xi_k} (1 - e^{-\xi_k t}) e^{\xi_k \tau_k}.$$

Lemma 4. *Let $\gamma \in (0, 2]$. For $\gamma > 1$ suppose that $\int_{(0,1)} x^{-\gamma} \pi(dx) < \infty$ and $\int_{(0,\infty)} z\lambda(dz) < \infty$. Then*

$$\lim_{t \rightarrow \infty} \frac{X_-^*(t)}{t^{1/\gamma}} = 0 \quad a.s.$$

Proof. Let first $\gamma \in (0, 1]$ and note that since $1 - e^{-x} \leq x$ we have

$$t^{-1} X_-^*(t) \leq t^{-1} \sum_{\tau_k < 0} \frac{\zeta_k}{\xi_k} \xi_k t e^{\xi_k \tau_k} \leq \sum_{\tau_k < 0} \zeta_k e^{\xi_k \tau_k}.$$

If the latter sum is finite a.s., then Lebesgue's dominated convergence theorem gives the result. The latter sum exists if and only if

$$\iiint_{(0, \infty)^3} (ze^{-sx} \wedge 1) \nu(dx, ds, dz) < \infty.$$

Simple calculation gives that

$$\begin{aligned} & \iiint_{(0, \infty)^3} (ze^{-sx} \wedge 1) \nu(dx, ds, dz) \\ &= \int_{(0, \infty)} x^{-1} \pi(dx) \left(\int_{(0, 1]} z \lambda(dz) + \int_{(1, \infty)} (1 + \log z) \lambda(dz) \right), \end{aligned}$$

thus the statement follows.

For $\gamma \in (1, 2)$, since $X_-^*(t)$ is a Poisson integral, we have

$$\begin{aligned} \mathbf{E} X_-^*(t) &= \iiint \frac{z}{x} (1 - e^{-tx}) e^{xs} \nu(dx, ds, dz) \\ &= m_1(\lambda) \int_{(0, \infty)} x^{-2} (1 - e^{-tx}) \pi(dx). \end{aligned}$$

By the assumption and Lebesgue's dominated convergence theorem

$$\begin{aligned} \int_{(0, \infty)} x^{-2} (1 - e^{-tx}) \pi(dx) &= t^{2-\gamma} \int_{(0, \infty)} x^{-\gamma} \frac{1 - e^{-tx}}{(tx)^{2-\gamma}} \pi(dx) \\ &= t^{2-\gamma} o(1), \quad t \rightarrow \infty, \end{aligned}$$

as the fraction in the integrand is bounded and tends to 0. Summarizing, as $t \rightarrow \infty$

$$\mathbf{E} X_-^*(t) = t^{2-\gamma} o(1).$$

Therefore, by Markov's inequality

$$\mathbf{P}(X_-^*(t) > \varepsilon t^{1/\gamma}) \leq \frac{\mathbf{E} X_-^*(t)}{\varepsilon t^{1/\gamma}} \leq t^{2-\gamma-\gamma^{-1}} o(1),$$

and by the Borel–Cantelli lemma, for $a > 0$ such that $\gamma + \gamma^{-1} - 2 > 1/a$, with $t_n = n^a$

$$\lim_{n \rightarrow \infty} t_n^{-1/\gamma} X_-^*(t_n) = 0.$$

The convergence now follows by monotonicity, as for $t \in [t_n, t_{n+1}]$

$$t^{-1/\gamma} X_-^*(t) \leq t_n^{-1/\gamma} X_-^*(t_{n+1}) = t_{n+1}^{-1/\gamma} X_-^*(t_{n+1}) \left(\frac{t_{n+1}}{t_n} \right)^{1/\gamma} \rightarrow 0.$$

The same computation shows that for $\gamma = 2$

$$\lim_{t \rightarrow \infty} \mathbf{E} X_-^*(t) = m_1(\lambda) m_{-2}(\pi).$$

Therefore the same reasoning works with $a = 3$. $\square$

Remark 2. Under the stronger assumption

$$\iint_{(0, \infty)^2} \left(\mathbf{1}(z \leq x) \frac{z}{x^2} + \mathbf{1}(z > x) \frac{1}{x} \left(1 + \log \frac{z}{x} \right) \right) \pi(dx) \lambda(dz) < \infty, \quad (17)$$

$\sup_{t \geq 0} X_-^*(t) < \infty$ a.s. Indeed, $\sup_{t \geq 0} X_-^*(t) \leq \sum_{\tau_k \leq 0} \frac{\zeta_k}{\xi_k} e^{\xi_k \tau_k}$, which is finite a.s. if and only if (17) holds.

Next we handle the main term.

Lemma 5. *Assume that for some $\gamma \in (0, 2)$*

$$\iint_{(0, \infty)^2} \left(\frac{z}{x} \right)^\gamma \mathbf{1}(z > x) \pi(dx) \lambda(dz) < \infty. \quad (18)$$

Then

$$\lim_{t \rightarrow \infty} \frac{X_{+,1}^*(t) - \mathbf{1}(\gamma \geq 1) t m_1(\lambda) m_{-1}(\pi)}{t^{1/\gamma}} = 0 \quad a.s. \quad (19)$$

If (18) holds with $\gamma = 2$, then the law of iterated logarithm holds, i.e.

$$\limsup_{t \rightarrow \infty} \frac{|X_{+,1}^*(t) - t m_1(\lambda) m_{-1}(\pi)|}{\sqrt{2t \log \log t}} = \sqrt{m_2(\lambda) m_{-2}(\pi)} \quad a.s.$$

Proof. Note that $X_{+,1}^*(t)$ is a subordinator with characteristic function

$$\mathbf{E} e^{iu X_{+,1}^*(t)} = \exp \left\{ t \int (e^{iuy} - 1) \eta_1(dy) \right\},$$

where the Lévy measure η_1 is given by

$$\eta_1((r, \infty)) = \bar{\eta}_1(r) = \pi \times \lambda(\{(x, z) : z > rx\}) = \int_{(0, \infty)} \bar{\lambda}(rx) \pi(dx).$$

Thus $\mathbf{E}X_{+,1}^*(1)^\gamma < \infty$ if and only if $\int_{(1,\infty)} r^\gamma \eta_1(dr) < \infty$ which holds if and only if (18) holds. Therefore, the Marcinkiewicz–Zygmund strong law of large numbers implies that (19) holds whenever (18) holds with $0 < \gamma < 2$. Indeed, if $\gamma < 1$, then by monotonicity the result follows from the usual partial sum version ([25, Theorem 6.7.1]). For $\gamma \in [1, 2)$, Theorem 2.1 in [47], the process version of the Marcinkiewicz–Zygmund SLLN, gives the result since

$$\mathbf{E}X_{+,1}^*(1) = \int_0^\infty \bar{\eta}_1(r) dr = m_1(\lambda)m_{-1}(\pi).$$

The statement for $\gamma = 2$ follows from the law of iterated logarithm for Lévy processes (see Proposition 48.9 in [44]) and the fact that

$$\mathbf{Var}(X_{+,1}^*(1)) = \iiint_{(0,\infty) \times (0,1] \times (0,\infty)} \frac{z^2}{x^2} \nu(dx, ds, dz) = m_2(\lambda)m_{-2}(\pi),$$

see e.g. [33, Theorem 2.7]. $\square$

We note that in case of infinite mean ($\gamma < 1$) more precise integral tests are known, see [6, Theorem III.13].

It remains to consider $X_{+,2}^*$. For $t > 0$, $r > 0$ and $r_2 > r_1 > 0$ introduce the notation

$$D(r, t) = \left\{ (x, s, z) : \frac{z}{x} e^{-x(t-s)} > r, \ s \in (0, t), x > 0, z > 0 \right\},$$

$$D(r_1, r_2, t) = \left\{ (x, s, z) : \frac{z}{x} e^{-x(t-s)} \in (r_1, r_2], \ s \in (0, t), x > 0, z > 0 \right\}.$$

Lemma 6. *For $t > 1$ and $0 < r_1 < r_2 < \infty$*

$$\nu(D(r_1, r_2, t)) \leq t \int_{(0, t^{-1}]} \bar{\lambda}(r_1 x) \pi(dx) + \log \frac{r_2}{r_1} \int_{(t^{-1}, \infty)} x^{-1} \bar{\lambda}(r_1 x) \pi(dx).$$

Proof. Since $\frac{z}{x} e^{-xu} \in (r_1, r_2]$ for $u > 0$ if and only if (i) $\frac{z}{x} \in (r_1, r_2]$ and $x^{-1} \log \frac{z}{r_1 x} > u$, or (ii) $\frac{z}{x} > r_2$ and $x^{-1} \log \frac{z}{r_2 x} \leq u < x^{-1} \log \frac{z}{r_1 x}$, we have

$$\begin{aligned} \nu(D(r_1, r_2, t)) &= \iint \mathbf{1}(r_1 x < z \leq r_2 x) \left(t \wedge x^{-1} \log \frac{z}{r_1 x} \right) \\ &\quad + \mathbf{1}(z > r_2 x) \left(t \wedge x^{-1} \log \frac{z}{r_1 x} - t \wedge x^{-1} \log \frac{z}{r_2 x} \right) \pi(dx) \lambda(dz) \\ &\leq \iint \mathbf{1}(r_1 x < z) \left(t \wedge x^{-1} \log \frac{r_2}{r_1} \right) \pi(dx) \lambda(dz). \end{aligned}$$

Bounding the expression in the bracket by t for $x \in (0, t^{-1}]$ and by $x^{-1} \log \frac{r_2}{r_1}$ for $x \in (t^{-1}, \infty)$, we obtain the statement. $\square$

If $\gamma = 1$, we need an extra assumption:

$$\exists \delta > 0 : \quad m_{-1-\delta}(\pi) < \infty. \quad (20)$$

Corollary 3. *Assume (18) with $\gamma \in [1, 2]$, and for $\gamma = 1$ also assume (20). There exists $C = C(\pi, \lambda, \gamma)$ such that for any $0 \leq a < b$*

$$\nu(D(n^a, n^b, n)) \leq \begin{cases} Cn^{2-\gamma-a} \log n, & \gamma > 1, \\ Cn^{1-a-\delta/2}, & \gamma = 1. \end{cases}$$

Proof. Note that (18) with $\gamma \in [1, 2]$ implies that $m_{-\gamma}(\pi) < \infty$ and $m_1(\lambda) < \infty$. For $\gamma > 1$ we have

$$\begin{aligned} n \int_{(0, n^{-1}]} \pi(dx) \int_{(n^a x, \infty)} \lambda(dz) &\leq n \int_{(0, n^{-1}]} n^{-a} x^{-1} \pi(dx) m_1(\lambda) \\ &\leq m_1(\lambda) n^{1-a} \int_{(0, n^{-1}]} x^{-\gamma} n^{1-\gamma} \pi(dx) \\ &= m_{-\gamma}(\pi) m_1(\lambda) n^{2-\gamma-a}, \end{aligned}$$

and similarly,

$$\begin{aligned} \int_{(n^{-1}, \infty)} x^{-1} \pi(dx) \int_{(n^a x, \infty)} \lambda(dz) &\leq n^{-a} \int_{(n^{-1}, \infty)} x^{-2} \pi(dx) m_1(\lambda) \\ &\leq m_{-\gamma}(\pi) m_1(\lambda) n^{2-\gamma-a}. \end{aligned}$$

The statement now follows from Lemma 6. For $\gamma = 1$ the proof is the same, only in the $d\pi$ integral we need $x^{-1-\delta}$. $\square$

Lemma 7. *Suppose that $\gamma \in (0, 1)$, or $\gamma \in [1, 2]$ and (18) holds. If $\gamma = 1$, assume additionally that (20) holds. Then*

$$\lim_{t \rightarrow \infty} t^{-1/\gamma} X_{+,2}^*(t) = 0 \quad \text{a.s.}$$

Proof. If $\gamma < 1$ the statement follows from Lemma 5 since $X_{+,1}^*(t) \geq X_{+,2}^*(t)$. Therefore we assume that $\gamma \geq 1$.

Put $Y_n = \sum_{\tau_k \in [n-1, n)} \frac{\zeta_k}{\xi_k}$. Then $Y_1, Y_2, \dots$ is an iid sequence with the same distribution as $X_{+,1}^*(1)$, thus $\mathbf{E}Y_1^\gamma < \infty$ by (18). Therefore,

$$n^{-1/\gamma} \max_{i \leq n} Y_i \rightarrow 0 \quad \text{a.s.} \quad (21)$$

(see e.g. [12, Theorem 3.5.1]). Furthermore, $\max_{\tau_k < n} \frac{\zeta_k}{\xi_k} \leq \max_{i \leq n} Y_i$.

First we prove that

$$\lim_{n \rightarrow \infty} n^{-1/\gamma} X_{+,2}^*(n) = 0 \quad \text{a.s.} \quad (22)$$

Assume that $\gamma > 1$. Let $c_0 = \gamma + \gamma^{-1} - 2 > 0$, and choose a_0 so that $2 - \gamma < a_0 < 2 - \gamma + c_0/2$, and let $m = \lfloor 2a_0/c_0 \rfloor$, with $\lfloor \cdot \rfloor$ standing for the lower integer part. Let N be an integer large enough to have $N(2 - \gamma - a_0) < -1$. Define the sets

$$\begin{aligned} A_{n,0} &= \{\mu(D(n^{a_0}, n)) > N\}, \\ A_{n,i+1} &= \{\mu(D(n^{a_{i+1}}, n^{a_i}, n)) > 2C \log n n^{2-\gamma-a_{i+1}}\}, \quad i = 0, \dots, m, \end{aligned} \quad (23)$$

where $a_{i+1} = a_i - c_0/2$ for $i = 0, 1, \dots, m-1$, $a_{m+1} = 0$, and $C = C(\pi, \lambda, \gamma)$ is the constant in Corollary 3. With this choice $a_m = a_0 - mc_0/2 \in [0, c_0/2)$.

We use the Poisson tail bounds (see [10, Lemma 3.1])

$$\begin{aligned} \mathbf{P}(N_\lambda \geq n) &\leq \frac{\lambda^n}{n!}, \\ \mathbf{P}(N_\lambda \geq x) &\leq e^{-0.19x}, \quad x \geq 2\lambda, \end{aligned} \quad (24)$$

where N_λ stands for a Poisson random variable with parameter λ . By Corollary 3 (with $a = a_0$ and $b = \infty$) and (24)

$$\mathbf{P}(A_{n,0}) \leq C n^{N(2-\gamma-a_0)} (\log n)^N,$$

and, similarly for $i = 1, 2, \dots, m+1$ using the second bound in (24)

$$\mathbf{P}(A_{n,i}) \leq \exp \left\{ -0.19 \cdot 2C \log n n^{2-\gamma-a_i} \right\}.$$

Clearly, all these bounds are summable, therefore, by the first Borel–Cantelli lemma the events $(A_{n,i})_{n \geq 1, i=0,1,\dots,m+1}$ occur finitely many times almost surely.

The contribution of the points (ξ_k, τ_k, ζ_k) for which $\zeta_k/\xi_k e^{-\xi_k(n-\tau_k)} \in (n^{a_{i+1}}, n^{a_i}]$, $\tau_k < n$, if n is large enough so that $A_{n,i+1}$ does not occur, is for $i = 0, 1, \dots, m$

$$\begin{aligned} \sum_{\substack{0 < \tau_k < n, \\ \zeta_k/\xi_k e^{-\xi_k(n-\tau_k)} \in (n^{a_{i+1}}, n^{a_i}]}} \frac{\zeta_k}{\xi_k} e^{-\xi_k(n-\tau_k)} &\leq n^{a_i} 2C \log n n^{2-\gamma-a_{i+1}} \\ &\leq 2C \log n n^{2-\gamma+c_0/2}, \end{aligned} \quad (25)$$

by the choice of c_0 , where the last inequality is in fact equality for $i \neq m$ (as $a_m - a_{m+1} = a_m \in [0, c_0/2)$).

The events in (23) take care of the contributions larger than 1. Next we deal with the small contributions. Introduce the events

$$B_{n,k} = \{\mu(D(2^{-k-1}, 2^{-k}, n)) > b_{n,k}\}, \quad k = 0, 1, \dots \quad (26)$$

with

$$\begin{aligned} b_{n,k} &= 6 \log(n(k+1)) + 2n \int_{(0, n^{-1}]} \pi(dx) \int_{(2^{-k-1}x, \infty)} \lambda(dz) \\ &\quad + 2 \int_{(n^{-1}, \infty)} x^{-1} \pi(dx) \int_{(2^{-k-1}x, \infty)} \lambda(dz). \end{aligned}$$

By Lemma 6 and (24) we have

$$\mathbf{P}(B_{n,k}) \leq \exp\{-6 \cdot 0.19 \cdot \log(n(k+1))\} \leq (n(k+1))^{-1.1},$$

which is summable for $k \geq 0$, $n \geq 1$. The first Borel–Cantelli lemma implies that all these events occur finitely many times a.s. Note that in $b_{n,k}$ the term $6 \log(n(k+1))$ ensures the summability of the probabilities even if the other terms in $b_{n,k}$ are small.

The sum of small contributions such that $\zeta_j/\xi_j e^{-\xi_j(n-\tau_j)} \in (2^{-k-1}, 2^{-k}]$, $\tau_j < n$, for $k = 0, 1, \dots$ and n large enough so that $B_{n,k}$ does not occur, is bounded by

$$\begin{aligned} \sum_{k=0}^{\infty} 2^{-k} b_{n,k} &\leq c \log n + 2n \int_{(0, n^{-1}]} \pi(dx) \sum_{k=0}^{\infty} 2^{-k} \int_{(2^{-k-1}x, \infty)} \lambda(dz) \\ &\quad + 2 \int_{(n^{-1}, \infty)} x^{-1} \pi(dx) \sum_{k=0}^{\infty} 2^{-k} \int_{(2^{-k-1}x, \infty)} \lambda(dz). \end{aligned}$$

The infinite sum can be bounded as

$$\begin{aligned} \sum_{k=0}^{\infty} 2^{-k} \int_{(2^{-k-1}x, \infty)} \lambda(dz) &= \int_{(0, \infty)} \sum_{k=0}^{\infty} 2^{-k} \mathbf{1}(z > 2^{-k-1}x) \lambda(dz) \\ &\leq 2 \int_{(0, \infty)} (1 \wedge zx^{-1}) \lambda(dz) \leq cx^{-1}. \end{aligned}$$

Substituting back, and using $\int_{(0, n^{-1}]} x^{-1} \pi(dx) \leq n^{1-\gamma} m_{-\gamma}(\pi)$, and

$$\int_{(n^{-1}, \infty)} x^{-2} \pi(dx) \leq n^{2-\gamma} m_{-\gamma}(\pi),$$

we have

$$\sum_{k=0}^{\infty} 2^{-k} b_{n,k} \leq c \log n + cn^{2-\gamma}. \quad (27)$$

Summarizing, by (25) and (27) the contribution of the terms in $X_{+,2}^*(n)$ can be bounded by

$$X_{+,2}^*(n) \leq W + N \max_{1 \leq k \leq n} Y_k + c \log n m n^{2-\gamma+c_0/2} + cn^{2-\gamma},$$

where the random variable W is coming from the Borel–Cantelli lemma, and the second term comes from $A_{n,0}$. By (21) all the terms are $o(n^{1/\gamma})$, thus (22) follows.

For $t \in (n, n+1)$, using (21)

$$t^{-1/\gamma} X_{+,2}^*(t) \leq n^{-1/\gamma} (X_{+,2}^*(n) + Y_{n+1}) \rightarrow 0 \quad \text{a.s.}$$

as claimed.

For $\gamma = 1$ the same idea works. Recall $\delta > 0$ from (20), and let $c_0 = \delta/2$, $a_0 \in (1 - \delta/2, 1 - \delta/4)$, $m = \lfloor 4a_0/\delta \rfloor$. Choose an integer N such that $N(1 - a_0 - \delta/2) < -1$. Define the events $A_{n,i}$'s similarly as in (23),

$$\begin{aligned} A_{n,0} &= \{\mu(D(n^{a_0}, n)) > N\}, \\ A_{n,i+1} &= \{\mu(D(n^{a_{i+1}}, n^{a_i}, n)) > 2C n^{1-\delta/2-a_{i+1}}\}, \quad i = 0, \dots, m, \end{aligned}$$

where $a_{i+1} = a_i - c_0/2$ for $i = 0, 1, \dots, m-1$, $a_{m+1} = 0$, and $C = C(\pi, \lambda, 1)$ is the constant in Corollary 3. As above, A_n 's occur finitely many times a.s., and the contribution is $o(n)$. For the small jumps define $B_{n,k}$'s as in (26). The only difference is that by assumption (20)

$$\int_{(0, n^{-1}]} x^{-1} \pi(dx) \leq n^{-\delta} m_{-1-\delta}(\pi),$$

and

$$\int_{(n^{-1}, \infty)} x^{-2} \pi(dx) \leq n^{1-\delta} m_{-1-\delta}(\pi). \quad \square$$

Proof of Theorem 1.

1. If $\lambda((-\infty, 0)) = 0$, then the proof follows directly from Lemmas 4, 5, 7 and decomposition (16). Note that the centering term in (19) is indeed $\mathbf{E}X^*(t)$ by (9). If $\lambda((-\infty, 0)) \neq 0$, then we decompose $X^*(t)$ as in (16) but separately for positive and negative jumps, that is considering $\mu\mathbf{1}(z > 0)$ and $\mu\mathbf{1}(z < 0)$ separately. This gives

$$X^*(t) = X_-^{*,+}(t) - X_-^{*, -}(t) + X_{+,1}^{*,+}(t) - X_{+,1}^{*, -}(t) - X_{+,2}^{*,+}(t) + X_{+,2}^{*, -}(t).$$

For $\gamma \in (0, 2)$ the statement now follows by applying Lemmas 4, 5 and 7 to positive and negative jump parts separately.

2. If $\gamma = 2$, then the proof follows as for $\gamma \in (0, 2)$. In general, we decompose $X_-^*(t)$ and $X_{+,2}^*(t)$ as in the previous case. When divided by $\sqrt{2t \log \log t}$ all of these terms converge to 0 a.s. The statement now follows from Proposition 48.9 in [44] applied to $X_{+,1}^*(t)$. $\square$

5 Proof of Theorem 2

We shall first prove the theorem by assuming that λ is supported on $(0, 1]$. At the very end we will then combine this with Theorem 1 to prove Theorem 2.

If $\int_{(0,1]} z\lambda(dz) = \infty$, then centering is needed and we have

$$X^*(t) = \iiint_{(0,\infty) \times (-\infty, t] \times (0,1]} f_t(x, s) z(\mu - \nu)(dx, ds, dz) =: X_+^*(t) - X_-^*(t),$$

where

$$\begin{aligned} X_+^*(t) &= \iiint_{(0,\infty) \times (0,t] \times (0,1]} \frac{z}{x} \left(1 - e^{-x(t-s)}\right) (\mu - \nu)(dx, ds, dz), \\ X_-^*(t) &= \iiint_{(0,\infty) \times (-\infty, 0] \times (0,1]} \frac{z}{x} \left(1 - e^{-xt}\right) e^{xs} (\mu - \nu)(dx, ds, dz). \end{aligned}$$

The following two statements, Lemmas 8 and 9, ensure that the paths are sufficiently smooth, so that it is enough to prove almost sure limits on the integers. Then we decompose the process further to a part with only small jumps $X_{\pm, < \kappa}^*$ and with only large jumps $X_{\pm, > \kappa}^*$. The small jump part is analysed via its moment generating function in Corollary 4, where the necessary bounds are established in Lemma 10. This part ruins the optimality, as an extra logarithmic factor appears. The large jump part is handled in Lemmas 11 and 13. The truncation idea resembles the proof of the classical Kolmogorov and Marcinkiewicz–Zygmund strong law of large numbers, see e.g. [25].

Lemma 8. *There exists a modification of $X_{\pm}^*$ such that for any $\theta \in [0, 1/2)$*

$$\sup_n \mathbf{E} \left(\sup_{s \neq t, s, t \in [n, n+1]} \frac{|X_+^*(t) - X_+^*(s)| + |X_-^*(t) - X_-^*(s)|}{|t - s|^\theta} \right)^2 < \infty.$$

Proof. For $t_1 < t_2$ we have

$$\begin{aligned} & X_+^*(t_2) - X_+^*(t_1) \\ &= \iiint_{(0, \infty) \times (0, t_1] \times (0, 1]} \frac{z}{x} \left(e^{-x(t_1-s)} - e^{-x(t_2-s)} \right) (\mu - \nu)(dx, ds, dz) \\ &+ \iiint_{(0, \infty) \times (t_1, t_2] \times (0, 1]} \frac{z}{x} \left(1 - e^{-x(t_2-s)} \right) (\mu - \nu)(dx, ds, dz), \end{aligned}$$

therefore, by Theorem 1 in [35]

$$\begin{aligned} & \mathbf{E} \left(X_+^*(t_2) - X_+^*(t_1) \right)^2 \\ & \leq 2 \iiint_{(0, \infty) \times (0, t_1] \times (0, 1]} \frac{z^2}{x^2} e^{-2x(t_1-s)} \left(1 - e^{-x(t_2-t_1)} \right)^2 \nu(dx, ds, dz) \\ &+ 2 \iiint_{(0, \infty) \times (t_1, t_2] \times (0, 1]} \frac{z^2}{x^2} \left(1 - e^{-x(t_2-s)} \right)^2 \nu(dx, ds, dz) =: I_1 + I_2. \end{aligned}$$

Writing $\Delta t = t_2 - t_1$, evaluating the ds integral, and using that $1 - e^{-y} \leq y$, the first term on the right-hand side can be bounded as

$$I_1 \leq 2(\Delta t)^2 \iint_{(0, \infty) \times (0, 1]} \frac{z^2}{2x} (1 - e^{-2xt_1}) \lambda(dz) \pi(dx) \leq m_2(\lambda) m_{-1}(\pi) (\Delta t)^2.$$

For the second term we have

$$\begin{aligned} I_2 &= 2m_2(\lambda) \int_{(0, \infty)} x^{-2} \pi(dx) \int_0^{\Delta t} (1 - e^{-xu})^2 du \\ &\leq 2m_2(\lambda) \Delta t \int_{(0, \infty)} x^{-2} (x \Delta t \wedge 1)^2 \pi(dx) \\ &\leq 2m_2(\lambda) m_{-1}(\pi) (\Delta t)^2. \end{aligned}$$

Summarizing

$$\mathbf{E} (X_+^*(t_2) - X_+^*(t_1))^2 \leq 3m_2(\lambda) m_{-1}(\pi) (\Delta t)^2. \quad (28)$$

Similarly, for X_-^* we have

$$\begin{aligned}
& \mathbf{E} \left(X_-^*(t_2) - X_-^*(t_1) \right)^2 \\
& \leq 2 \iiint_{(0,\infty) \times (-\infty,0] \times (0,1]} \frac{z^2}{x^2} e^{2xs} e^{-2xt_1} (1 - e^{-x\Delta t})^2 \nu(dx, ds, dz) \\
& = m_2(\lambda) \int_{(0,\infty)} \frac{1}{x^3} e^{-2xt_1} (1 - e^{-x\Delta t})^2 \pi(dx) \\
& \leq m_2(\lambda) m_{-1}(\pi) (t_2 - t_1)^2.
\end{aligned} \tag{29}$$

The claim follows from a version of Kolmogorov's continuity theorem ([27, Theorem 4.3] with $[a, b] = [n, n+1]$, $\gamma = p = 2$, $\theta < 1/2$). Note that (28) and (29) hold for any $t_1, t_2 \in (0, \infty)$, which implies that the bound above is uniform in n . $\square$

Lemma 9. *Assume that for a process $(Y(t))_{t \in \mathbb{R}}$*

$$\sup_{n \in \mathbb{N}} \mathbf{E} \left(\sup_{t \in [n, n+1]} |Y(t) - Y(n)| \right)^2 < \infty,$$

and for an increasing function $a(t)$ for which

$$\sum_n a(n)^{-2} < \infty, \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{a(n)}{a(n+1)} = 1,$$

we have $\limsup_{n \rightarrow \infty} |Y(n)|/a(n) \leq c$ a.s., with some $c \geq 0$. Then

$$\lim_{t \rightarrow \infty} \frac{|Y(t)|}{a(t)} \leq c \quad \text{a.s.}$$

Proof. By Markov's inequality

$$\mathbf{P} \left(\sup_{t \in [n, n+1]} |Y(t) - Y(n)| > a(n)\varepsilon \right) \leq \frac{\mathbf{E} \left(\sup_{t \in [n, n+1]} |Y(t) - Y(n)| \right)^2}{a(n)^2 \varepsilon^2},$$

thus the first Borel-Cantelli lemma implies the statement. $\square$

We shall frequently use the following estimates

$$\begin{aligned}
\int_{(y,1]} z^u \lambda(dz) &\leq y^{u-\beta} \int_{(0,1]} z^\beta \lambda(dz), \quad u \leq \beta, \\
\int_{(0,y]} z^u \lambda(dz) &\leq y^{u-\beta} \int_{(0,1]} z^\beta \lambda(dz), \quad u \geq \beta, \\
\int_{(y,1]} x^{-v} \pi(dx) &\leq y^{1+\alpha-v} \int_{(0,1]} x^{-1-\alpha} \pi(dx), \quad v \geq 1 + \alpha, \\
\int_{(0,y]} x^{-v} \pi(dx) &\leq y^{1+\alpha-v} \int_{(0,1]} x^{-1-\alpha} \pi(dx), \quad v \leq 1 + \alpha,
\end{aligned} \tag{30}$$

with α and β defined in the paragraph below (11).

We decompose X_+^* and X_-^* further. For $\kappa \in (0, 1]$, put

$$\begin{aligned}
A(t; \kappa) &= \left\{ (x, s, z) : \frac{z}{x} \left(1 - e^{-x(t-s)} \right) \leq t^\kappa, s \in (0, t], x > 0, z \in (0, 1] \right\}, \\
B(t; \kappa) &= \left\{ (x, s, z) : \frac{z}{x} e^{xs} \left(1 - e^{-xt} \right) \leq t^\kappa, s \leq 0, x > 0, z \in (0, 1] \right\},
\end{aligned}$$

and define the truncated processes

$$\begin{aligned}
X_{+,<,\kappa}^*(t) &= \iiint_{A(t;\kappa)} \frac{z}{x} \left(1 - e^{-x(t-s)} \right) (\mu - \nu)(dx, ds, dz), \\
X_{+,>,\kappa}^*(t) &= \iiint_{A(t;\kappa)^c} \frac{z}{x} \left(1 - e^{-x(t-s)} \right) (\mu - \nu)(dx, ds, dz),
\end{aligned}$$

and

$$\begin{aligned}
X_{-,<,\kappa}^*(t) &= \iiint_{B(t;\kappa)} \frac{z}{x} e^{xs} \left(1 - e^{-xt} \right) (\mu - \nu)(dx, ds, dz), \\
X_{-,>,\kappa}^*(t) &= \iiint_{B(t;\kappa)^c} \frac{z}{x} e^{xs} \left(1 - e^{-xt} \right) (\mu - \nu)(dx, ds, dz).
\end{aligned}$$

Here, to ease notation the complement is understood as $A(t; \kappa)^c = (0, \infty) \times (0, t] \times (0, 1] \setminus A(t; \kappa)$, and $B(t; \kappa)^c = (0, \infty) \times (-\infty, 0] \times (0, 1] \setminus B(t; \kappa)$. Note that the truncated processes do depend on κ . The truncation level $\kappa = 1$ means no truncation, that is $X_{\pm,<,1}^*(t) = X_{\pm}^*(t)$.

Lemma 10. For $|\theta| \leq 4t^{-\kappa}$

$$\log \mathbf{E} e^{\theta X_{+,<,\kappa}^*(t)} \leq \begin{cases} c\theta^2 t^{\beta-\alpha+\kappa(2-\beta)}, & \text{if } \alpha \leq \beta - 1, \\ c\theta^2 t^{1+\kappa(1-\alpha)_+}, & \text{if } \alpha \geq \beta - 1, \end{cases}$$

and

$$\log \mathbf{E} e^{\theta X_{-, <, \kappa}^*(t)} \leq \begin{cases} c\theta^2 t^{\beta-\alpha+\kappa(2-\beta)}, & \text{if } \alpha \leq \beta, \\ c\theta^2 t^{\kappa(2-\alpha)+}, & \text{if } \alpha \geq \beta. \end{cases}$$

Proof. For the moment generating function we have

$$\mathbf{E} e^{\theta X_{\pm, <, \kappa}^*(t)} = \exp \left\{ \int \left(e^{\theta y} - 1 - \theta y \right) \Pi_t^\pm(dy) \right\},$$

where the Lévy measures $\Pi_t^\pm$ are given by

$$\begin{aligned} \Pi_t^+(D) &= \nu \left(\left\{ (x, s, z) \in A(t; \kappa) : \frac{z}{x} (1 - e^{-x(t-s)}) \in D \right\} \right), \\ \Pi_t^-(D) &= \nu \left(\left\{ (x, s, z) \in B(t; \kappa) : \frac{z}{x} e^{xs} (1 - e^{-xt}) \in D \right\} \right), \end{aligned}$$

for any Borel set D . The moment generating function is finite for any $\theta \in \mathbb{R}$, see e.g. [44, Section 26]. Using the inequality $e^u - 1 - u \leq 4u^2$ for $|u| \leq 4$, and that $\Pi_t^+((t^\kappa, \infty)) = 0$, we obtain for $|t^\kappa \theta| \leq 4$

$$\begin{aligned} \log \mathbf{E} e^{\theta X_{+, <, \kappa}^*(t)} &\leq \int 4\theta^2 y^2 \Pi_t^+(dy) \\ &= \iiint_{(0, \infty) \times (0, t] \times (0, 1]} 4\theta^2 \frac{z^2}{x^2} (1 - e^{-xu})^2 \mathbb{1} \left(\frac{z}{x} (1 - e^{-xu}) \leq t^\kappa \right) \nu(dx, du, dz) \\ &\leq 4\theta^2 \iiint_{(0, \infty) \times (0, t] \times (0, 1]} \left[\mathbb{1}(xu \leq 1, zu \leq 2t^\kappa) z^2 u^2 \right. \\ &\quad \left. + \mathbb{1}(xu > 1, z \leq 2t^\kappa x) z^2 x^{-2} \right] \nu(dx, du, dz) \\ &=: 4\theta^2 (I_1 + I_2). \end{aligned}$$

For I_1 we have from (30)

$$\begin{aligned} I_1 &\leq \int_0^1 u du \int_{(0, 1]} z^2 \lambda(dz) m_{-1}(\pi) + \int_1^t u^2 du \int_{(0, u^{-1}]} \pi(dx) \int_{(0, 2t^\kappa/u]} z^2 \lambda(dz) \\ &\leq c \left(1 + \int_1^{2t^\kappa} u^{1-\alpha} du + \mathbb{1}(2t^\kappa < t) \int_{2t^\kappa}^t u^{1-\alpha+\beta-2} t^{\kappa(2-\beta)} du \right) \\ &\leq c \left(1 + \mathbb{1}(\alpha < 2) t^{\kappa(2-\alpha)} + \mathbb{1}(\alpha = 2) \log t \right. \\ &\quad \left. + [\mathbb{1}(\kappa < 1, \beta = \alpha) \log t + t^{(\beta-\alpha)+}] t^{\kappa(2-\beta)} \right), \end{aligned}$$

and for I_2

$$\begin{aligned}
I_2 &\leq \int_{(t^{-1}, \infty)} tx^{-2} \pi(dx) \int_{(0, 2t^\kappa x]} z^2 \lambda(dz) \\
&\leq c \mathbf{1}(2t^\kappa < t) t^{1+\kappa(2-\beta)} \int_{(t^{-1}, t^{-\kappa}/2]} x^{-\beta} \pi(dx) + ct \int_{(t^{-\kappa}/2, \infty)} x^{-2} \pi(dx) \\
&\leq c \left(\mathbf{1}(1 + \alpha \geq \beta) t^{1+\kappa(1-\alpha)} + \mathbf{1}(1 + \alpha \leq \beta) t^{\beta-\alpha+\kappa(2-\beta)} + t^{1+\kappa(1-\alpha)+} \right) \\
&\leq \begin{cases} ct^{\beta-\alpha+\kappa(2-\beta)}, & \alpha \leq \beta - 1, \\ ct^{1+\kappa(1-\alpha)+}, & \alpha \geq \beta - 1, \end{cases}
\end{aligned}$$

proving the first statement.

Similarly, for $X_{-, <, \kappa}^*$ if $|t^\kappa \theta| \leq 4$

$$\begin{aligned}
\log \mathbf{E} e^{\theta X_{-, <, \kappa}^*(t)} &\leq \int 4\theta^2 y^2 \Pi_t^-(dy) \\
&= \iiint 4\theta^2 \frac{z^2}{x^2} e^{2xs} (1 - e^{-xt})^2 \mathbf{1}(B(t; \kappa)) \nu(dx, ds, dz) \\
&\leq 4\theta^2 \iiint_{(0, \infty) \times (-\infty, 0] \times (0, 1]} \left[\mathbf{1}(xt \leq 1, e^{xs} \leq 2t^{\kappa-1} z^{-1}) z^2 t^2 e^{2xs} \right. \\
&\quad \left. + \mathbf{1}\left(xt > 1, e^{xs} \frac{z}{x} \leq 2t^\kappa\right) z^2 x^{-2} e^{2xs} \right] \nu(dx, ds, dz) \\
&=: 4\theta^2 (J_1 + J_2).
\end{aligned}$$

Integrating with respect to s , again by (30) we get

$$\begin{aligned}
J_1 &= t^2 \int_{(0, t^{-1}]} x^{-1} \pi(dx) \left[\int_{(0, 2t^{\kappa-1}]} \frac{1}{2} z^2 \lambda(dz) + \int_{(2t^{\kappa-1}, 1]} 2t^{2(\kappa-1)} \lambda(dz) \right] \\
&\leq ct^{2-\alpha} \left[t^{-(2-\beta)(1-\kappa)} + t^{-(2-\beta)(1-\kappa)} \right] = ct^{\beta-\alpha+\kappa(2-\beta)},
\end{aligned}$$

and

$$\begin{aligned}
J_2 &= \int_{(t^{-1}, \infty)} \frac{1}{2x^3} \pi(dx) \int_{(0, 1]} z^2 \lambda(dz) \left[\mathbf{1}(2t^\kappa x \geq z) + \frac{(2t^\kappa x)^2}{z^2} \mathbf{1}(2t^\kappa x < z) \right] \\
&\leq c \mathbf{1}(2t^\kappa < t) \int_{(t^{-1}, t^{-\kappa}/2]} x^{-3} (t^\kappa x)^{2-\beta} \pi(dx) + c \int_{(t^{-\kappa}/2, \infty)} x^{-3} \pi(dx) \\
&\leq c \left(\mathbf{1}(\alpha \leq \beta) t^{\beta-\alpha+\kappa(2-\beta)} + \mathbf{1}(\alpha \geq \beta) t^{\kappa(2-\alpha)+} + t^{\kappa(2-\alpha)+} \right).
\end{aligned}$$

□

Corollary 4. For any $\kappa \in [1/2, 1]$ such that

$$\kappa \geq \begin{cases} 1 - \frac{\alpha}{\beta}, & \text{if } \alpha \leq \beta - 1, \\ \frac{1}{1+\alpha}, & \text{if } \beta - 1 \leq \alpha \leq 1, \\ \frac{1}{2}, & \text{if } \alpha \geq 1, \end{cases}$$

it holds that

$$\limsup_{t \rightarrow \infty} \frac{|X_{\pm, <, \kappa}^*(t)|}{t^\kappa \log t} \leq \frac{1}{4} \quad a.s.$$

Proof. Fix $\varepsilon > 0$. Applying Markov's inequality together with Lemma 10 with $\theta = 4t^{-\kappa}$, we obtain

$$\begin{aligned} \mathbf{P}(X_{+, <, \kappa}^*(t) > (1/4 + \varepsilon)t^\kappa \log t) &= \mathbf{P}\left(e^{\theta X_{+, <, \kappa}^*(t)} > e^{\theta(1/4 + \varepsilon)t^\kappa \log t}\right) \\ &\leq \begin{cases} \exp\{c\theta^2 t^{\beta - \alpha + \kappa(2 - \beta)} - \theta(1/4 + \varepsilon)t^\kappa \log t\} \leq ct^{-(1 + 4\varepsilon)}, & \alpha \leq \beta - 1, \\ \exp\{c\theta^2 t^{1 + \kappa(1 - \alpha)_+} - \theta(1/4 + \varepsilon)t^\kappa \log t\} \leq ct^{-(1 + 4\varepsilon)}, & \alpha \geq \beta - 1, \end{cases} \end{aligned} \quad (31)$$

where we used that $\beta - \alpha + \kappa(2 - \beta) - 2\kappa \leq 0$ for $\alpha \leq \beta - 1$, and $1 + \kappa(1 - \alpha)_+ - 2\kappa \leq 0$ for $\alpha \geq \beta - 1$. The same bound for $-X_{+, <, \kappa}^*(t)$ combined with the Borel–Cantelli lemma shows that the convergence takes place on the subsequence $t_n = n$. From the proof of Lemma 8 it is clear that the statement remains true for $X_{\pm, <, \kappa}^*(t)$, thus the result follows by Lemma 9.

For $X_{-, <, \kappa}^*$ the same argument applies. In the corresponding version of (31) one has to check that $\beta - \alpha + \kappa(2 - \beta) - 2\kappa \leq 0$ for $\alpha \leq \beta$, and $\kappa(2 - \alpha)_+ - 2\kappa \leq 0$ for $\alpha \geq \beta$. Short calculations give that these inequalities hold whenever κ satisfies the assumptions of the statement. $\square$

Lemma 11. If for some $\gamma \geq 1$

$$\iint_{(0, \infty) \times (0, 1]} \left(\frac{z}{x}\right)^\gamma \mathbf{1}(z > x) \pi(dx) \lambda(dz) < \infty,$$

then almost surely $\mu(A(t; 1/\gamma)^c) = 0$ and $\mu(B(t; 1/\gamma)^c) = 0$ for t large enough.

Proof. We have

$$\nu(\{(x, s, z) : z > xn^{1/\gamma}, s \in (n, n + 1)\}) = \iint \mathbf{1}(z > xn^{1/\gamma}) \pi(dx) \lambda(dz),$$

thus

$$\begin{aligned}
& \sum_{n=1}^{\infty} \nu(\{(x, s, z) : z > xn^{1/\gamma}, s \in (n, n+1)\}) \\
&= \iint \sum_{n=1}^{\infty} \mathbb{1}(z > xn^{1/\gamma}) \pi(dx) \lambda(dz) \\
&\leq \iint \left(\frac{z}{x}\right)^{\gamma} \mathbb{1}(z > x) \pi(dx) \lambda(dz) < \infty,
\end{aligned}$$

by the assumption. Therefore, the Borel–Cantelli lemma implies that for n large enough

$$\mu(\{(x, s, z) : z > xn^{1/\gamma}, |s| \in (n, n+1)\}) = 0,$$

from which the statement follows. $\square$

Next we handle the compensation of $X_{\pm, >, \kappa}^*$.

Lemma 12. *For any $\kappa \in (0, 1)$*

$$\iiint_{A(t; \kappa)^c} \frac{z}{x} (1 - e^{-x(t-s)}) \nu(dx, ds, dz) \leq \begin{cases} ct^{\beta-\alpha-\kappa(\beta-1)}, & \alpha \leq \beta - 1, \\ ct^{1-\kappa\alpha}, & \alpha \geq \beta - 1, \end{cases} \quad (32)$$

and

$$\nu(A(t; \kappa)^c) \leq \begin{cases} ct^{\beta-\alpha-\kappa\beta}, & \alpha \leq \beta - 1, \\ ct^{1-\kappa(1+\alpha)}, & \alpha \geq \beta - 1, \end{cases} \quad (33)$$

while for the negative jumps

$$\iiint_{B(t; \kappa)^c} \frac{z}{x} (1 - e^{-xt}) e^{xs} \nu(dx, ds, dz) \leq ct^{\beta-\alpha-\kappa(\beta-1)}, \quad (34)$$

and

$$\nu(B(t; \kappa)^c) \leq c \log t t^{\beta-\alpha-\kappa\beta}. \quad (35)$$

Proof. Let K denote the left-hand side in (32). Then

$$\begin{aligned}
K &\leq \iiint \left[zu \mathbb{1}(ux \leq 1, zu > t^\kappa) + \frac{z}{x} \mathbb{1}(ux > 1, z > t^\kappa x) \right] \nu(dx, du, dz) \\
&=: I_1 + I_2.
\end{aligned}$$

Since in I_1 necessarily $u \geq t^\kappa$, we have by (30)

$$\begin{aligned} I_1 &\leq \int_{t^\kappa}^t u \, du \int_{(0, u^{-1}]} \pi(\mathrm{d}x) \int_{(t^\kappa/u, 1]} z \lambda(\mathrm{d}z) \\ &\leq ct^{-\kappa(\beta-1)} \int_{t^\kappa}^t u^{\beta-1-\alpha} \mathrm{d}u \leq \begin{cases} ct^{\beta-\alpha-\kappa(\beta-1)}, & \alpha < \beta, \\ c \log t \, t^{\kappa(1-\alpha)}, & \alpha = \beta, \\ ct^{\kappa(1-\alpha)}, & \alpha < \beta. \end{cases} \end{aligned}$$

Similarly, for I_2 after integration with respect to s we have

$$I_2 \leq t \int_{(t^{-1}, t^{-\kappa}]} x^{-1} \pi(\mathrm{d}x) \int_{(t^\kappa x, 1]} z \lambda(\mathrm{d}z) \leq \begin{cases} ct^{\beta-\alpha-\kappa(\beta-1)}, & \alpha \leq \beta - 1, \\ ct^{1-\kappa\alpha}, & \alpha \geq \beta - 1, \end{cases}$$

and (32) follows. In exactly the same way we obtain (33). The only difference is that in both I_1 and in I_2 a factor $t^{-\kappa}$ appear.

Denoting by L the left-hand side of (34), we similarly have

$$\begin{aligned} L &\leq \iiint \left[zte^{xs} \mathbf{1}(xt \leq 1, zte^{xs} > t^\kappa) \right. \\ &\quad \left. + \frac{z}{x} e^{xs} \mathbf{1}(xt > 1, ze^{xs} > t^\kappa x) \right] \nu(\mathrm{d}x, \mathrm{d}s, \mathrm{d}z) \\ &=: J_1 + J_2. \end{aligned}$$

In both J_1 and J_2 we bound the $\mathrm{d}s$ integral by x^{-1} , thus we obtain by using (30)

$$J_1 \leq t \int_{(0, t^{-1}]} x^{-1} \pi(\mathrm{d}x) \int_{(t^{\kappa-1}, 1]} z \lambda(\mathrm{d}z) \leq ct^{\beta-\alpha-\kappa(\beta-1)},$$

and

$$J_2 \leq \int_{(t^{-1}, t^{-\kappa}]} x^{-2} \pi(\mathrm{d}x) \int_{(t^\kappa x, 1]} z \lambda(\mathrm{d}z) \leq \begin{cases} ct^{\beta-\alpha-\kappa(\beta-1)}, & \alpha \leq \beta, \\ ct^{\kappa(1-\alpha)}, & \alpha \geq \beta, \end{cases}$$

proving (34). The estimate (35) follows in the same way, except that we bound the $\mathrm{d}s$ integral in both terms by $(1 - \kappa)x^{-1} \log t$. $\square$

Lemma 13. *Assume that $1 + \alpha \leq \beta$. Then for any $\kappa > 1 - \frac{\alpha}{\beta}$ almost surely $\mu(A(t; \kappa)^c) = 0$ and $\mu(B(t; \kappa)^c) = 0$ for t large enough.*

Proof. Fix $\kappa' \in (1 - \alpha/\beta, \kappa)$. Then for some $\delta = \delta(\kappa') > 0$, by (33) in Lemma 12

$$\nu(A(t; \kappa')^c) \leq ct^{-\delta}.$$

Thus if $d\delta > 1$, by the Borel–Cantelli lemma $\mu(A(n^d, \kappa')^c) = 0$ a.s. for n large enough. Assume that $\mu(A(t; \kappa)^c) \geq 1$ for some $t \in ((n-1)^d, n^d]$. Then there exists a point $(\xi, \tau, \zeta) \in A(t; \kappa)^c$, such that, if n is large enough,

$$t^\kappa \leq \frac{\zeta}{\xi} \left(1 - e^{-\xi(t-\tau)}\right) \leq \frac{\zeta}{\xi} \left(1 - e^{-\xi(n^d-\tau)}\right) \leq n^{d\kappa'},$$

which is a contradiction. Therefore $\mu(A(t; \kappa)^c) = 0$ for t large enough, as claimed.

The same proof works for $B(t; \kappa)^c$ since the extra $\log t$ factor in (35) plays no role here. $\square$

We now combine the previous results to obtain the statement for λ supported on $[-1, 1]$.

Corollary 5. *Assume that $\int_{|z| \leq 1} |z| \lambda(dz) = \infty$, $a = 0$, $b = 0$ in (2) and λ is supported on $[-1, 1]$. Then*

$$\limsup_{t \rightarrow \infty} \frac{|X^*(t)|}{t \log t} \leq 1 \quad a.s. \quad (36)$$

Furthermore, if $\beta \leq 1 + \alpha$ and for some $\gamma \in [1, 2]$

$$\int_{(0, \infty)} \int_{|z| \leq 1} \frac{|z|^\gamma}{x^\gamma} \mathbb{1}(|z| > x) \lambda(dz) \pi(dx) < \infty, \quad (37)$$

then

$$\limsup_{t \rightarrow \infty} \frac{|X^*(t)|}{t^{1/\gamma} \log t} \leq 1 \quad a.s.$$

If $\beta \geq 1 + \alpha$, then for any $\gamma < 1/(1 - \frac{\alpha}{\beta})$

$$\lim_{t \rightarrow \infty} \frac{X^*(t)}{t^{1/\gamma}} = 0 \quad a.s.$$

Proof. Assume first that λ is supported on $(0, 1]$. Then (36) follows from Corollary 4 with $\kappa = 1$. Recall that $\kappa = 1$ corresponds to no truncation, thus $X_{\pm, <, 1}^*(t) = X_\pm^*(t)$.

Next assume (37). Then $1 + \alpha \geq \gamma$, and by assumption $\beta \leq 1 + \alpha$. Lemma 11 implies that the terms $X_{\pm, >, 1/\gamma}^*(t)$ do not contribute to the limit. The result now follows from Corollary 4 and Lemma 12 by taking $\kappa = 1/\gamma$.

For $\beta \geq 1 + \alpha$ the result follows from Corollary 4 and Lemma 12 with $\kappa = 1 - \frac{\alpha}{\beta}$, and from Lemma 13.

For λ supported on $[-1, 1]$, we apply the previous to $\mu\mathbf{1}(z > 0)$ and $\mu\mathbf{1}(z < 0)$ separately. $\square$

Proof of Theorem 2. If we make a decomposition based on $\mu\mathbf{1}(|z| \leq 1)$ and $\mu\mathbf{1}(|z| > 1)$, we obtain the results by applying Corollary 5 and Theorem 1, respectively. $\square$

6 Proof of Theorem 4

We will use a general result for Gaussian processes from [38]. Since X is Gaussian, we have that

$$\mathbf{E}e^{i\theta\Lambda(A)} = e^{-\frac{b}{2}\theta^2(\pi \times \text{Leb}(A))}$$

and by Proposition 2.6. in [42]

$$\log \mathbf{E}e^{i\theta X^*(t)} = -\frac{b}{2}\theta^2 \int_{(0,\infty)} \int_0^t (1 - e^{-xu}) x^{-2} du \pi(dx).$$

Thus X^* is a mean 0 Gaussian process with

$$Q(t) := \frac{1}{2} \mathbf{Var}(X^*(t)) = \frac{b}{2} \int_{(0,\infty)} \int_0^t (1 - e^{-xu}) x^{-2} du \pi(dx).$$

Hence, Q is non-decreasing and $Q(0) = 0$.

1. If $m_{-2}(\pi) < \infty$, then

$$Q(t) = \frac{b}{2}t \int_{(0,\infty)} \frac{1 + tx - e^{-tx}}{tx} x^{-2} \pi(dx) \sim \frac{b}{2}tm_{-2}(\pi),$$

as $t \rightarrow \infty$. Now the result follows by Theorem 1.1 in [38] with $v(t) = \frac{b}{2}tm_{-2}(\pi)$ (see also Remark 1.3 in [38]).

2. As in [20, Eq. (5.8)], we can write $Q(t)$ in the form

$$Q(t) = \frac{b}{2} \frac{\Gamma(1 + \alpha)}{(2 - \alpha)(1 - \alpha)} \ell_1(t) t^{2-\alpha},$$

with ℓ_1 slowly varying at infinity such that $\ell_1(t) \sim \ell(t)$. By Potter's bounds [7, Theorem 1.5.6], for $\delta < \alpha/3$, there is $s_0 > 0$ such that

$$\left(\frac{t}{s}\right)^{2-\alpha-\delta} Q(s) \leq Q(t) \leq \left(\frac{t}{s}\right)^{2-\alpha+\delta} Q(s), \quad t \geq s > s_0.$$

Moreover,

$$Q(t) \leq \frac{b}{2} \int_{(0,\infty)} \int_0^t u dx x^{-1} \pi(dx) = \frac{b}{4} m_{-1}(\pi) t^2.$$

Thus, the assumptions of Theorem 1.1 in [38] hold with $v = Q$.

Appendix A: A Fubini theorem

Here we show that the stochastic Fubini theorem from [2] is indeed applicable in our setup. If π is a finite measure, this was justified in [22, Lemma 4.1] (see also [46, Lemma 3.1] for a similar argument).

Put $g_t(x, s) = e^{-x(t-s)} \mathbf{1}(s \leq t)$. If $\mathbf{E}|X(1)| < \infty$, then we can directly use a stochastic Fubini theorem given in [2, Theorem 3.1] which guarantees the integrals exist. Indeed, the conditions of Theorem 3.1 and Remark 3.2 in [2] boil down to showing that

(i) for every $u \in [0, t]$, $g_u(\cdot, \cdot)$ is in the Musielak-Orlicz space L^{ϕ_1} , that is

$$\iint_{(0,\infty) \times \mathbb{R}} \left(b g_u(x, s)^2 + \int_{\mathbb{R}} (|z g_u(x, s)|^2 \wedge |z g_u(x, s)|) \lambda(dz) \right) \pi(dx) ds < \infty,$$

(ii) it holds that

$$\begin{aligned} \int_0^t \left[\iint_{(0,\infty) \times \mathbb{R}} \left(b g_u(x, s)^2 \right. \right. \\ \left. \left. + \int_{\mathbb{R}} (|z g_u(x, s)|^2 \wedge |z g_u(x, s)|) \lambda(dz) \right) \pi(dx) ds \right] du < \infty. \end{aligned}$$

By [42, Theorem 3.3], L^{ϕ_1} coincides with the space of Λ -integrable functions g such that $\mathbf{E}|\int g d\Lambda| < \infty$. The supOU process is well defined, i.e. $g_t(\cdot, \cdot)$ is Λ -integrable and since we have assumed $\mathbf{E}|X(u)| < \infty$, we conclude that condition (i) holds. By the change of variables, first $r = s - u$

and then $r = t - s$

$$\begin{aligned}
& \int_0^t \int_{(0,\infty)} \int_{\mathbb{R}} \left(be^{-2x(u-s)} \mathbf{1}(s \leq u) \right. \\
& \quad \left. + \int_{\mathbb{R}} (|ze^{-x(u-s)}|^2 \wedge |ze^{-x(u-s)}|) \mathbf{1}(s \leq u) \lambda(dz) \right) \pi(dx) ds du \\
&= \int_0^t \int_{(0,\infty)} \int_{\mathbb{R}} \left(be^{2xr} \mathbf{1}(r \leq 0) \right. \\
& \quad \left. + \int_{\mathbb{R}} (|ze^{xr}|^2 \wedge |ze^{xr}|) \mathbf{1}(r \leq 0) \lambda(dz) \right) \pi(dx) dr du \\
&= t \int_{(0,\infty)} \int_{\mathbb{R}} \left(be^{2xr} \mathbf{1}(r \leq 0) + \int_{\mathbb{R}} (|ze^{xr}|^2 \wedge |ze^{xr}|) \mathbf{1}(r \leq 0) \lambda(dz) \right) \pi(dx) dr \\
&= t \int_{(0,\infty)} \int_{\mathbb{R}} \left(be^{-2x(t-s)} \mathbf{1}(s \leq t) \right. \\
& \quad \left. + \int_{\mathbb{R}} (|xe^{-x(t-s)}|^2 \wedge |xe^{-x(t-s)}|) \mathbf{1}(s \leq t) \lambda(dz) \right) \pi(dx) ds \\
&= \int_{(0,\infty)} \int_{\mathbb{R}} \left(bg_t(x, s)^2 + \int_{\mathbb{R}} (|xg_t(x, s)|^2 \wedge |xg_t(x, s)|) \lambda(dz) \right) \pi(dx) ds < \infty,
\end{aligned}$$

hence, (ii) follows from (i).

Suppose now that $\mathbf{E}|X(1)| = \infty$. By the Lévy-Itô decomposition of Λ , we may write it in the form

$$\Lambda(A) = \Lambda_1(A) + \Lambda_2(A),$$

where Λ_1 and Λ_2 are independent random measures such that Λ_1 has the characteristic quadruple $(0, 0, \lambda_1, \pi)$, $\lambda_1(dz) = \lambda(dz) \mathbf{1}(|z| > 1)$ and Λ_2 has characteristic quadruple (a, b, λ_2, π) , $\lambda_2(dz) = \lambda(dz) \mathbf{1}(|z| \leq 1)$. By [43, Proposition 3.3.9] we have that

$$\begin{aligned}
& \int_{(0,\infty) \times \mathbb{R}} g_u(x, s) \Lambda(dx, ds) \\
&= \int_{(0,\infty) \times \mathbb{R}} g_u(x, s) \Lambda_1(dx, ds) + \int_{(0,\infty) \times \mathbb{R}} g_u(x, s) \Lambda_2(dx, ds), \text{ a.s.}
\end{aligned}$$

For the integral with respect to Λ_2 we can apply [2, Theorem 3.1] as in the previous case. It remains to consider Λ_1 , which is a compound Poisson random measure. If $(\xi_k, \tau_k, \zeta_k)_{k \geq 0}$ are the points of Λ_1 , then the corresponding supOU process $X_1(u)$ can be written as

$$X_1(u) = \sum_{\tau_k \leq 0} \zeta_k e^{-\xi_k(u-\tau_k)} + \sum_{0 < \tau_k \leq u} \zeta_k e^{-\xi_k(u-\tau_k)}.$$

The second sum has finitely many terms a.s. and for the first one we have

$$\int_0^t \sum_{\tau_k \leq 0} |\zeta_k| e^{-\xi_k(u-\tau_k)} du \leq t \sum_{\tau_k \leq 0} |\zeta_k| e^{\xi_k \tau_k},$$

which is finite a.s. since

$$\iiint_{(0,\infty) \times (-\infty,0) \times \mathbb{R}} (|z|e^{xs} \wedge 1) \pi(dx) ds \mu_1(dz) < \infty.$$

Appendix B: An exotic example

Here we give an example for π and λ such that $\beta > 1 + \alpha$ and

$$\iint_{(0,\infty) \times \mathbb{R}} \frac{|z|^\gamma}{x^\gamma} \mathbf{1}(|z| > x) \pi(dx) \lambda(dz) < \infty,$$

still holds with $\gamma > 0$. Let $a_n = 2^{-2^n}$ and $a \geq 0$. For $r \in (a_n, a_{n-1}]$ put $\pi((0, r]) = a_n^{1+a}$. Then $\pi(\{a_n\}) = a_n^{1+a} - a_{n+1}^{1+a} \sim a_n^{1+a}$, thus

$$\int_{(0,1]} x^{-1-\alpha} \pi(dx) = \sum_{n=1}^{\infty} a_n^{-1-\alpha} \pi(\{a_n\}) < \infty$$

if and only if $a > \alpha$. In particular, $\alpha_0 = a$ and $\alpha < a$.

Let $b > 0$ and for $r \in (a_n, a_{n-1}]$ let $\bar{\lambda}(r) = \lambda((r, 1]) = a_{n-1}^{-b}$. Then $\lambda(\{a_n\}) = a_n^{-b} - a_{n-1}^{-b} \sim a_n^{-b}$, therefore,

$$\int_{(0,1]} z^\beta \lambda(dz) = \sum_{n=1}^{\infty} a_n^\beta \pi(\{a_n\}) < \infty$$

if and only if $b < \beta$. Thus $\beta_0 = b$ and $\beta > b$. Furthermore, for $\gamma < b$, $r \in (0, 1)$

$$\int_{(r,1]} z^\gamma \lambda(dz) \sim \sum_{a_n > r} a_n^{\gamma-b} \sim a_{\psi(r)}^{\gamma-b},$$

where $\psi(r) = \max\{k : a_k > r\}$.

Thus

$$\iint \frac{z^\gamma}{x^\gamma} \mathbf{1}(z > x) \lambda(dz) \pi(dx) \sim \sum_n a_n^{1+a-\gamma} a_{\psi(a_n)}^{\gamma-b} = \sum_n a_{n-1}^{2(1+a-\gamma)+\gamma-b},$$

where we used that $\psi(a_n) = n - 1$ and $a_n = a_{n-1}^2$. The latter sum is finite if and only if $2(1 + a - \gamma) + \gamma - b > 0$, which is equivalent to $\gamma < 2(1 + a) - b$. This is possible even if $b > 1 + a$.

Acknowledgement. We thank the referees and the AE for constructive remarks and suggestions which greatly improved our paper. DG was partially supported by the Croatian Science Foundation (HRZZ) grant Scaling in Stochastic Models (IP-2022-10-8081). PK was supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences, and by the NKFIH grant FK124141.

References

- [1] O. E. Barndorff-Nielsen. Superposition of Ornstein-Uhlenbeck type processes. *Teor. Veroyatnost. i Primenen.*, 45(2):289–311, 2000.
- [2] O. E. Barndorff-Nielsen and A. Basse-O'Connor. Quasi Ornstein-Uhlenbeck processes. *Bernoulli*, 17(3):916–941, 2011.
- [3] O. E. Barndorff-Nielsen, F. E. Benth, and A. E. D. Veraart. *Ambit Stochastics*, volume 88 of *Probability Theory and Stochastic Modelling*. Springer, Cham, 2018.
- [4] O. E. Barndorff-Nielsen and R. Stelzer. Multivariate supOU processes. *Ann. Appl. Probab.*, 21(1):140–182, 2011.
- [5] O. E. Barndorff-Nielsen and R. Stelzer. The multivariate supOU stochastic volatility model. *Math. Finance*, 23(2):275–296, 2013.
- [6] J. Bertoin. *Lévy Processes*, volume 121 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1996.
- [7] N. H. Bingham, C. M. Goldie, and J. L. Teugels. *Regular Variation*, volume 27 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1989.
- [8] R. A. Carmona and S. A. Molchanov. Parabolic Anderson problem and intermittency. *Mem. Amer. Math. Soc.*, 108(518):viii+125, 1994.
- [9] C. Chong and P. Kevei. Intermittency for the stochastic heat equation with Lévy noise. *Ann. Probab.*, 47(4):1911–1948, 2019.

- [10] C. Chong and P. Kevei. The almost-sure asymptotic behavior of the solution to the stochastic heat equation with Lévy noise. *Ann. Probab.*, 48(3):1466–1494, 2020.
- [11] I. V. Curato and R. Stelzer. Weak dependence and GMM estimation of supOU and mixed moving average processes. *Electron. J. Stat.*, 13(1):310–360, 2019.
- [12] P. Embrechts, C. Klüppelberg, and T. Mikosch. *Modelling Extremal Events*, volume 33 of *Applications of Mathematics (New York)*. Springer-Verlag, Berlin, 1997. For insurance and finance.
- [13] V. Fasen. Extremes of regularly varying Lévy-driven mixed moving average processes. *Adv. in Appl. Probab.*, 37(4):993–1014, 2005.
- [14] V. Fasen and C. Klüppelberg. Extremes of supOU processes. In *Stochastic analysis and applications*, volume 2 of *Abel Symp.*, pages 339–359. Springer, Berlin, 2007.
- [15] F. Fuchs and R. Stelzer. Mixing conditions for multivariate infinitely divisible processes with an application to mixed moving averages and the supOU stochastic volatility model. *ESAIM Probab. Stat.*, 17:455–471, 2013.
- [16] J. Gärtner, W. König, and S. Molchanov. Geometric characterization of intermittency in the parabolic Anderson model. *Ann. Probab.*, 35(2):439–499, 2007.
- [17] D. Grahovac and P. Kevei. Tail behavior and almost sure growth rate of supOU processes. *arXiv:2402.08584*, 2024.
- [18] D. Grahovac, N. N. Leonenko, A. Sikorskii, and M. S. Taqqu. The unusual properties of aggregated superpositions of Ornstein-Uhlenbeck type processes. *Bernoulli*, 25(3):2029–2050, 2019.
- [19] D. Grahovac, N. N. Leonenko, and M. S. Taqqu. Intermittency of trawl processes. *Statist. Probab. Lett.*, 137:235–242, 2018.
- [20] D. Grahovac, N. N. Leonenko, and M. S. Taqqu. Limit theorems, scaling of moments and intermittency for integrated finite variance supOU processes. *Stochastic Process. Appl.*, 129(12):5113–5150, 2019.
- [21] D. Grahovac, N. N. Leonenko, and M. S. Taqqu. The multifaceted behavior of integrated supOU processes: the infinite variance case. *J. Theoret. Probab.*, 33(4):1801–1831, 2020.

- [22] D. Grahovac, N. N. Leonenko, and M. S. Taqqu. Intermittency and infinite variance: the case of integrated supOU processes. *Electron. J. Probab.*, 26:Paper No. 56, 31, 2021.
- [23] D. Grahovac, N. N. Leonenko, and M. S. Taqqu. Intermittency and multiscaling in limit theorems. *Fractals*, 30(07):2250137, 2022.
- [24] C. W. J. Granger. Long memory relationships and the aggregation of dynamic models. *J. Econometrics*, 14(2):227–238, 1980.
- [25] A. Gut. *Probability: A Graduate Course*. Springer Texts in Statistics. Springer, New York, second edition, 2013.
- [26] B. C. Kelly, T. Treu, M. Malkan, A. Pancoast, and J.-H. Woo. Active galactic nucleus black hole mass estimates in the era of time domain astronomy. *The Astrophysical Journal*, 779:187, 2013.
- [27] D. Khoshnevisan. A primer on stochastic partial differential equations. In *A minicourse on stochastic partial differential equations*, volume 1962 of *Lecture Notes in Math.*, pages 1–38. Springer, Berlin, 2009.
- [28] D. Khoshnevisan. *Analysis of Stochastic Partial Differential Equations*, volume 119 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2014.
- [29] D. Khoshnevisan, K. Kim, and C. Mueller. Dissipation in parabolic SPDEs II: Oscillation and decay of the solution. *Ann. Inst. Henri Poincaré Probab. Stat.*, 59(3):1610–1641, 2023.
- [30] D. Khoshnevisan, K. Kim, C. Mueller, and S.-Y. Shiu. Dissipation in parabolic SPDEs. *J. Stat. Phys.*, 179(2):502–534, 2020.
- [31] D. Khoshnevisan, K. Kim, and Y. Xiao. Intermittency and multifractality: a case study via parabolic stochastic PDEs. *Ann. Probab.*, 45(6A):3697–3751, 2017.
- [32] N. Kôno. Iterated log type strong limit theorems for self-similar processes. *Proc. Japan Acad. Ser. A Math. Sci.*, 59(3):85–87, 1983.
- [33] A. E. Kyprianou. *Fluctuations of Lévy Processes with Applications*. Universitext. Springer, Heidelberg, second edition, 2014. Introductory lectures.

- [34] R. Leipus and D. Surgailis. Random coefficient autoregression, regime switching and long memory. *Adv. in Appl. Probab.*, 35(3):737–754, 2003.
- [35] C. Marinelli and M. Röckner. On maximal inequalities for purely discontinuous martingales in infinite dimensions. In *Séminaire de Probabilités XLVI*, volume 2123 of *Lecture Notes in Math.*, pages 293–315. Springer, Cham, 2014.
- [36] M. Moser and R. Stelzer. Tail behavior of multivariate Lévy-driven mixed moving average processes and supOU stochastic volatility models. *Adv. in Appl. Probab.*, 43(4):1109–1135, 2011.
- [37] G. Oppenheim and M.-C. Viano. Aggregation of random parameters Ornstein-Uhlenbeck or AR processes: some convergence results. *J. Time Ser. Anal.*, 25(3):335–350, 2004.
- [38] S. Orey. Growth rate of certain Gaussian processes. In *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability (Univ. California, Berkeley, Calif., 1970/1971), Vol. II: Probability theory*, pages 443–451. Univ. California Press, Berkeley, CA, 1972.
- [39] M. S. Pakkanen, R. Passeggeri, O. Sauri, and A. E. D. Veraart. Limit theorems for trawl processes. *Electron. J. Probab.*, 26:Paper No. 116, 36, 2021.
- [40] J. Pedersen. The Lévy-Itô decomposition of an independently scattered random measure. *MaPhySto Research Report, University of Aarhus*, 2003. <http://www.maphysto.dk>.
- [41] A. Philippe, D. Puplinskaite, and D. Surgailis. Contemporaneous aggregation of triangular array of random-coefficient AR(1) processes. *J. Time Series Anal.*, 35(1):16–39, 2014.
- [42] B. S. Rajput and J. Rosiński. Spectral representations of infinitely divisible processes. *Probab. Theory Related Fields*, 82(3):451–487, 1989.
- [43] G. Samorodnitsky. *Stochastic processes and long range dependence*. Springer Series in Operations Research and Financial Engineering. Springer, Cham, 2016.
- [44] K. Sato. *Lévy Processes and Infinitely Divisible Distributions*, volume 68 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1999. Translated from the 1990 Japanese original, Revised by the author.

- [45] K. Takashima. Sample path properties of ergodic self-similar processes. *Osaka J. Math.*, 26(1):159–189, 1989.
- [46] A. Talarczyk and Ł. Tszczotko. Limit theorems for integrated trawl processes with symmetric Lévy bases. *Electron. J. Probab.*, 25:Paper No. 117, 24, 2020.
- [47] J. Tiefeng, M. B. Rao, W. Xiangchen, and L. Deli. Laws of large numbers and moderate deviations for stochastic processes with stationary and independent increments. *Stochastic Process. Appl.*, 44(2):205–219, 1993.
- [48] H. Watanabe. An asymptotic property of Gaussian processes. I. *Trans. Amer. Math. Soc.*, 148:233–248, 1970.
- [49] Y. B. Zel’dovich, S. A. Molchanov, A. A. Ruzmaïkin, and D. D. Sokolov. Intermittency in random media. *Uspekhi Fiz. Nauk*, 152(1):3–32, 1987.